

Central Lancashire Online Knowledge (CLoK)

Title	Reduced CD4 T Lymphocytes in Lymph Nodes of the Mouse Model of Autism	
	Induced by Valproic Acid	
Type	Article	
URL	https://clok.uclan.ac.uk/id/eprint/24078/	
DOI	https://doi.org/10.1159/000491395	
Date	2018	
Citation	Baronio, Diego, Bauer-Negrini, Guilherme, Castro, Kamila, Della-Flora Nunes, Gustavo, Riesgo, Rudimar, Mendes-da-Cruz, Daniella Arêas, Savino, Wilson, Gottfried, Carmem and Bambini-Junior, Victorio (2018) Reduced CD4 T Lymphocytes in Lymph Nodes of the Mouse Model of Autism Induced by Valproic Acid. Neuroimmunomodulation. ISSN 1021-7401	
Creators	Baronio, Diego, Bauer-Negrini, Guilherme, Castro, Kamila, Della-Flora Nunes, Gustavo, Riesgo, Rudimar, Mendes-da-Cruz, Daniella Arêas, Savino, Wilson, Gottfried, Carmem and Bambini-Junior, Victorio	

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1159/000491395

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the http://clok.uclan.ac.uk/policies/

Table 1. Total T cell counts in the thymuses and spleen from animals of the VPA model of autism and controls*

Number of thymocytes (x 10 ⁶)			
	Control $(n = 9)$	VPA (n = 9)	
CD4-CD8-	8.96 ± 5.76	6.87 ± 2.73	
CD4+CD8+	55.51 ± 9.84	63.32 ± 10.97	
CD4+CD8-	6.12 ± 1.21	5.77 ± 1.44	
CD4-CD8+	2.63 ± 0.77	2.40 ± 0.94	
Number of spleen T lymphocytes (x 10 ⁶)			
CD3+CD4+	5.86 ± 1.11	4.94 ± 1.14	
CD3+CD8+	1.93 ± 0.37	1.37 ± 0.33	

^{*}No statistically significant differences were found when comparing control versus VPA

T-cell subsets in the thymus or spleen