

Central Lancashire Online Knowledge (CLOK)

Title	Identification of strata for a trial of a targeted multimodal physiotherapy intervention in patellofemoral pain patients
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/25190/
DOI	https://doi.org/10.1186/1745-6215-16-S2-P158
Date	2015
Citation	Sutton, Chris J, Dey, Paola, Janssen, Jessica, Richards, James, Hogarth, Russell and Selfe, James (2015) Identification of strata for a trial of a targeted multimodal physiotherapy intervention in patellofemoral pain patients. <i>Trials</i> , 16 (S2). p. 158.
Creators	Sutton, Chris J, Dey, Paola, Janssen, Jessica, Richards, James, Hogarth, Russell and Selfe, James

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1186/1745-6215-16-S2-P158>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

POSTER PRESENTATION

Open Access

Identification of strata for a trial of a targeted multimodal physiotherapy intervention in patellofemoral pain patients

Chris Sutton^{1*}, Paola Dey¹, Jessie Janssen¹, Michael Callaghan², Jim Richards¹, Maria Stokes³, Denis Martin⁴, John Dixon⁴, Erik Witvrouw^{5,6}, Russell Hogarth¹, Vasilios Baltzopoulos⁷, Elizabeth Ritchie⁸, Nigel Arden⁹, Rich Masters¹⁰, Remco Polman¹¹, David Turner¹², James Selfe¹

From 3rd International Clinical Trials Methodology Conference
Glasgow, UK. 16-17 November 2015

Background

Patellofemoral Pain (PFP) is a musculoskeletal disorder causing significant pain and dysfunction around the knee, commonly leading to long term limitations. People with PFP are commonly referred for physiotherapy, although current multimodal physiotherapy approaches tend to be rather ad-hoc and are failing in the long term. Identification of patient strata (or subgroups) who may respond differentially to interventions has been recognised as an international priority. It has been proposed that there are PFP patient strata, classified based on clinical tests, who would respond to modes of interventions targeted at the individual's stratum.

Methods

Participants with PFP underwent clinical assessment of muscle weakness, muscle length, patellar mobility and foot posture. The presence of strata was explored using two classification techniques: hierarchical cluster analysis and latent profile analysis (LPA); for LPA, the Bayesian Information Criterion was used to guide model selection.

Results

One hundred and twenty seven (of 130) recruited participants had complete assessment data. A three strata solution appeared optimal from modelling and clinical perspectives. The three suggested strata were characterised as: stronger; weaker and tighter; weaker with pronated feet. However, although membership of the stronger stratum was consistent, there were substantial

differences in the other composition of the other two strata between the two classification approaches.

Conclusion

PFP patients can be classified into clinical strata, although further investigation is needed to identify thresholds upon which stratification should be performed. Interventions are currently being mapped onto the strata and feasibility and evaluation trials of these interventions should then be performed.

Authors' details

¹University of Central Lancashire, Preston, Lancashire, UK. ²University of Manchester, Manchester, UK. ³University of Southampton, Southampton, Hampshire, UK. ⁴Teesside University, Middlesbrough, Teesside, UK. ⁵Ghent University, Ghent, Belgium. ⁶Aspetar, Doha, Qatar. ⁷Brunel University, London, UK. ⁸Harrogate and District National Health Services, Harrogate, N Yorkshire, UK. ⁹University of Oxford, Oxford, Oxfordshire, UK. ¹⁰University of Waikato, Hamilton, New Zealand. ¹¹Bournemouth University, Bournemouth, Dorset, UK. ¹²University of East Anglia, Norwich, Norfolk, UK.

Published: 16 November 2015

doi:10.1186/1745-6215-16-S2-P158

Cite this article as: Sutton *et al.*: Identification of strata for a trial of a targeted multimodal physiotherapy intervention in patellofemoral pain patients. *Trials* 2015 **16**(Suppl 2):P158.

¹University of Central Lancashire, Preston, Lancashire, UK
Full list of author information is available at the end of the article