

Central Lancashire Online Knowledge (CLoK)

Title	Morphine self-administration alters the expression of translational machinery genes in the amygdala of male Lewis rats
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/27967/
DOI	https://doi.org/10.1177/0269881119836206
Date	2019
Citation	Ucha, Marcos, Coria, Santiago M, Núñez, Adrián E, Santos, Raquel, Roura-Martínez, David, Fernández-Ruiz, Javier, Higuera-Matas, Alejandro and Ambrosio, Emilio (2019) Morphine self-administration alters the expression of translational machinery genes in the amygdala of male Lewis rats. <i>Journal of Psychopharmacology</i> , 33 (7). pp. 882-893. ISSN 0269-8811
Creators	Ucha, Marcos, Coria, Santiago M, Núñez, Adrián E, Santos, Raquel, Roura-Martínez, David, Fernández-Ruiz, Javier, Higuera-Matas, Alejandro and Ambrosio, Emilio

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1177/0269881119836206>


For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Supplementary Information:

Alternative analyses of the delay-discounting task with area under the curve (AUC) and normalized AUC

There is a debate in the preclinical literature on impulsivity about the best way to analyze impulsivity using delay-discounting curves (the two debated options are curve-fitting approaches or to compute the area under the delay-discounting curve). In addition to extracting the k parameter (see Methods section) we have also computed the AUC (normalized and non-normalized). When using these AUCs as input variables, the clustering approach was inadequate as it created two groups with a very unequal number of animals, even after removing two outliers (see SI Fig.1). In spite of this, the correlation obtained for *Cnr1* and the k parameter had an equivalent correlation between both the AUC (Magnard et al., 2018) and the normalized AUC (Myerson et al., 2001), with or without two identified outliers, and *Cnr1*, supporting the robustness of our results.



SI. Fig 1. The relationship between *Cnr1* gene expression and impulsive behavior measured (alternative analysis)

Impulsive choice was positively correlated with impulsivity measured by both measures of AUC, with or without outliers (a smaller AUC indicates a faster switch in the preference in favor of the immediate lever i.e. higher impulsivity) A) Impulsive choice (as defined by the normalized AUC, without removing outliers) was positively correlated with *Cnr1* (CB₁ cannabinoid receptor) gene expression in the mOFC (Kendall's $\tau=-0.584$; $p=0.02$). B) Impulsive choice (as defined by the AUC, without removing the outliers) was positively

correlated with *Cnr1* (CB₁ cannabinoid receptor) gene expression the mOFC (Kendall's τ -0.629=; p=0.012). C) Impulsive choice (as defined by the normalized AUC without the two outliers) was positively correlated with *Cnr1* (CB₁ cannabinoid receptor) gene expression the mOFC (Pearson's r=-0.747; p=0.033). D) Impulsive choice (as defined by the AUC without the outliers) was positively correlated with *Cnr1* (CB₁ cannabinoid receptor) gene expression in the mOFC (Pearson's r=-0.740; p=0.036).

SI Table 1. Primer sets used for RT-qPCR amplification.

Gene	Description	Forward primer	Reverse primer
<i>Grin1</i>	glutamate ionotropic receptor NMDA type subunit 1	AACCTGCAGAACCGCAAG	GCTTGATGAGCAGGTCTATGC
<i>Grin2a</i>	glutamate ionotropic receptor NMDA type subunit 2A	TGTGAAGAAATGCTGCAAGG	GAACGCTCCTCATTGATGGT
<i>Gria1</i>	glutamate ionotropic receptor AMPA type subunit 1	AGAGGCTGGTGGTGGTTGACT	ACCCTGGTATGGTCTCGGGA
<i>Gria2</i>	glutamate ionotropic receptor AMPA type subunit 2	GGCGTGTAATCCTGGACTGT	ACACCAGGGAATCGTCGTAG
<i>Gabrg2</i>	gamma-aminobutyric acid type A receptor gamma 2 subunit	CGGAAACCAAGCAAGGATAA	ACAGTCCTTGCCATCCAAAC
<i>Gabrd</i>	gamma-aminobutyric acid type A receptor delta subunit	GCTGGACCTGGAGAGCTATG	CCGAAGCTGGAAGGTAAAGC
<i>Gabra1</i>	gamma-aminobutyric acid type A receptor alpha 1 subunit	TTGACTGTGAGAGCCGAATG	AAACGTGACCCATCTTCTGC
<i>Gabra2</i>	gamma-aminobutyric acid type A receptor alpha 2 subunit	CCATGCACTTGGAGGACTT	ACTGGCCCAGCAAATCATAC
<i>Cnr1</i>	cannabinoid receptor 1	GTCGATCCTAGATGGCCTTGC	GTCATTGAGCCACGTCAGAG
<i>Dagla</i>	diacylglycerol lipase, alpha	CTTTGCTGAATTTTCCGTGACC	TTGTTTGCCTCATCCAGCAC
<i>Mgll</i>	monoacylglycerol lipase	CTACCTGCTCATGGAATC	GACACCCACGTATTTATTTTC
<i>Napepld</i>	N-acyl phosphatidylethanolamine phospholipase D	AGATATGGACTCAAGAGTGAAGACTTC	TCCTCAAAGGCTTTGTCATCGG
<i>Faah</i>	fatty acid amide hydrolase	GTTACAGAGTGGAGAGCTGTCC	GTCTCACAGTCGGTCAGATAGG
<i>Gapdh</i>	glyceraldehyde-3-phosphate dehydrogenase	TCCCTGTTCTAGAGACAG	CCACTTTGTCACAAGAGA