

Central Lancashire Online Knowledge (CLoK)

Title	Adherence and persistence to direct oral anticoagulants in atrial fibrillation: a population-based study
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/29866/
DOI	https://doi.org/10.1136/heartjnl-2019-315307
Date	2019
Citation	Banerjee, Amitava, Benedetto, Valerio, Gichuru, Phillip, Burnell, Jane, Antoniou, S, Schilling, R, Strain, WD, Ryan, R, Watkins, Caroline Leigh et al (2019) Adherence and persistence to direct oral anticoagulants in atrial fibrillation: a population-based study. Heart. ISSN 1355-6037
Creators	Banerjee, Amitava, Benedetto, Valerio, Gichuru, Phillip, Burnell, Jane, Antoniou, S, Schilling, R, Strain, WD, Ryan, R, Watkins, Caroline Leigh, Marshall, T and Sutton, Chris J

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1136/heartjnl-2019-315307>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

TITLE PAGE

Title: Adherence and persistence to direct oral anticoagulants in atrial fibrillation: a population-based study.

Authors: Amitava Banerjee^{1, 2}, Valerio Benedetto³, Phillip Gichuru⁴, Jane Burnell³, Sotiris Antoniou⁵, Richard Schilling⁵, W David Strain⁶, Ronan Ryan⁷, Caroline Watkins³, Tom Marshall⁷, Chris Sutton⁸

¹Institute of Health Informatics, University College London

²School of Health Sciences, University of Central Lancashire

³Faculty of Health and Wellbeing, University of Central Lancashire

⁴Edge Hill University

⁵Department of Cardiology, St Bartholomews Hospital, Barts NHS Trust

⁵University of Exeter

⁶Department of Primary Care, University of Birmingham

⁷ School of Health Sciences, University of Manchester

Corresponding author: Dr Amitava Banerjee, Institute of Health Informatics, University College London
222 Euston Road, London,
United Kingdom
NW1 2DA

Corresponding author's email address: ami.banerjee@ucl.ac.uk

Contributorship Statement: The study was conceived by AB. VB, PG, JB, AB and CS wrote the statistical analysis plan, carried out the analysis, collected data and produced the initial draft of the manuscript. AB was guarantor. All authors contributed to revision of the manuscript and have accepted the final version.

Transparency Statement: The corresponding author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; no important aspects of the study have been omitted.

Data Sharing Statement: No data is available for this study due to ethical constraints.

Funding Statement: The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement no. 339239. CS and PG were based at UCLan for part of this work.

WDS was supported by the NIHR Exeter Clinical Research Facility and the NIHR Collaboration for Leadership in Applied Health Research and Care (CLAHRC) for the South West Peninsula. CW and VB were supported by the NIHR CLAHRC North West Coast. TM is supported by the NIHR CLAHRC West Midlands. The views expressed in this publication are those of the authors and not necessarily those of the respective funders.

Ethical approval: Research carried out using The Health Improvement Network data was approved by the NHS South-East Multicentre Research Ethics Committee (MREC) in 2003, subject to independent scientific approval. Approval for this analysis was obtained in 2015 (SRC reference number 15THIN002).

Permission statement: The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in HEART editions and any other BMJPGL products to exploit all subsidiary rights.

Abstract

Background: Despite simpler regimes than vitamin K antagonists(VKA) for stroke prevention in atrial fibrillation(AF), adherence(taking drugs as prescribed) and persistence(continuation of drugs) to direct oral anticoagulants(DOACs) are suboptimal, yet understudied in electronic health records(EHR).

Objective: We investigated: (i)time trends at individual and system levels; (ii)risk factors for and associations between adherence and persistence.

Methods: In UK primary care EHR(The Health Information Network; 2011-2016), we investigated adherence and persistence at 1-year for OACs in adults with incident AF. Baseline characteristics were analysed by OAC and adherence/persistence status. Risk factors for non-adherence and non-persistence were assessed using Cox and logistic regression. Patterns of adherence and persistence were analysed.

Results: Among 36652 individuals with incident AF, cardiovascular comorbidities(median CHA₂DS₂VASc 3) and polypharmacy(median number of drugs 6) were common. Adherence was 55.2%(95 CI 54.6-55.7), 51.2%(50.6-51.8), 66.5%(63.7-69.2), 63.1%(61.8-64.4) and 64.7(63.2-66.1) for all OACs, VKA, dabigatran, rivaroxaban and apixaban. One-year persistence was 65.9%(65.4-66.5), 63.4%(62.8-64.0), 61.4%(58.3-64.2), 72.3%(70.9-73.7) and 78.7%(77.1-80.1) for all OACs, VKA, dabigatran, rivaroxaban and apixaban. Risk of non-adherence and non-persistence increased over time at individual and system levels. Increasing comorbidity was associated with reduced risk of non-adherence and non-persistence, across all OACs. Overall rates of “primary non-adherence”(stopping after first prescription), “non-adherent non-persistence” and “persistent adherence” were 3.5%, 26.5% and 40.2%, differing across OACs.

Conclusions: Adherence and persistence to OACs are low at one year with heterogeneity across drugs and over time, at individual and system levels. Better understanding of contributory factors will inform interventions to improve adherence and persistence across OACs in individuals and populations.

Key words: atrial fibrillation, anticoagulant, adherence, persistence, stroke

Key questions

What is already known?

- Despite proven efficacy for stroke prevention in atrial fibrillation, adherence and persistence are suboptimal for oral anticoagulants
- Adherence and persistence are rarely studied together in the same population across all anticoagulants, taking into account all baseline factors in electronic health records.

What does this study add?

- This is the first study evaluating the time trends, predictive factors and associations between adherence and persistence of anticoagulants in atrial fibrillation in a population-based study in electronic health records
- Persistence and adherence to OACs are relatively low at 1-year and there is heterogeneity across different OACs.
- There are significant variations over the time that a patient is on a DOAC, and population-level time trends in adherence and persistence after new drugs (DOACs in this case) are introduced.

How might this impact on clinical practice?

- Interventions should focus on improving adherence and persistence together and across drugs.
- These data may help to better understand the determinants of adherence and persistence, and to design and target interventions

Adherence and persistence to direct oral anticoagulants in atrial fibrillation: a population-based study.

Introduction

For sixty years, vitamin K antagonists(VKA), mainly warfarin, dominated stroke prevention in atrial fibrillation(AF), the commonest arrhythmia globally(1, 2). Successive approval of four direct oral anticoagulants(DOACs: dabigatran(3), apixaban(4), rivaroxaban(5), edoxaban(6)) changed the landscape, with early adoption in guidelines(7, 8) and quality improvement initiatives(9). DOACs are often preferred over vitamin K antagonist(VKA) due to reduced international normalised ratio(INR) monitoring, but only if OAC services are fully decommissioned, and DOACs are taken appropriately. Paradoxically, removal of need for INR monitoring also removes additional patient-clinician engagement that encourages adherence(taking drugs as prescribed) and persistence(continuation of therapy)(10); both pertinent to OAC with a lifelong therapeutic indication.

Despite its importance in the context of population ageing, declining cognitive function, multimorbidity and polypharmacy, adherence was unreported in trials of DOACs(3-6), despite short half-lives, particularly dabigatran and apixaban which require twice-daily dosing(11). Reported trial persistence was highest for dabigatran(79.3% at low-dose) and lowest for edoxaban(65.6% at high-dose)(3-6)(**Web Table 1**). All DOACs have proven efficacy compared with VKA, albeit at much lower time in therapeutic range(TTR) in trials than usual clinical practice, but “head-to-head” DOAC trial comparisons are unlikely. However, several studies have shown suboptimal adherence and persistence for DOACs in different countries and settings, even compared with VKA, and effective interventions are lacking(12-14). Underlying causes include factors at social, economic, health system, clinician and patient levels. Although all patient-level factors are not captured, electronic health records(EHR) allow population-level studies of adherence and persistence together across all DOACs in the same dataset, which are rare(13).

Only one study to-date has considered all metrics of drug utilisation(“initiation”, “implementation” and “discontinuation”) together rather than “adherence” or “persistence” in isolation(15) for OAC in AF(13). Steps in drug utilisation may be described as the “prescription-persistence cascade”(from “recommendation” to “persistence”/“continuation”), estimable from EHR. For OAC in AF, the steps are “*recommendation*”(eligible for OAC); “*initial prescription*”(≥1 OAC prescription), “*dispensing*”(no EHR data), “*initiation by patient*”(no EHR data), “*adherence*”/“*implementation*”(adherent to OAC), and “*persistence*”/“*continuation*”(persistent to OAC). Interaction between adherence and persistence is often overlooked, e.g. “persistent and non-adherent”(i.e. continuing medications but not taking as prescribed) versus “non-persistent and non-adherent”(i.e. discontinued medications and also not taking as prescribed).

The UK has universal primary healthcare, enabling large-scale, representative datasets where uptake, adherence and persistence for different DOACs can be studied. We used The Health Improvement Network(THIN) database in the UK to investigate adherence and persistence for OACs in individuals with AF, focusing on: (i)time trends since DOAC introduction at health system level and after initiation in individuals; (ii)relative impact of sociodemographic and baseline risk factors and treatment characteristics; and (iii)associations between adherence and persistence.

Methods

The study conformed to STROBE recommendations(16)

Data source

The Health Improvement Network (THIN) database includes longitudinal, anonymised EHRs from over 500 UK general practices using Visionsoftware(INPS, www.inps4.co.uk/), representative of the UK population(17).

Study population

Our retrospective cohort included individuals aged ≥ 18 years with first-ever, non-valvular AF diagnosis between January 2011 and December 2016 and first prescription of VKA/DOAC on or after the date of AF diagnosis. The date of first prescription became the index date. For inclusion, patients needed ≥ 90 days of follow-up. Individuals with ≥ 1 prescription of VKA/DOAC were eligible for inclusion in adherence/persistence analyses. Exclusion criteria were taking OAC for other indications(e.g. deep-vein thrombosis and pulmonary embolism). Follow-up was until outcome event, death, the patient leaving the database or the most recent data upload.

Baseline covariates

Baseline factors were assessed: demography(age, sex, Townsend Deprivation Index Quintile-Level 1 the least deprived category), comorbidities(heart failure, hypertension, diabetes mellitus, stroke/TIA, vascular disease, liver disease, hypercholesterolaemia-i.e. on statin and/or had hypercholesterolaemia), social history(alcohol misuse, smoking status), drug history(aspirin, statin, BP-lowering drugs and mean number of drugs including OAC, prescribed in ≤ 365 days until, but not including, the episode start date). CHA₂DS₂VASc (18) and "HASBLED-1" (rather than HASBLED(19), since INR and "labile INR" were not available) scores were calculated from available variables, and categorised based on current guidelines.

Outcomes

Outcomes were adherence to and persistence with OACs. Adherence was estimated by proportion of days covered(PDC) over the year following first prescription of VKA/DOAC, more accurately reflects patient behaviour and treatment continuity than other adherence measures(20):

$$\text{PDC} = \frac{\text{Number of days with drug supplied (up to maximum of 365)}}{\text{Number of days between first and (last+30 days) prescription or 365 days*}}$$

*(whichever is shorter)

Each prescription was assumed to last 30 days, unless a new prescription was issued within 30 days, in which case the original prescription's duration was assumed to equal the gap between the two prescriptions. Patients with only one OAC prescription were classed as 'missing' and not included in the estimation. Impact of varying PDC threshold to 70% and 90% was examined for all OACs. Adherence was defined as PDC > 80% like previous studies(21) and produced more stable estimates. Individuals prescribed VKA/DOAC were deemed persistent until a prescription gap > 90 days on that or an alternative OAC('switch') in which case they were non-persistent, or there was no further longitudinal data(in which case persistence status was unknown beyond that time). For each DOAC, proportion of switching to VKA or another DOAC was analysed over 12-months. Impact of varying prescription gaps to 60 and 120 days was examined for all OACs. Like previous studies(22), the 90-day prescription gap was used, providing more stable estimates. As prescription gaps lengthened, persistence improved more with VKA than DOACs(**Web Table 3**).

Statistical analysis

Baseline characteristics were analysed by OAC. If any OAC/DOAC group consisted of <100 individuals, sample size was deemed too low to undertake meaningful analysis. Persistence was estimated using Kaplan-Meier product-limit estimator. Crude persistence for different OACs was estimated through survival life tables(adopting different prescription gaps) and ascertaining number and percentage(95% confidence intervals) of patients still in the study(i.e. persistent or uncensored) after 1 year. After stratification by adherence/persistence status at 12 months, baseline characteristics were determined. Chi-squared test and ANOVA test were used for categorical and continuous covariates respectively. Relative effects of OACs on non-adherence and non-persistence were modelled using univariable and multivariable logistic regression

and Cox proportional hazard regression (simple and multiple) respectively. For multivariable analyses, we adjusted for date of first OAC prescription (relative to study start date), CHA₂DS₂VASc, HASBLED-1, Townsend Deprivation Index quintile and number of drugs. Optimal adjustments for CHA₂DS₂VASc, HASBLED-1, number of drugs and date of first OAC prescription were investigated using continuous variables (including potential quadratic effects) or clinically appropriate categorisation. Models were compared using Bayes Information Criterion (BIC) and the optimal model chosen based on lowest BIC. For Cox regression, the proportional hazards assumption was investigated by adding interactions with 'time in study'. Interactions were included if they improved the model (by BIC criterion). For non-adherence, a sensitivity analysis was performed in those who had ≥6 months' potential OAC coverage (i.e. ≥6 months between date of first prescription and date of last prescription plus 30 days) and ≥12 months' OAC potential coverage to reduce potential bias in estimated adherence in short treatment periods, leading to overestimate of PDC (**Web Table 2(a)** and **2(b)**). For non-persistence, two sensitivity analyses were based around the chosen Cox regression model by: (i) adding interactions (linear; linear and quadratic) between OAC and timing of first OAC prescription (relative to study initiation); and (ii) reclassifying those switching to another OAC as censored (cessation of observation) rather than non-persistent to first OAC prescribed.

Results

Among 4,354,740 individuals, 154,180 with AF were identified, of whom 36,652 met inclusion criteria for analysis: VKA($n=25,008$, 68.2%), dabigatran($n=1285$, 3.5%), rivaroxaban($n=5855$, 16.0%) and apixaban($n=4504$, 12.3%)(**Figure 1**). Major exclusions were no new AF diagnosis($n=95237$) and absence of OAC prescription data($n=17065$). Edoxaban was UK-approved in September 2015, leading to inadequate sample size for analysis($n=45$) in the study period.

The study population had mean age 74.4(standard deviation 10.5) years and was 45% female. Cardiovascular comorbidities were common; hypertension(62.6%) and hypercholesterolaemia(71.8%), with median CHA₂DS₂VASc of 3(interquartile range, IQR 2-4). Polypharmacy was common(number of drugs: median 6, IQR 4-7). Individuals on dabigatran had lower CHA₂DS₂VASc scores, rates of hypertension and current smoking, with no other significant baseline differences across different OACs(**Table 1**).

Adherence

Adherence was 55.2%(54.6-55.7) overall, and 51.2%(50.6-51.8), 66.5%(63.7-69.2), 63.1%(61.8-64.4) and 64.7(63.2-66.1) for VKA, dabigatran, rivaroxaban and apixaban respectively(**Web Table 3**).

In univariable analysis, likelihood of non-adherence with DOACs was lower than with VKA(OR 0.53,0.47-0.60; 0.61,0.58-0.65; and 0.57,0.54-0.61 for dabigatran, rivaroxaban and apixaban respectively)(**Table 2** and **Web Table 2(d)**). In multivariable analysis, likelihood of non-adherence was similar for dabigatran(0.54,0.48-0.62), but higher for rivaroxaban(0.76,0.71-0.82) and apixaban(0.77,0.71-0.84). Increasing comorbidity(by CHA₂DS₂VASc) was associated with decreased likelihood of non-adherence(1.00,0.93-1.08; 0.94,0.88-1.01 and 0.81,0.74-0.89 for CHA₂DS₂VASc=2, 3-4 and 5-9 respectively, compared with CHA₂DS₂VASc=0-1), but not for HASBLED-1 score(0.98,0.93-1.05 for HASBLED-1 $\geq$ 3, compared with HASBLED-1 0-2). Number of drugs and Townsend quintile were not associated with non-adherence(**Table 2**). Age $\geq$ 75 years, diabetes, female gender and anaemia were associated with reduced risk of non-adherence, while hypertension and vascular disease were associated with increased risk(**Web Table 2(c)**). Non-adherence was non-linearly associated with time since introduction of DOACs, increasing for approximately two years(to early 2013) before starting to decrease, returning to its original level by early 2015 (**Web Figure 1(a)**) and then dropping below its original level. Baseline characteristics by adherence status are in **Web Table 4**. **Web Table 2(e)** illustrates no important differences in the effect of time since introduction of DOACs between different OACs (**Web Figure 1(b)**). Sensitivity analysis in only those who had at least 6 or 12 months of OAC prescriptions showed little impact on relative non-adherence for dabigatran, and VKA; however estimated ORs for rivaroxaban and apixaban both decreased to 0.65(for 12-month restriction)(**Web Tables 2(a)** and **(b)**).

Persistence

One-year persistence was 65.9%(65.4-66.5) overall, and 63.4%(62.8-64.0), 61.4%(58.3-64.2), 72.3%(70.9-73.7) and 78.7%(77.1-80.1) for VKA, dabigatran, rivaroxaban and apixaban respectively. Persistence reduced over three years for all OAC and was highest for apixaban and lowest for VKA and dabigatran(**Figure 2**).

In univariable analysis, apixaban had lowest(HR 0.53,0.50-0.57) and dabigatran had highest(HR 1.02,0.93-1.11) risk of non-persistence at 1 year, relative to VKA. **Table 3** illustrates the optimal multivariable model, where the effects of CHA₂DS₂VASc and OAC required an interaction term with time since first OAC prescription. The interaction between time and date of first prescription shows that immediately following first prescription, apixaban still had the lowest risk(0.53,0.46-0.60) and dabigatran the highest(1.24,1.08-1.42), relative to VKA. Risk of non-persistence did not change over time for apixaban(0.91,0.78-1.06 per year), but reduced over time for dabigatran(0.75,0.65-0.86 per year) and rivaroxaban(0.69,0.62-0.77 per year). Immediately after first prescription, increasing comorbidity, when measured by CHA₂DS₂VASc score, was associated with reduced risk of non-persistence(0.71,0.66-0.76; 0.66,0.62-0.71 and 0.69,0.63-0.76 for CHA₂DS₂VASc=2, 3-4 and 5-9 respectively, compared with CHA₂DS₂VASc=0-1). However, this risk was lessened over time(1.05,0.99-1.12 and 1.13,1.06-1.20 per year for CHA₂DS₂VASc=2 and 5-9 respectively). There was no significant effect of HASBLED-1(1.04,0.99-1.08 for HASBLED-1 $\geq$ 3, compared with HASBLED-1 0-2), nor for number of drugs nor Townsend quintile(**Table 3**). Overall, risk of non-persistence increased from 2011 until 2016(1.03,1.01-1.05 per year). Heart failure, vascular disease, chronic kidney disease, prior bleeding and alcohol misuse were associated with increased risk of non-persistence, while hypertension and

age>65 years were associated with reduced risk. Non-persistence was more likely for dabigatran soon after initiation, but the effect relative to VKA and to apixaban declined over the period of an individual's prescription, although this did not happen relative to rivaroxaban(**Web Table 5(a)** and **(b)**). Baseline characteristics are presented by 1-year persistence status in **Web Table 6**.

Persistence and adherence

Of 36652 individuals, 31.0% had <1 year of data, and of these, 15.8% had primary non-adherence, 27.9% were non-adherent and 56.2% were adherent. Among 25263 individuals with ≥1 year of data, primary non-adherence(3.5%) was less common than non-adherent, non-persistent(21.2%), adherent, non-persistent(8.6%), non-adherent, persistent(26.5%) and persistent, adherent(40.2%). Differences between OACs were significant($p<0.001$). Primary non-adherence was highest with dabigatran(7.8%)and lowest(2.7%) with apixaban, while persistent adherence was highest with apixaban(50.7%) and lowest with VKA (38.2%). Non-adherent, non-persistence was greatest with VKA(23.4%), and least with apixaban(12.4%). Non-adherent persistence was highest with apixaban(29.0%) and lowest with dabigatran(19.5%)(**Table 4**).

Switching

In non-persistent individuals, switching rates were 20.3%, 18.8%, 40.3%, 27.0% and 18.5% for all OACs, VKA, dabigatran, rivaroxaban and apixaban respectively. In primary non-adherent individuals, corresponding rates were 45.2%, 40.9%, 69.1%, 51.2% and 44.7% respectively(**Web Table 7**). When“switching” was censored rather than “non-persistent”, differences between OACs were more consistent over time(since first prescription) and non-persistence was generally lower for dabigatran than VKA, with an increasing trend in the difference with time on OAC. The effect was less for other DOACs relative to VKA(Web Fig 1 and Web Table 8).

Discussion

In this study of long-term persistence and adherence across all OACs in AF, we have four findings. First, primary non-adherence is uncommon (3.5%) and over time on a DOAC, likelihood of non-adherence and non-persistence increases. Second, the proportion of individuals on OACs who are both adherent and persistent at 1 year is low (40.2%) with heterogeneity across different OACs. Third, population-level time trends in adherence and persistence exist after new drugs (DOACs in this case) are introduced. Fourth, increased comorbidities were associated with reduced risk of non-adherence and non-persistence for DOACs, but number of drugs was not.

Observed rates of primary nonadherence for OAC are comparable to a recent Spanish study using large-scale regional EHR (12). Primary nonadherence varies across different drugs but our estimates appear lower than other chronic disease medications ($\leq 20.8\%$ for lipid-lowering drugs (23)). Rates are generally lower in European populations (23) than in North America, probably due to greater provision of prescription medication in public-funded health systems. Our results suggest that improved adherence and persistence require longer-term monitoring, rather than current strategies emphasising drug adherence post-initiation. Greater switching with dabigatran than other OACs may reflect greater discontinuation (due to dyspepsia or other side-effects (3)), or prescription patterns favouring other DOACs (24).

For drugs to be effective, both adherence and persistence are prerequisites. It is therefore concerning that adherent persistence at 1 year after initial prescription ranged from only 38.2% to 50.7% for VKA and apixaban respectively. Non-adherent, non-persistent individuals constituted 23.4% and 12.4% for users of VKA and apixaban respectively (**Table 4**). Adherence and persistence should be considered in combination, yet this is rare in both research and clinical practice (25). Furthermore, observations that non-adherent, persistence is more common than adherent, non-persistence and that these proportions vary by type of OAC highlight need for measurement of both metrics and potential for personalising approaches to improved drug utilisation. Factors which influence choice of drug in the same class include pharmacodynamics, pharmacokinetic, tolerability and cost (26), to which adherence and persistence may be added.

DOACs are unusual for several new drugs in the same class entering the market in a short timeframe. Other examples are statins, antihypertensives and novel hypoglycaemic agents, but four new drugs in a five-year period is extraordinary. DOACs have proven efficacy and effectiveness over VKA, and appropriate prescribing of OACs in AF has improved in the UK between 2000 and 2016 (27). However, there have been variations in prescription across DOACs over time (28). Our analyses add that when DOACs were first prescribed, persistence to all DOACs appears to have been initially higher than to VKA and in some cases increase further over time on the OAC, with clear differences between different OACs. Findings were sensitive to how switched of drugs was considered, with persistence to dabigatran, which had highest rates of 'switching', appearing much lower than to VKA when 'switchers' were classed as 'non-persistent' versus censored ('no longer observed for that OAC'). Other possible reasons for differences between OACs include side effect profile, marketing strategies, and varying procurement and prescription practices. Adherence and persistence can have far-reaching implications on drug cost, effectiveness and policy at population level (29) and should be monitored at population level.

Our findings are consistent with previous studies which have shown associations between polypharmacy (12, 30) and increased comorbidities (25) and reduced risk of non-adherence or non-persistence for OACs and other cardiovascular medications. Understanding each of the multiple steps in the prescription-persistence cascade may aid design and implementation of better interventions to improve drug utilisation. In routine clinical practice, EHR-based methods may be used to highlight individuals at greater risk of non-adherence or non-persistence, e.g. by suggesting that persistence should be more of a focus for improvement than adherence, or for monitoring long-term adherence/persistence.

The major strength of our analysis is consideration of the relationship between adherence and persistence together across all DOACs and VKA in the same population. As well as sociodemographic, health and medication characteristics, the influence of time was also analysed. Even in large-scale datasets with prescription data, there are several limitations. First, we did not have dispensing data, and therefore used previously validated methods to estimate adherence/persistence from prescription data. Our methods may be more uncertain for VKA than for DOACs (e.g. differential impact of varying prescription gaps on persistence), possibly due to patients on warfarin often having longer duration prescriptions than DOACs.

Second, missing prescription data meant that not all eligible individuals could be included due to incompleteness of follow-up. On the other hand, these are real-world data, which are routinely available and nationally representative. Third, relatively small numbers of patients could be included for DOACs, but numbers were comparable with other studies. Fourth, TTR would be a better measure of adherence for VKA but could not be estimated in our analysis, due to lack of INR data (which also limited our HASBLED analysis). Finally, we focused on initial OAC prescription in OAC-naïve patients to minimise bias in adherence/persistence based on previous OAC use, but did not focus on second and subsequent OACs used, where there would be more bias, and greater consideration of overall treatment pattern which requires consideration of multi-drug, multi-disease adherence/persistence over time.

Conclusions

Our study shows changes in adherence and persistence for DOACs over time in AF. Since these are usually lifelong therapies, more emphasis should be placed on long-term adherence and persistence in clinical practice and research. Standardisation is required for EHR methods of adherence and persistence estimation across drugs, diseases and datasets. Persistence and adherence may have different determinants and should be studied together in EHR. Better understanding of these factors will lead to interventions which are more likely to improve adherence and persistence at individual and population levels across OACs and other drugs. Post-marketing surveillance should take into account adherence and persistence particularly for multiple drugs in the same class where head-to-head trials are unlikely.

References

1. Chugh SS, Roth GA, Gillum RF, Mensah GA. Global burden of atrial fibrillation in developed and developing nations. *Glob Heart*. 2014;9(1):113-9.
2. Gage BF, Cardinali AB, Albers GW, Owens DK. Cost-effectiveness of warfarin and aspirin for prophylaxis of stroke in patients with nonvalvular atrial fibrillation. *JAMA*. 1995;274(23):1839-45.
3. Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, et al. Dabigatran versus warfarin in patients with atrial fibrillation. *The New England journal of medicine*. 2009;361(12):1139-51.
4. Granger CB, Alexander JH, McMurray JJV, Lopes RD, Hylek EM, Hanna M, et al. Apixaban versus Warfarin in Patients with Atrial Fibrillation. *New Engl J Med*. 2011;365(11):981-92.
5. Patel MR, Mahaffey KW, Garg J, Pan GH, Singer DE, Hacke W, et al. Rivaroxaban versus Warfarin in Nonvalvular Atrial Fibrillation. *New Engl J Med*. 2011;365(10):883-91.
6. Giugliano RP, Ruff CT, Braunwald E, Murphy SA, Wiviott SD, Halperin JL, et al. Edoxaban versus warfarin in patients with atrial fibrillation. *N Engl J Med*. 2013;369(22):2093-104.
7. Kirchhof P, Curtis AB, Skanes AC, Gillis AM, Samuel Wann L, John Camm A. Atrial fibrillation guidelines across the Atlantic: a comparison of the current recommendations of the European Society of Cardiology/European Heart Rhythm Association/European Association of Cardiothoracic Surgeons, the American College of Cardiology Foundation/American Heart Association/Heart Rhythm Society, and the Canadian Cardiovascular Society. *Eur Heart J*. 2013;34(20):1471-4.
8. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Jr., et al. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2014;64(21):e1-76.
9. Lewis WR, Piccini JP, Turakhia MP, Curtis AB, Fang M, Suter RE, et al. Get With The Guidelines AFIB: novel quality improvement registry for hospitalized patients with atrial fibrillation. *Circ Cardiovasc Qual Outcomes*. 2014;7(5):770-7.
10. Raebel MA, Schmittiel J, Karter AJ, Konieczny JL, Steiner JF. Standardizing terminology and definitions of medication adherence and persistence in research employing electronic databases. *Med Care*. 2013;51(8 Suppl 3):S11-21.
11. Clemens A, Noack H, Brueckmann M, Lip GY. Twice- or once-daily dosing of novel oral anticoagulants for stroke prevention: a fixed-effects meta-analysis with predefined heterogeneity quality criteria. *PLoS One*. 2014;9(6):e99276.
12. Rodriguez-Bernal CL, Peiro S, Hurtado I, Garcia-Sempere A, Sanfelix-Gimeno G. Primary Nonadherence to Oral Anticoagulants in Patients with Atrial Fibrillation: Real-World Data from a Population-Based Cohort. *J Manag Care Spec Pharm*. 2018;24(5):440-8.
13. Mueller T, Alvarez-Madrado S, Robertson C, Bennie M. Use of direct oral anticoagulants in patients with atrial fibrillation in Scotland: Applying a coherent framework to drug utilisation studies. *Pharmacoepidemiol Drug Saf*. 2017;26(11):1378-86.
14. Paquette M, Riou Franca L, Teutsch C, Diener HC, Lu S, Dubner SJ, et al. Persistence With Dabigatran Therapy at 2 Years in Patients With Atrial Fibrillation. *J Am Coll Cardiol*. 2017;70(13):1573-83.
15. Vrijens B, De Geest S, Hughes DA, Przemyslaw K, Demonceau J, Ruppar T, et al. A new taxonomy for describing and defining adherence to medications. *Br J Clin Pharmacol*. 2012;73(5):691-705.
16. Gallo V, Egger M, McCormack V, Farmer PB, Ioannidis JP, Kirsch-Volders M, et al. STrengthening the Reporting of OBservational studies in Epidemiology--Molecular Epidemiology (STROBE-ME): an extension of the STROBE Statement. *PLoS Med*. 2011;8(10):e1001117.
17. Blak BT, Thompson M, Dattani H, Bourke A. Generalisability of The Health Improvement Network (THIN) database: demographics, chronic disease prevalence and mortality rates. *Inform Prim Care*. 2011;19(4):251-5.
18. Lip GY, Nieuwlaat R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the euro heart survey on atrial fibrillation. *Chest*. 2010;137(2):263-72.
19. Pisters R, Lane DA, Nieuwlaat R, de Vos CB, Crijns HJ, Lip GY. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. *Chest*. 2010;138(5):1093-100.
20. Forbes CA, Deshpande S, Sorio-Vilela F, Kutikova L, Duffy S, Gouni-Berthold I, et al. A systematic literature review comparing methods for the measurement of patient persistence and adherence. *Curr Med Res Opin*. 2018;34(9):1613-25.
21. Karve S, Cleves MA, Helm M, Hudson TJ, West DS, Martin BC. Good and poor adherence: optimal cut-point for adherence measures using administrative claims data. *Curr Med Res Opin*. 2009;25(9):2303-10.

22. Cataldo N, Pegoraro V, Ripellino C, Bertasi V, Di Turi R, Pastorello M, et al. Non-persistence risk and health care resource utilization of Italian patients with non-valvular atrial fibrillation. *Recenti Prog Med*. 2018;109(2):113-21.
23. Lemstra M, Nwankwo C, Bird Y, Moraros J. Primary nonadherence to chronic disease medications: a meta-analysis. *Patient Prefer Adherence*. 2018;12:721-31.
24. Loo SY, Dell'Aniello S, Huiart L, Renoux C. Trends in the prescription of novel oral anticoagulants in UK primary care. *British journal of clinical pharmacology*. 2017;83(9):2096-106.
25. Manzoor BS, Lee TA, Sharp LK, Walton SM, Galanter WL, Nutescu EA. Real-World Adherence and Persistence with Direct Oral Anticoagulants in Adults with Atrial Fibrillation. *Pharmacotherapy*. 2017;37(10):1221-30.
26. Brown MJ. A rational basis for selection among drugs of the same class. *Heart*. 2003;89(6):687-94.
27. Adderley NJ, Ryan R, Nirantharakumar K, Marshall T. Prevalence and treatment of atrial fibrillation in UK general practice from 2000 to 2016. *Heart*. 2019;105(1):27-33.
28. Zhu J, Alexander GC, Nazarian S, Segal JB, Wu AW. Trends and Variation in Oral Anticoagulant Choice in Patients with Atrial Fibrillation, 2010-2017. *Pharmacotherapy*. 2018;38(9):907-20.
29. Cutler RL, Fernandez-Llimos F, Frommer M, Benrimoj C, Garcia-Cardenas V. Economic impact of medication non-adherence by disease groups: a systematic review. *BMJ Open*. 2018;8(1):e016982.
30. Ofori-Asenso R, Ilomaki J, Tacey M, Si S, Curtis AJ, Zomer E, et al. Predictors of first-year nonadherence and discontinuation of statins among older adults: a retrospective cohort study. *Br J Clin Pharmacol*. 2019;85(1):227-35.

Tables and Figures

Figure 1. Flow diagram of study population for adherence and persistence analyses.

Figure 2. Kaplan-Meier analysis of persistence over time

Table 1. Baseline characteristics of study population.

Table 2. Likelihood of non-adherence by oral anticoagulant.

Table 3. Risk of non-persistence by anticoagulant.

Table 4. Adherence and persistence by oral anticoagulant.

Table 1. Baseline characteristics of study population.

	Overall (N=36,652)	VKA (n=25,008)	Dabigatran (n=1,285)	Rivaroxaban (n=5,855)	Apixaban (n=4,504)	P
Characteristic [n(%)]						
Age [Mean(SD)]	74.4(10.5)	74.3 (10.2)	73.4(11.2)	74.8(11.0)	74.8(11.0)	<0.001
Female	16,494(45.0)	11,186(44.7)	517(40.2)	2,671(45.6)	2,120(47.1)	<0.001
<i>Townsend Quintile [Median(IQR)]</i>	2(1-4)	2(1-4)	2(1-3)	3(1-4)	3(1-4)	
1	8,862(24.2)	6,124(24.5)	345(26.9)	1,347(23.0)	1,046(23.2)	<0.001
2	8,269(22.6)	5,706(22.8)	321(25.0)	1,300(22.2)	942(20.9)	
3	7,330(20.0)	4,964(19.9)	270(21.0)	1,231(21.0)	865(19.2)	
4	5,864(16.0)	4,065(16.3)	185(14.4)	897(15.3)	717(15.9)	
5	3,800(10.4)	2,540(10.2)	98(7.6)	574(9.8)	588(13.1)	
Missing	2,527(6.9)	1,609(6.4)	66(5.1)	506(8.6)	346(7.7)	
COMORBIDITIES [n(%)]						
Heart failure	2,700(7.4)	1,908 (7.6)	79(6.2)	387(6.6)	326(7.2)	0.016
Hypertension	22,955(62.6)	15,824(63.3)	751(58.4)	3,612(61.7)	2,768(61.5)	<0.001
Diabetes mellitus	6,691(18.3)	4,594(18.4)	210(16.3)	1,058(18.1)	829(18.4)	0.31
Stroke/TIA	4,622(12.6)	3,096(12.4)	160(12.5)	735(12.6)	631(14.0)	0.026
Vascular disease	4,793(13.1)	3,316(13.3)	149(11.6)	717(12.3)	611(13.6)	0.052
Alcohol misuse	977(2.7)	589(2.4)	36(2.8)	175(3.0)	177(3.9)	<0.001
CKD	7,844(21.4)	5,426(21.7)	221(17.2)	1,229(21.0)	968(21.5)	0.002
Liver disease	100(0.3)	67(0.3)	4(0.3)	15(0.3)	14(0.3)	0.94
Hypercholesterolaemia	26,328(71.8)	17,979(71.9)	915(71.2)	4,172(71.3)	3,262(72.4)	0.56
Smoking Status						
Current smoker	3,374(9.2)	2,272(9.1)	103(8.0)	585(10.0)	414(9.2)	<0.001
Ex-smoker	13,928(38.0)	9,711(38.8)	484(37.7)	2,125(36.3)	1,608(35.7)	
Never smoked	18,484(50.4)	12,472(49.9)	666(51.8)	2,993(51.1)	2,353(52.2)	
Not indicated	866(2.4)	553(2.2)	32(2.5)	152(2.6)	129(2.9)	
RISK SCORES [n(%)]						
CHA₂DS₂-VASc						
[0-1]	5,856(16.0)	3,887(15.5)	263(20.5)	984(16.8)	722(16.0)	<0.001
[2]	7,192(19.6)	4,939(19.8)	279(21.7)	1,106(18.9)	868(19.3)	
[3-4]	17,894(48.8)	12,324(49.3)	571(44.4)	2,850(48.7)	2,149(47.7)	
[5-9]	5,710(15.6)	3,858(15.4)	172(13.4)	915(15.6)	765(17.0)	
HASBLED-1						
[0-2]	28,279(77.2)	19,298(77.2)	1,047(81.5)	4,508(77.0)	3,426(76.1)	<0.001
[3-8]	8,373(22.8)	5,710(22.8)	238(18.5)	1,347(23.0)	1,078(23.9)	
DRUGS [n(%)]						
Aspirin	20,510(56.0)	14,175(56.7)	683(53.2)	3,116(53.2)	2,536(56.3)	<0.001
Statin	17,185(46.9)	11,803(47.2)	551(42.9)	2,651(45.3)	2,180(48.4)	<0.001
BP-lowering drugs	29,136(79.5)	20,007(80.0)	961(74.8)	4,563(77.9)	3,605(80.0)	<0.001
Number of Drugs [Mean(SD)]	5.5(2.2)	5.5(2.2)	5.3(2.2)	5.5(2.2)	5.7(2.2)	<0.001

Table 2 Likelihood of non-adherence by oral anticoagulant.

Odds ratios (95% CI)	Univariable	Multivariable	
N	33,960	31,615	P-value
VKA	1.00 (-)	1.00 (-)	
Dabigatran	0.53 (0.47 to 0.60)	0.54 (0.48 to 0.62)	<0.001
Rivaroxaban	0.61 (0.58 to 0.65)	0.76 (0.71 to 0.82)	
Apixaban	0.57 (0.54 to 0.61)	0.77 (0.71 to 0.84)	
CHA ₂ DS ₂ Vasc			
[0,1]		1.00 (-)	
[2]		1.00 (0.93 to 1.08)	<0.001
[3,4]		0.94 (0.88 to 1.01)	
[5,9]		0.81 (0.74 to 0.89)	
HASBLED-1			
[0,2]		1.00 (-)	
[3,9]		0.98 (0.93 to 1.05)	0.62
Number of drugs			
Continuous/ Linear		0.99 (0.98 to 1.00)	0.067
Townsend Quintile			
[1]		1.00 (-)	
[2]		0.93 (0.87 to 0.99)	<0.001
[3]		0.86 (0.80 to 0.91)	
[4]		0.91 (0.85 to 0.97)	
[5]		0.86 (0.80 to 0.94)	
Date of first prescription ^a (years after 1 st January 2011)			
Continuous/ Linear		1.29 (1.22 to 1.37)	<0.001
Continuous/ Quadratic		0.94 (0.93 to 0.95)	
BIC	46263.63	42880.41	

^a Time difference (in years) between the date of the first ever OAC prescription for each patient and the start date of the study (1st January 2011). This suggests that the maximum effect of calendar time occurs at $-\ln(1.29)(2^*/\ln(0.94))=2.13$ years

Table 3 Risk of non-persistence by oral anticoagulant.

Hazard ratios (95% CI)	Univariable	Multivariable
N	36,636	34,109
OAC (effect on first prescribing)		
VKA	1.00 (-)	1.00 (-)
Dabigatran	1.02 (0.93 to 1.11)	1.24 (1.08 to 1.42)
Rivaroxaban	0.71 (0.67 to 0.74)	0.85 (0.77 to 0.93)
Apixaban	0.53 (0.50 to 0.57)	0.53 (0.46 to 0.60)
Time-dependent effect of OAC (per year of prescriptions)		
VKA		1.00 (-)
Dabigatran		0.75 (0.65 to 0.86)
Rivaroxaban		0.69 (0.62 to 0.77)
Apixaban		0.91 (0.78 to 1.06)
CHA₂DS₂Vasc (effect on first prescribing)		
[0,1]		1.00 (-)
[2]		0.71 (0.66 to 0.76)
[3,4]		0.66 (0.62 to 0.71)
[5,9]		0.69 (0.63 to 0.76)
Time-dependent effect of CHA₂DS₂Vasc (per year of prescriptions)		
[0,1]		1.00 (-)
[2]		1.05 (0.99 to 1.12)
[3,4]		1.12 (1.07 to 1.18)
[5,9]		1.13 (1.06 to 1.20)
HASBLED-1		
[0,2]		1.00 (-)
[3,9]		1.04 (0.99 to 1.08)
Number of drugs		1.00 (0.99 to 1.01)
Townsend Quintile		
[1]		1.00 (-)
[2]		0.99 (0.95 to 1.03)
[3]		0.95 (0.91 to 1.00)
[4]		0.97 (0.93 to 1.02)
[5]		0.97 (0.92 to 1.03)
Date of first prescription ^a (years after 1st January 2011)		1.03 (1.02 to 1.05)

^a Time difference (in years) between the date of the first ever OAC prescription for each patient and the start date of the study (1st January 2011) .

^b Time-dependent covariates are interacted with survival analysis time (e.g. time on a specific OAC), i.e. (end date of persistence or exit from/end of study - date of first ever OAC prescription)/365.25.

Table 4. Adherence and persistence by oral anticoagulant.

	Overall	VKA	Dabigatran	Rivaroxaban	Apixaban
Total [n(%)]	36652	25008	1285	5855	4504
Zero follow-up	16(0.04)	7(0.03)	1(0.08)	5(0.09)	3(0.07)
<i>Total <1 yr of data</i>	<i>11373(31.0)</i>	<i>5307(21.2)</i>	<i>441(34.3)</i>	<i>3,014 (51.5)</i>	<i>2,611(58.0)</i>
Primary non-adherence	1,800(15.8)	818(15.4)	78(17)	531(17.6)	373(14.3)
Non-adherent	3178(27.9)	1,654(31.2)	94(21.3)	771(25.6)	659(25.2)
Adherent	6,395(56.2)	2,835(53.4)	269(61.0)	1,712(56.8)	1,579(60.5)
<i>Total ≥1 yr of data</i>	<i>25,263(68.9)</i>	<i>19694(78.8)</i>	<i>843(65.6)</i>	<i>2,836(48.4)</i>	<i>1,890(42.0)</i>
Primary non-adherence	876(3.5)	631(3.2)	66(7.8)	128(4.5)	51(2.7)
Non-adherent, non-persistent	5,352 (21.2)	4,616(23.4)	124(14.7)	378(13.3)	234(12.4)
Adherent, non-persistent	2,173 (8.6)	1711(8.7)	132(15.7)	232(8.2)	98 (5.2)
Non-adherent, persistent	6,699 (26.5)	5221(26.5)	164(19.5)	766(27.0)	548 (29.0)
Persistent, adherent	10,163 (40.2)	7515(38.2)	357(42.3)	1,332(47.0)	959 (50.7)