

Central Lancashire Online Knowledge (CLOK)

Title	The effect of match fatigue in elite badminton players using plantar pressure measurements and the implications to injury mechanisms
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/31518/
DOI	https://doi.org/10.1080/14763141.2020.1712469
Date	2020
Citation	Valldecabres, Raul, Richards, James and de Benito, Ana-Maria (2020) The effect of match fatigue in elite badminton players using plantar pressure measurements and the implications to injury mechanisms. Sports Biomechanics. ISSN 1476-3141
Creators	Valldecabres, Raul, Richards, James and de Benito, Ana-Maria

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1080/14763141.2020.1712469>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

**Main title: The effect of match fatigue in elite badminton players using
plantar pressure measurements and the implications to injury
mechanisms**

Running title: The effect of match fatigue in elite badminton players

Raúl Valldecabres^{1,2} (orcid.org/0000-0002-7478-7526)

Jim Richards³ (orcid.org/0000-0002-4004-3115)

Ana-María de Benito² (orcid.org/0000-0002-8836-4073)

¹ Doctorate School, Valencia Catholic University San Vicente Mártir. Valencia, Spain.

² Physical Activity and Sports Science Faculty, Valencia Catholic University San Vicente
Mártir. Valencia, Spain.

³ Allied Health Research Unit, University of Central Lancashire, Preston, United
Kingdom.

Corresponding author:

Raúl Valldecabres

raul.valldecabres@ucv.es

Main title: The effect of match fatigue in elite badminton players using plantar pressure measurements and the implications to injury mechanisms

Running title: The effect of match fatigue in elite badminton players

Word count: 7557

Abstract

The purpose of this study was to investigate the differences in plantar pressure under the lead and trail foot between two lunge tasks to the net in the dominant (LD) and non-dominant (LND) directions, and to explore how fatigue affects the plantar pressure patterns whilst performing movements before and after a competitive match. Peak and mean pressure were measured with the Biofoot-IBV in-shoe system from five repetitions of each task, with sensors positioned under the calcaneus, midfoot and phalanges on the lead and trail foot. Data were collected pre and immediately post playing an official 1st national league competition match. The study was conducted with a sample of thirteen 1st league badminton players. A 2x2 repeated ANOVA found significant differences between the two tasks and between pre and post match (fatigued state). Players also had different foot pressure distributions for the LD and LND tasks, which indicated a difference in loading strategy. In a fatigued state the plantar pressure shifted to the medial aspect of the midfoot in the trail limb, indicating a reduction in control and a higher injury risk during non-dominant lunge tasks.

Abstract word count: 172

Keywords: biomechanics; kinetics; insole-system; competition

Introduction

Badminton is reported to be the fastest racket sport when considering ball velocity (Jaitner & Gawin, 2007). Its popularity has grown since its inclusion at the '92 Olympic Games in Spain and the use of a new scoring system in 2006 (Chen, Wu, & Chen, 2011), with up to 200 million players worldwide (Kwan, Cheng, Tang, & Rasmussen, 2010). The sport is characterised by short duration high intensity actions placing a high demand on the anaerobic system (Cabello & González-Badillo, 2003). The sport requires good footwork including; rapid turning, pivoting, jumps, lunges and running; in order for players to reach and hit the shuttlecock and to return to a defensive position in the centre of the court (Gibbs, 1988) which has not been analysed previously. There is a lack of quantification the court movements on the literature in a real competition context.

It has previously been reported that up to 70% of injuries are to the lower limbs, with the remaining 30% being due to loss of balance, slips and falls, landing or collisions with other players (Hensley & Paup, 1979; Krøner et al., 1990; Reeves, Hume, Gianotti, Wilson, & Ikeda, 2015). However, as in other racquet and field sports, badminton cannot be appropriately simulated in the laboratory setting (Faude et al., 2007). The majority of match based studies on badminton performance have focussed on notational performance analysis (Abián, Castanedo, Feng, Sampedro, & Abián-Vicén, 2014; Barreira, Chiminazzo, & Fernandes, 2016), game characteristics (Phomsoupha & Laffaye, 2014), or physiological and ground reaction force data (Chen et al., 2011; Ramos, Del Castillo, Polo, Ramón, & Bosch, 2016).

Players footwork during lunge movements has been previously reported (Kuntze, Mansfield, & Sellers, 2010), however the nature of on court movements had not been considered until recently (Valdecabres, de Benito, Casal, & Pablos, 2017). Valdecabres et al. (2017) divided the court into 12 parts, and described 3 different on court footwork movements which were commonly used to hit the shuttle, these were; diagonal, transverse and longitudinal, of which more than 50% were diagonal court movements.

Several studies have investigated injuries during badminton (Hensley & Paup, 1979; Jørgensen & Winge, 1990), with a higher risk of injury during competitions when compared to training sessions (Jørgensen & Winge, 1990). Of these injuries, 43% have

84 been reported as patellar tendinopathy (Shariff, George, & Ramlan, 2009), which is
85 considered a common injury in sports with jumps, cutting manoeuvres and explosive
86 running tasks (Tibesku & Pässler, 2005). The clinical injury risk incidence has also been
87 analysed and found that 26% of players had previous symptoms before getting injured
88 (Fahlström, Björnstig, & Lorentzon, 1998) being overuse sprains and strains the most
89 musculoskeletal common injury (Goh, Mokhtar, & Mohaman, 2013; Hensley & Paup,
90 1979; Shariff, George, & Ramlan, 2009). Lower limb dominance is an important
91 consideration during sports tasks, and has been identified as a factor associated with
92 potential Anterior Cruciate Ligament injury risk (Negrete, Schick, & Cooper, 2007).
93 However, lower limb dominance seems to be more related to specific movement tasks
94 which require different side to side movements (Peters, 1988), such as one side being
95 used for more postural stabilization (Velotta, Weyer, Ramirez, & Winstead, 2011). This
96 was highlighted by Sinsurin, Vachalathiti, Srisangboriboon & Richards (2018) who found
97 better coordination during multi-direction jump landings on the non-dominant limb. The
98 literature also suggests differences in the injury incidence between the dominant and non-
99 dominant limbs for many sports, which has been attributed to side to side differences in
100 loading as a result of differences in movement strategies between the sides (Vauhnik et
101 al., 2008). For example, Kimura et al. (2010) reported that the knee on the lead limb side
102 was the most commonly injured in badminton players, in particular during cutting and
103 side-step movements to the racket-hand side. In addition, Krajnc et al. (2010) showed that
104 the non-dominant leg, defined as the one that is not used to kick a ball, suffered more
105 knee injuries than the dominant leg in soccer players, with players requiring more
106 operations with greater pain or discomfort after surgery on the non-dominant limb. Side
107 to side differences have also been identified in badminton (Kimura et al., 2010), with a
108 higher number of injuries seen on the non-dominant leg.

109
110 The incidence of lower limb injuries has also been shown to be linked to fatigue, which
111 has been associated with a decrease in neuromuscular control, impaired kinesthesia and
112 proprioception of joints, and a diminishing maximum voluntary strength (Dickin & Doan,
113 2008; Rozzi, Yuktanandana, & Pincivero, 2000; Saragiotto, Di Pierro, & Lopes, 2014;
114 Whyte, Burke, White, & Moran, 2015). Links between fatigue and exercise have been
115 investigated previously, however the exact definition of fatigue varies between studies
116 (López-Calbet & Dorado-García, 2006). Fatigue may be considered as multidimensional
117 or multifactorial (Hunter & Smith, 2007; Millet et al., 2011), which may be characterised

by the decrease of the capacity or ability to generate force or muscle power. This originates from physiological, mechanical and psychological modifications (Stirling, Von Tscharnner, Fletcher, & Nigg, 2012) and a reduction of performance, which has been described as a conservative response to maintain tissue integrity (Millet et al., 2011). Previous work considering fatigue in badminton players has shown that dynamic postural control and the quality of athletes' performance is lower in a post fatigue state (Sarshin, Mohammadi, Shahrabad, & Sedighi, 2011). Players' perceived fatigue may be assessed using the rating of perceived exertion scale (Borg, 1982a), which has been shown to be valid for the assessment of exercise intensity during badminton matches (Fernández, de la Aleja, Moya, Cabello, & Méndez, 2013).

Previous investigations into sports related tasks have shown plantar pressure is an important method of quantifying the magnitude and location of the force applied beneath the foot, which may be measured using in shoe pressure systems (Falda-Buscaiot, Hintzy, Rougier, Lacouture, & Coulmy, 2017; Navarro, Zahonero, Huertas, Vera, & Barrios, 2012). Fu, Liu & Wei (2009) studied badminton lunge tasks, and found the maximum peak pressure was distributed under the forefoot with lower pressures under the midfoot. Conversely, Hu, Li, Hong & Wang (2015) found lower pressures under the forefoot with the maximum pressure under the heel, indicating either differences in movement strategy or test protocol. Changes in distribution of foot pressure, for example midfoot loading, has been associated with foot pronation and patellofemoral pain (Thijs, Van Tiggelen, Roosen, De Clercq, & Witvrouw, 2007; Powers, 2003). In addition, the static assessment of foot posture using the Foot Posture Index (Redmond, Crosbie, & Ouvrier, 2006) has been shown to be important in the prediction of clinical subgroups in people with patellofemoral pain (Selfe et al., 2016). Therefore, the use of foot pressure may allow an assessment of changes in loading strategy and dynamic postural control on the trail and lead limbs which could be performed in the competition arena. This could be useful in the assessment of training and injury prevention, or when considering when to return to sport after an injury.

The aims of the present study were to investigate the differences in plantar pressure under the trail and lead foot between two lunge tasks to the net, and to explore how fatigue affects the plantar pressure patterns whilst performing movements before and after a competitive match. It was hypothesised that lunge tasks to the dominant side (LD) and

lunge tasks to the non-dominant side (LND) would show different foot loading strategies and that players would show changes in plantar pressure patterns in a fatigue state.

Methods

Participants

The inclusion criteria for both genders were; players in the 1st Spanish badminton league who played at least 3 times a week, with no injuries to the upper and lower limbs in the previous 6 months. In addition, all participants had no history of surgery or traumatic injury to the lower extremities or lower back, and no history of medical conditions that limit physical activity. Exclusion criteria included; presence of neuromuscular or vestibular conditions, visual impairment or back pain.

Thirteen right-handed badminton players (5 males and 8 females) participated in the study. Anthropometric measurements of height and weight were recorded; age 25.93 ± 10.05 years and bodyweight 64.30 ± 8.66 kg. Hand laterality was assessed using a previously validated questionnaire (Chapman & Chapman, 1987), which includes questions such as *which hand do you use to throw a ball* or *which foot do you use to kick a ball*, which has been used in previous studies (Brophy, Silvers, Gonzales, & Mandelbaum, 2010; English, Brannock, Chik, Eastwood, & Uhl, 2006). In addition, Foot Posture Index (FPI), a clinical tool that measures foot type between +12 (pronated) and -12 (supinated) was recorded (Redmond, Crosbie, & Ouvrier, 2006). This study was approved by the Valencia Catholic University San Vicente Mártir Ethics Committee and all data collection conformed to the Declaration of Helsinki. Volunteers gave written informed consent and parental consent for the four participants who were minors was obtained prior to participation.

Equipment

In shoe plantar pressure data were recorded using the Biofoot-IBV in-shoe system (Valencia, Spain), which consists of 64 sensors of 0.5mm thickness and 5mm diameter and has been previously validated and used to assess sports tasks (Martínez, Hoyos, Brizuela, Ferrús, & González, 1988; Marhuenda, Fuentes, Costa, Ferrús, & González, 2011). Not normalised peak and mean pressure data were used according to previous

studies (Patrick, & Donovan, 2018; Taylor, Nguyen, Griffin, & Ford, 2018). In addition, the regions of interest were normalised to foot size following Oliveira, Sousa, Santos, & Tavares (2012) with values for foot shape proposed by Hu et al. (2015).

Procedure

In order to familiarise the players, submaximal lunge tasks were performed to the right and left sides, LD and LND respectively for a right-handed player (figure 1). During the lunge, the role of the non-stepping limb is to support the bodyweight during the forward movement of the lead limb (Hofmann, Holyoak, & Juris, 2017) and reaches a maximum knee flexion during landing, this is followed by a movement backwards in order to recover to the initial starting position (Kuntze et al., 2010). The lunge tasks were performed at 45° to the net in the defensive position (Gibbs, 1988), and were defined as controlled movements of the knee such that the knee did not move in front of the ankle joint. The most natural start position for the lunge task was identified for each player in order to hit a shuttlecock that was hung using a fine thread at a height of 165cm, 10 cm from the net (figure 2). Prior to data collection, a standardised 10 minute warm-up was performed, which included active stretching of the quadriceps and hamstring muscles (Lam et al., 2017), specifically this involved five repetitions of 30 seconds per muscle; and a familiarization period which involved the participants performing as many repetitions of the lunge tasks as they needed to feel comfortable (Gribble, Hertel, & Plisky, 2012).

[Figures 1 and 2 near here]

Pressure sensors were then placed in both shoes by fixing them to the insole to avoid them moving within the shoe. Changes in foot pressure data were recorded at a sampling frequency of 265Hz. in directions. Each lunge task trial consisted of a lunge to the net, hitting the shuttlecock with a top-spin shot and moving back to the start position as fast as possible to simulate a match shot. Approximately 5 minutes after participants had played a competition match the lunge tests were repeated. In addition, the level of match intensity was assessed using the Rated Perceived Exertion (RPE) Scale (Borg, 1982a). This determined the perceived intensity of the exercise, from 0 (nothing at all) to 10 (very, very heavy) (Borg, 1982b).

The plantar surface of the foot was divided into 9 areas previously reported by Navarro et al. (2012) (figure 3), these included; hallux, 2nd and 3rd phalanges and 4th and 5th phalanges, 1st metatarsal, 2nd and 3rd metatarsals and 4th and 5th metatarsals, medial midfoot and lateral midfoot, and calcaneus. The peak pressure (PP) and average pressure (PMEAN) of the whole foot were recorded and an average value from 5 repetitions was found for each parameter for each movement direction for both the lead and trail feet.

[Figure 3 near here]

Statistical Analysis

Based on previous papers by Lam et al. (2017) and Park et al. (2017) and an alpha level of 0.05 and an 80% power, a sample size of at least 13 was found to be required to explore the differences between sides and fatigue state. The data were checked for normality using the Shapiro-Wilk test and found to be suitable for parametric testing. In addition, partial eta squared (η^2) was computed to determine the effect size which was interpreted as small 0.1, medium 0.3, and large 0.5. Statistical analysis was performed with SPSS 21.0 (IBM, Armonk, NT, USA).

A 2x2 repeated ANOVA found significant differences between the two tasks and between pre and post match (fatigued state).

Results

The players had a mean RPE after playing the competition match of 7.54 (SD 2.10), with a range of perception of fatigue between strong and extremely strong. The average match length was 26 minutes 29 seconds with a standard deviation of 8 minutes and 9 seconds. The FPI showed mean values of 1.17 (SD 1.79) for the lead foot and 1.50 (SD 1.72) for the trail foot, which corresponds to a neutral foot posture (Redmond, Crosbie, & Ouvrier, 2006). All players were at a similar level, and were all Spanish 1st league competitors and were playing for the Championship.

Significant main effects were seen between pre and post fatigue on the lead foot. These revealed significant differences under the 4th and 5th Phalangeal areas between pre and post fatigue for the peak and mean pressure ($p=0.035$, $\eta^2=0.32$; $p=0.031$, $\eta^2=0.33$) respectively, with lower values in a fatigue state. In addition, a reduction of lead foot peak pressure under the lateral side was seen with a corresponding increase in peak pressure under the medial side of the midfoot, although the latter was not statistically significant (table 1). A significant interaction was seen between the factors of lunge task and fatigue state under the 2nd and 3rd Metatarsals ($p=0.011$, $\eta^2=0.43$). Further post hoc analysis explored differences between pre and post fatigue for each lunge task separately, which showed an increase in peak pressure post fatigue for the LD task and a decrease with the LND task (table 1). In addition, the calcaneus mean pressure for the lead foot in the LD task was lower than the LND task ($p=0.032$, $\eta^2=0.33$) (Table 2), indicating a different strategy during landing between lunge tasks.

[Tables 1 and 2 near here]

Significant differences were also seen under the 1st Metatarsal on the trail foot pre and post fatigue for the peak and mean pressure ($p=0.048$, $\eta^2=0.29$; $p=0.046$, $\eta^2=0.29$) respectively, with a decrease in pressure post fatigue in both LD and LND tasks, and a corresponding significant increase in medial midfoot pressure. Finally, the medial midfoot showed an increase in peak and mean pressure post fatigue for both the LD and LND tasks ($p=0.022$, $\eta^2=0.37$; $p=0.046$, $\eta^2=0.29$) respectively (Tables 3 and 4), which would imply a shift in pressure to the medial midfoot as the players fatigue.

Significant main effects were seen between LD and LND tasks. These revealed differences in the peak and mean pressure under the Hallux ($p=0.008$, $\eta^2=0.45$; $p=0.020$, $\eta^2=0.38$) respectively. This showed the Hallux pressure was lower in the LND task than the LD task in the trail foot (tables 3 and 4), with a corresponding lower pressure in the LND task versus the LD task for the peak and mean pressure under the 4th and 5th Phalangeal areas ($p=0.026$, $\eta^2=0.35$; $p=0.017$, $\eta^2=0.39$) respectively (tables 3 and 4). In addition, significant differences were found between the LD and LND tasks in the mean pressure under the Hallux ($p=0.038$, $\eta^2=0.31$) (table 4). Furthermore, a significant interaction was also seen for the peak pressure under the 2nd and 3rd Phalangeal areas in

the trail foot between lunge task and fatigue state ($p=0.017$, $\eta^2=0.39$), with a greater pressure being seen during the LD task, although this was not significant (Table 3). A further post hoc analysis revealed a significant difference in peak pressure between pre and post fatigue in the LD task on the trail foot only ($p=0.045$, $\eta^2=0.30$).

[Tables 3 and 4 near here]

Discussion and Implications

A fatigue state has been shown to reduce athletes' performance and can lead to poor joint control (Chang, 2015) and a decrease in dynamic postural control in badminton players (Sarshin et al., 2011). Furthermore, the biomechanics and movement patterns have been shown to be modified as a result of fatigue (Chang, 2015), which could lead to a subsequent increase in injury rate. Footwork manoeuvres are therefore extremely important for badminton players and the consideration of how landing strategies change on the different limbs after fatigue may help us to understand injury mechanisms (Phomsoupha & Laffaye, 2014). Diagonal lunge tasks have been previously shown to account for approximately 50% of on court manoeuvres in both male and female players (Valldecabres et al., 2017). This movement is required to allow the players to move to hit the shuttlecock, and the speed and reaction time are important to allow the player to recover a defensive court position to be ready for the next shot (Gibbs, 1988). Therefore, the aim of this study was to explore differences in foot contact pressures under the lead and trail foot during two badminton specific lunge tasks to the net, and to determine how plantar pressure patterns change as the players' fatigue.

The Foot Posture Index has been used to determine foot type in athletes in different sports including; basketball, handball and running (Martínez-Nova et al., 2014). Our results show a neutral foot type across the participants. However, when athletes fatigue, their foot posture tends to move towards a more pronated position (Cowley & Marsden, 2013), which could be an indicator of arch collapse leading to an increase of plantar pressure under the medial midfoot. Jørgensen & Winge (1990) reported that more than 30% of lower limb injuries are to the foot or ankle joint. This could be related to a flattening of the arch with a more pronated foot position (Lange, Chipchase, & Evans, 2004), leading to associated injuries (Barton, Bonanno, Levinger, & Menz, 2010; Beeson, Phillips, Corr,

& Ribbans, 2009; Graham, Jawrani, & Goel, 2011; Raissi, Cherati, Mansoori, & Razi, 2009).

This study found an increase in pressure under the medial aspect of the lead foot, indicating a more pronated foot or a collapse of the arch post fatigue with the adoption of a flatter foot posture, which is in line with previous findings (Wei, Liu, Tian, & Fu, 2009). This would arguably increase the pronation and pronation velocity which have been previously identified as key factors in patellofemoral pain and patellar tendinopathy in runners (Powers, 2003; Thijs et al., 2007), the latter of which has been identified as an issue in 43% of badminton players (Shariff et al., 2009). In addition, the side-to-side differences found may help to explain the higher injury rates which have been reported on the non-dominant leg (Krajnc et al., 2010). One consideration to help mitigate this effect is the use of foot orthoses. These could be used to directly manage foot posture by supporting the arch and therefore decreasing the pronation and pronation velocity when athletes are in a fatigued state. This in turn may reduce the incidence of patellar tendinopathy (Mündermann, Nigg, Humble, & Stefanyshyn, 2003). However, more research is needed to explore the use of foot orthotics in badminton players, and their effect on potential injury mechanisms when athletes are in a fatigued state.

When comparing LD and LND tasks on lead foot, there are differences in pressure under the Hallux, a lower pressure under the 4th and 5th Phalangeal areas and a lower calcaneus mean pressure, which indicates a possible change in strategy during landing as Vauhnik et al. (2008) reported in other court sports. Lower pressure on the forefoot and lateral areas could be due to the involvement of pelvic rotation which has been reported by players who find dominant side movements easier than non-dominant side ones (Bazipoor, Shojaeddin, Shahhoseini, & Abdollahi, 2017). Further exploration of such compensations was beyond the scope of this current study

After fatigue, a significant decrease in mean pressure under the Hallux and 1st Metatarsal areas and a significant increase under medial midfoot were seen in both LD and LND tasks. This would support the shift in pressure to the medial midfoot and collapse of the arch when players reach a fatigued state which is in agreement with previous literature (Weist, Eils, & Rosenbaum, 2004), and supports that changes in plantar pressure distribution occur due to fatigue (Bisiaux & Moretto, 2008). This has implications to the

foot movement used to recover the initial position in the centre of the court with cutting and pivoting creating a higher tibia rotation torque (Oh, Kreinbrink, Wojtys, & Ashton-Miller, 2012) which has been linked with patella mal-alignment and an increased risk of patellofemoral pain (Sinclair & Dillon, 2016).

During both LND and LD lunge tasks the Hallux, 4th and 5th Metatarsals all showed lower mean and peak pressure on the trail limb (tables 3 and 4). The explanation for this could be different hip movement strategies when lunging towards the net, with movement into hip external rotation during the LD task, and a move into hip internal rotation when performing the LND task which has been previously reported by Valdecabres, de Benito, Littler & Richards (2018). In addition, the trail limb is in contact with the ground for longer and is responsible for the eccentric control as the person lunges forwards, which is similar to findings reported during fencing, where the trail/rear limb demonstrated lower pressures (Trautmann, Martinelli, & Rosenbaum, 2011).

This current study is the first to explore the plantar pressure patterns in the lead and trail foot. Previously published work by Hu et al. (2015) on plantar pressures during badminton movements has only taken into account the dominant/lead limb during lunge tasks to the net. This work offers a contribution to knowledge on the changes in movement strategies due to fatigue within a competition environment in elite players. However, more work is required to assess how fatigue affects pronation and pronation velocity, knee joint stability and compensations in pelvic rotation in badminton players.

This study was not without its limitations which included a large standard deviation in the age of the players recruited. A large standard deviation was seen in many of the parameters reported, this was due to variations in the magnitude of many of the parameters between the participants, however the direction of the changes seen were common among the participants which accounts for the significant differences seen, and supports the conclusion that a change in strategy occurs due to fatigue. In addition, no direct measures of the amount of pronation or pronation velocity were taken. Furthermore, greater fatigue has been previously reported as a competition progresses, however this current study did not report how many matches were played at the time of

assessment, although the match duration was similar to that reported by Abián-Vicén, Castanedo, Abián, & Sampedro (2013).

Conclusion

In conclusion, this study found that players have different strategies when moving to the dominant and non-dominant sides during lunge to the net tasks. Under a fatigue state the plantar pressure shifts to the medial aspect of the midfoot in the trail limb indicating a reduction in control of the midfoot and collapse of the arch. This could account for the higher injury risk in the non-dominant trail limb which has been previously reported in badminton players, therefore further work should consider both the lead and trail foot. These results provide new insights into the changes in foot function in a fatigued state, which could possibly be managed using foot orthoses.

Disclosure statement

There is no conflict of interest reported by authors.

Funding details

This study has been made possible by Generalitat Valenciana ACIF 121/2015 grants for PhD students and Valencia Catholic University San Vicente Mártir grant to new research groups.

Acknowledgments

We would like to thank Badminton Spanish Federation for support and access to the real competition context and Club Bádminton Alicante for allowing us a court in sports centre. Data collection was possible thanks to (in alphabetical order): Omar Aguar, Carlos Baeschlin, Joaquín Barrachina, Yvan Barthelemy, Claudio Alberto Casal, Diego Ceca, Raúl Fernández, Encarnación Liébana and Cristina Menescardi.

References

- Abián, P., Castanedo, A., Feng, X. Q., Sampedro, J., & Abián-Vicén, J. (2014). Notational comparison of men's singles badminton matches between Olympic Games in Beijing and London. *International Journal of Performance Analysis in Sport*, 14(1), 42-53. doi:10.1080/24748668.2014.11868701
- Abián-Vicén, J., Castanedo, A., Abián, P., & Sampedro, J. (2013). Temporal and notational comparison of badminton matches between men's singles and women's singles. *International Journal of Performance Analysis in Sport*, 13(2), 310–320. doi:10.1080/24748668.2013.11868650
- Barreira, J., Chiminazzo, J. G. C., & Fernandes, P. T. (2016). Analysis of point difference established by winners and losers in games of badminton. *International Journal of Performance Analysis in Sport*, 16(2), 687–694. doi:10.1080/24748668.2016.11868916
- Barton, C. J., Bonanno, D., Levinger, P., & Menz, H. B. (2010). Foot and ankle characteristics in patellofemoral pain syndrome: A case control and reliability study. *Journal of Orthopaedic & Sports Physical Therapy*, 40(5), 286-296. doi:10.2519/jospt.2010.3227
- Bazipoor, P., Shojaeddin, S. S., Shahhoseini, A., & Abdollahi, I. (2017). A comparison of foot plantar pressure in badminton players with normal and high-arched feet during the two-way lunge. *Journal of Rehabilitation Sciences and Research*, 1, 20-25. doi:10.30476/JRSR.2017.41114
- Beeson, P., Phillips, C., Corr, S., & Ribbans, W. J. (2009). Hallux rigidus: A cross-sectional study to evaluate clinical parameters. *The Foot*, 19(2), 80-92. doi:10.1016/j.foot.2008.12.001
- Bisiaux, M., & Moretto, P. (2008). The effects of fatigue on plantar pressure distribution in walking. *Gait & Posture*, 28(4), 693-698. doi:10.1016/j.gaitpost.2008.05.009

- 434 Borg, G. (1982a). A category scale with ratio properties for intermodal and
435 interindividual comparisons. In: Geissler HG, Petzold P, eds. *Psychophysical*
436 *Judgment and the Process of Perception*, 25-34.
- 437 Borg, G. (1982b). Psychophysical bases of perceived exertion. *Medicine and Science in*
438 *Sports Exercice*, 15(5), 377-381. doi:10.1249/00005768-198205000-00012
- 439 Brophy, R., Silvers, H. J., Gonzales, T., & Mandelbaum, B. R. (2010). Gender influences:
440 the role of leg dominance in ACL injury among soccer players. *British Journal of*
441 *Sports Medicine*, 44(10), 694-697. doi:10.1136/bjsm.2008.051243
- 442 Cabello, D., & González-Badillo, J. J. (2003). Analysis of the characteristics of
443 competitive badminton. *British Journal of Sports Medicine*, 37(1), 62–66.
444 doi:10.1136/bjsm.37.1.62
- 445 Chang, C. (2015). Research on the biomechanics analysis of technical movement in
446 fatigue period for badminton athletes. *International Journal of Simulation*
447 *Systems, Science & Technology*, 16(4B), 13.1-13.6.
448 doi:10.5013/IJSSST.a.16.4B.13
- 449 Chapman, L. J., & Chapman, J. P. (1987). The measurement of handedness. *Brain and*
450 *Cognition*, 6(2), 175-183. doi:10.1016/0278-2626(87)90118-7
- 451 Chen, H. L., Wu, C. J., & Chen, T. C. (2011). Physiological and notational comparison
452 of new and old scoring systems of singles matches in men's badminton. *Asian*
453 *Journal of Physical Education & Recreation*, 17(1).
- 454 Cowley, E., & Marsden, J. (2013). The effects of prolonged running on foot posture: a
455 repeated measures study of half marathon runners using the foot posture index
456 and navicular height. *Journal of Foot and Ankle Research*, 6(1), 1-6.
457 doi:10.1186/1757-1146-6-20
- 458 Dickin, D. C., & Doan, J. B. (2008). Postural stability in altered and unaltered sensory

459 environments following fatiguing exercise of lower extremity joints: Postural
 460 control following fatiguing exercise of lower extremity joints. *Scandinavian*
 461 *Journal of Medicine & Science in Sports*, 18(6), 765-772. doi:10.1111/j.1600-
 462 0838.2007.00760.x

463 English, R., Brannock, M., Chik, W. T., Eastwood, L. S., & Uhl, T. (2006). The
 464 relationship between lower extremity isokinetic work and single-leg functional
 465 hop-work test. *Journal of Sport Rehabilitation*, 15(2), 95-104.
 466 doi:10.1123/jsr.15.2.95

467 Fahlström, M., Yeap, J. S., Alfredson, H., & Söderman, K. (2006). Shoulder pain—a
 468 common problem in world-class badminton players. *Scandinavian Journal of*
 469 *Medicine & Science in Sports*, 16(3), 168-173. doi:10.1111/j.1600-
 470 0838.2004.00427.x

471 Falda-Buscaiot, T., Hintzy, F., Rougier, P., Lacouture, P., & Coulmy, N. (2017).
 472 Influence of slope steepness, foot position and turn phase on plantar pressure
 473 distribution during giant slalom alpine ski racing. *PLoS One*, 12(5), e0176975.
 474 doi:10.1371/journal.pone.0176975

475 Faude, O., Meyer, T., Rosenberger, F., Fries, M., Huber, G., & Kindermann, W. (2007).
 476 Physiological characteristics of badminton match play. *European Journal of*
 477 *Applied Physiology*, 100(4), 479-485. doi:10.1007/s00421-007-0441-8

478 Fernández, J., de la Aleja, J. G., Moya, M., Cabello, D., & Méndez, A. (2013). Gender
 479 differences in game responses during badminton match play: *Journal of Strength*
 480 *and Conditioning Research*, 27(9), 2396-2404.
 481 doi:10.1519/JSC.0b013e31827fcc6a

482 Fu, W. J., Liu, Y., & Wei, Y. (2009). The characteristics of plantar pressure in typical
 483 footwork of badminton. *Footwear Science*, 1(sup1), 113-115.

doi:10.1080/19424280903063630

Gibbs, M. J. (1988). Badminton-Teaching concepts. *Journal of Physical Education, Recreation & Dance*, 59(8), 92-94. doi:10.1080/07303084.1988.10606299

Graham, M. E., Jawrani, N. T., & Goel, V. K. (2011). Evaluating plantar fascia strain in hyperpronating cadaveric feet following an extra-osseous talotarsal stabilization procedure. *The Journal of Foot and Ankle Surgery*, 50(6), 682-686. doi:10.1053/j.jfas.2011.07.005

Gribble, P. A., Hertel, J., & Plisky, P. (2012). Using the star excursion balance test to assess dynamic postural-control deficits and outcomes in lower extremity injury: A literature and systematic review. *Journal of Athletic Training*, 47(3), 339-357. doi:10.4085/1062-6050-47.3.08

Hensley, L. D., & Paup, D. C. (1979). A survey of badminton injuries. *British Journal of Sports Medicine*, 13(4), 156–160. doi:10.1136/bjism.13.4.156

Hofmann, C. L., Holyoak, D. T., & Juris, P. M. (2017). Trunk and shank position influences patellofemoral joint stress of the lead and trail limb during the forward lunge exercise. *Journal of Orthopaedic & Sports Physical Therapy*, 47(1), 31-40. doi:10.2519/jospt.2017.6336

Hu, X., Li, J. X., Hong, Y., & Wang, L. (2015). Characteristics of plantar loads in maximum forward lunge tasks in badminton. *PLoS One*, 10(9), e0137558. doi:10.1371/journal.pone.0137558

Hunter, I., & Smith, G. A. (2007). Preferred and optimal stride frequency, stiffness and economy: changes with fatigue during a 1-h high-intensity run. *European Journal of Applied Physiology*, 100(6), 653-661. doi:10.1007/s00421-007-0456-1

Jaitner, T., & Gawin, W. (2007). Analysis of badminton smash with a mobile measure device based on accelerometry. In *ISBS-Conference Proceedings Archive* (Vol.

509 1).

510 Jørgensen, U., & Winge, S. (1990). Injuries in badminton. *Sports Medicine*, 10(1), 59-

511 64. doi:10.2165/00007256-199010010-00006

512 Kimura, Y., Ishibashi, Y., Tsuda, E., Yamamoto, Y., Tsukada, H., & Toh, S. (2010).

513 Mechanisms for anterior cruciate ligament injuries in badminton. *British Journal*

514 *of Sports Medicine*, 44(15), 1124-1127. doi:10.1136/bjsm.2010.074153

515 Krajnc, Z., Vogrin, M., Rečnik, G., Crnjac, A., Drobnič, M., & Antolič, V. (2010).

516 Increased risk of knee injuries and osteoarthritis in the non-dominant leg of

517 former professional football players. *Wiener Klinische Wochenschrift*, 122(S2),

518 40-43. doi:10.1007/s00508-010-1341-1

519 Krøner, K., Schmidt, S. A., Nielsen, A. B., Yde, J., Jakobsen, B. W., Møller-Madsen, B.,

520 & Jensen, J. (1990). Badminton injuries. *British Journal of Sports Medicine*,

521 24(3), 169–172. doi:10.1136/bjsm.24.3.169

522 Kuntze, G., Mansfield, N., & Sellers, W. (2010). A biomechanical analysis of common

523 lunge tasks in badminton. *Journal of Sports Sciences*, 28(2), 183-191.

524 doi:10.1080/02640410903428533

525 Kwan, M., Cheng, C. L., Tang, W. T., & Rasmussen, J. (2010). Measurement of

526 badminton racket deflection during a stroke. *Sports Engineering*, 12(3), 143-153.

527 doi:10.1007/s12283-010-0040-5

528 Lam, W. K., Ryue, J., Lee, K. K., Park, S. K., Cheung, J. T. M., & Ryu, J. (2017). Does

529 shoe heel design influence ground reaction forces and knee moments during

530 maximum lunges in elite and intermediate badminton players? *PloS one*, 12(3),

531 e0174604. doi:10.1371/journal.pone.0174604

- 532 Lange, B., Chipchase, L., & Evans, A. (2004). The effect of low-dye taping on plantar
533 pressures, during gait, in subjects with navicular drop exceeding 10 mm. *Research*
534 *Report*, 34(4), 9. doi:10.2519/jospt.2004.34.4.201
- 535 López Calbet, J., & Dorado Garcia, C. (2006). Fatiga, dolor muscular tardío y
536 sobreentrenamiento. *López Chicharro, J. y Fernández Vaquero, A. Fisiología del*
537 *ejercicio. 3ª edición. Madrid: Editorial Médica Panamericana.*
- 538 Marhuenda, A. M., Fuentes, J. V. H., Costa, G. B., Ferrús, E., & González, J. C. (2011).
539 Biofoot-IBV. Una técnica de registro y análisis de la distribución de presiones
540 plantares aplicable a la mejora del rendimiento deportivo. *Colección ICD:*
541 *Investigación en Ciencias del Deporte*, 19.
- 542 Martínez, A., Hoyos, J. V., Brizuela, G., Ferrús, E., & González, J. C. (1988). Biofoot-
543 IBV. Una técnica de registro y análisis de la distribución de presiones plantares
544 aplicable a la mejora del rendimiento deportivo. *Colección ICD: Investigación en*
545 *Ciencias del Deporte*, 19, 73-88.
- 546 Martínez-Nova, A., Gómez-Blázquez, E., Escamilla-Martínez, E., Pérez-Soriano, P.,
547 Gijón-Nogueron, G., & Fernández-Segúin, L. M. (2014). The foot posture index
548 in men practicing three sports different in their biomechanical gestures. *Journal*
549 *of the American Podiatric Medical Association*, 104(2), 154-158.
550 doi:10.7547/0003-0538-104.2.154
- 551 Millet, G. Y., Banfi, J. C., Kerherve, H., Morin, J. B., Vincent, L., Estrade, C., Geyssant,
552 A. & Feasson, L. (2011). Physiological and biological factors associated with a
553 24 h treadmill ultra-marathon performance: Factors associated with ultra-
554 marathon performance. *Scandinavian Journal of Medicine & Science in Sports*,
555 21(1), 54-61. doi:10.1111/j.1600-0838.2009.01001.x
- 556 Mündermann, A., Nigg, B. M., Humble, R. N., & Stefanyshyn, D. J. (2003). Foot

557 orthotics affect lower extremity kinematics and kinetics during running. *Clinical*
558 *Biomechanics*, 18(3), 254-262. doi:10.1016/S0268-0033(02)00186-9

559 Navarro, D., Zahonero, J., Huertas, F., Vera, P., & Barrios, C. (2012). *Efecto de las*
560 *plantillas podológicas de acomodación selectiva en la presión plantar de*
561 *ciclistas profesionales. Estudio Preliminar*. Presented at I Simposio sobre
562 Biomecánica y pie durante la Actividad Física, Santander.

563 Oh, Y. K., Kreinbrink, J. L., Wojtys, E. M., & Ashton-Miller, J. A. (2012). Effect of
564 axial tibial torque direction on ACL relative strain and strain rate in an in vitro
565 simulated pivot landing. *Journal of Orthopaedic Research*, 30(4), 528-534.
566 doi:10.1002/jor.21572

567 Oliveira, F. P. M., Sousa, A., Santos, R., & Tavares, J. M. R. S. (2012). Towards an
568 efficient and robust foot classification from pedobarographic images. *Computer*
569 *Methods in Biomechanics and Biomedical Engineering*, 15(11), 1181-1188.
570 doi:10.1080/10255842.2011.581239

571 Park, S. K., Lam, W. K., Yoon, S., Lee, K. K., & Ryu, J. (2017). Effects of forefoot
572 bending stiffness of badminton shoes on agility, comfort perception and lower
573 leg kinematics during typical badminton movements. *Sports Biomechanics*,
574 16(3), 374-386. doi:10.1080/14763141.2017.1321037

575 Patrick, K., & Donovan, L. (2018). Test–retest reliability of the Tekscan® F-Scan® 7 in-
576 shoe plantar pressure system during treadmill walking in healthy recreationally
577 active individuals. *Sports Biomechanics*, 17(1), 83-97.
578 doi:10.1080/14763141.2017.1355010

579 Peters, M. (1988). Footedness: Asymmetries in foot preference and skill and
580 neuropsychological assessment of foot movement. *Psychological Bulletin*,
581 103(2), 179-192. doi:10.1037/0033-2909.103.2.179

582 Phomsoupha, M., & Laffaye, G. (2014). The Science of badminton: game characteristics,
583 anthropometry, visual fitness and biomechanics. *Sports Medicine*, 11, 1-45.
584 doi:10.1007/s40279-014-0287-2

585 Powers, C. M. (2003). The influence of altered lower-extremity kinematics on
586 patellofemoral joint dysfunction: a theoretical perspective. *Journal of*
587 *Orthopaedic & Sports Physical Therapy*, 33(11), 639–646.
588 doi:10.2519/jospt.2003.33.11.639

589 Raissi, G. R. D., Cherati, A. D. S., Mansoori, K. D., & Razi, M. D. (2009). The
590 relationship between lower extremity alignment and Medial Tibial Stress
591 Syndrome among non-professional athletes. *BMC Sports Science, Medicine and*
592 *Rehabilitation*, 1(1). doi:10.1186/1758-2555-1-11

593 Ramos, J. J., Del Castillo, M. J., Polo, C., Ramón, M., & Bosch, A. (2016). Analysis of
594 the physiological parameters of young Spanish badminton players. *Revista*
595 *Internacional de Medicina y Ciencias de la Actividad Física y del Deporte*,
596 16(61), 45-54. doi:10.15366/rimcafd2016.61.004

597 Redmond, A. C., Crosbie, J., & Ouvrier, R. A. (2006). Development and validation of a
598 novel rating system for scoring standing foot posture: The foot posture index.
599 *Clinical Biomechanics*, 21(1), 89-98. doi:10.1016/j.clinbiomech.2005.08.002

600 Reeves, J., Hume, P., Gianotti, S., Wilson, B., & Ikeda, E. (2015). A retrospective review
601 from 2006 to 2011 of lower extremity injuries in badminton in New Zealand.
602 *Sports*, 3(2), 77-86. doi:10.3390/sports3020077

603 Rozzi, S., Yuktanandana, P., & Pincivero, D. (2000). Role of fatigue on proprioception
604 and neuromuscular control. In S. Lephart & F. Fu (Eds.), *Proprioception and*
605 *Neuromuscular Control Joint Stability* (pp. 375-383). Champaign, IL: Human
606 Kinetics.

607 Saragiotto, B. T., Di Pierro, C., & Lopes, A. D. (2014). Risk factors and injury prevention
608 in elite athletes: a descriptive study of the opinions of physical therapists, doctors
609 and trainers. *Brazilian Journal of Physical Therapy*, 18(2), 137-143.
610 doi:10.1590/S1413-35552012005000147

611 Sarshin, A., Mohammadi, S., Shahrabad, H. B. P., & Sedighi, M. (2011). The effects of
612 functional fatigue on dynamic postural control of badminton players. *Biology of*
613 *Exercise*, 7(2), 25-34. doi:10.4127/jbe.2011.0047

614 Selfe, J., Janssen, J., Callaghan, M., Witvrouw, E., Sutton, C., Richards, J., Stokes,
615 M., Martin, D., Dixon, J., Hogarth, R., Baltzopoulos, V., Ritchie, E., Arden, N., &
616 Dey, P. (2016). Are there three main subgroups within the patellofemoral pain
617 population? A detailed characterisation study of 127 patients to help develop
618 targeted intervention (TIPPs). *British Journal of Sports Medicine*, 50(14), 873-
619 880. doi:10.1136/bjsports-2015-094792

620 Shariff, A. H., George, J., & Ramlan, A. A. (2009). Musculoskeletal injuries among
621 Malaysian badminton players. *Singapore Medical Journal*, 50(11), 1095.

622 Sinclair, J., & Dillon, S. (2016). Influence of run and cut manoeuvres on patellofemoral
623 kinetics and kinematics. *British Journal of Sports Medicine*, 50(22), e4.14-e4.
624 doi:10.1136/bjsports-2016-096952.21

625 Sinsurin, K., Vachalathiti, R., Srisangboriboon, S., & Richards, J. (2018). Knee joint
626 coordination during single-leg landing in different directions. *Sports*
627 *Biomechanics*, 1-13. doi:10.1080/14763141.2018.1510024

628 Stirling, L. M., Von Tscharnier, V., Fletcher, J. R., & Nigg, B. M. (2012). Quantification
629 of the manifestations of fatigue during treadmill running. *European Journal of*
630 *Sport Science*, 12(5), 418-424. doi:10.1080/17461391.2011.568632

631 Taylor, J. B., Nguyen, A.-D., Griffin, J. R., & Ford, K. R. (2018). Effects of turf and cleat

632 footwear on plantar load distributions in adolescent American football players
633 during resisted pushing. *Sports Biomechanics*, 17(2), 227-237.
634 doi:10.1080/14763141.2016.1271448

635 Thijs, Y., Van Tiggelen, D., Roosen, P., De Clercq, D., & Witvrouw, E. (2007). A
636 prospective study on gait-related intrinsic risk factors for patellofemoral pain.
637 *Clinical Journal of Sport Medicine*, 17(6), 437-445.
638 doi:10.1097/JSM.0b013e31815ac44f

639 Tibesku, C., & Pässler, H. (2005). Jumper's knee. *Sportverletzung Sportschaden*, 19(02),
640 63-71. doi:10.1055/s-2005-858141

641 Trautmann, C., Martinelli, N., & Rosenbaum, D. (2011). Foot loading characteristics
642 during three fencing-specific movements. *Journal of Sports Sciences*, 29(15),
643 1585-1592. doi:10.1080/02640414.2011.605458

644 Valdecabres, R., de Benito, A. M., Littler, G., & Richards, J. (2018). An exploration of
645 the effect of proprioceptive knee bracing on biomechanics during a badminton
646 lunge to the net, and the implications to injury mechanisms. *PeerJ*, 6(e6033), 13.
647 doi:10.7717/peerj.6033

648 Valdecabres, R., de Benito, A. M., Casal, C. A., & Pablos, C. (2017). 2015 Badminton
649 world championship: Singles final men's vs. women's behaviours. *Journal of*
650 *Human Sport and Exercise*, 12(3proc), 775-788.
651 doi:10.14198/jhse.2017.12.Proc3.01

652 Vauhnik, R., Morrissey, M. C., Rutherford, O. M., Turk, Z., Pilih, I. A., & Pohar, M.
653 (2008). Knee anterior laxity: a risk factor for traumatic knee injury among
654 sportswomen? *Knee Surgery, Sports Traumatology, Arthroscopy*, 16(9), 823-
655 833. doi:10.1007/s00167-008-0559-1

- Velotta, J., Weyer, J., Ramirez, A., Winstead, J., & Bahamonde, R. (2011). Relationship between leg dominance tests and type of task. *Portuguese Journal of Sports Sciences*, 11(Suppl.2), 1035-1038.
- Wei, Y., Liu, Y., Tian, M., & Fu, W. (2009). Effects of different footwear on the metatarsophalangeal joint during push-off in critical badminton footwork. *Journal of Medical and Biological Engineering*, 29(4), 172-176. doi:10.1080/19424280902977038
- Weist, R., Eils, E., & Rosenbaum, D. (2004). The influence of muscle fatigue on electromyogram and plantar pressure patterns as an explanation for the incidence of metatarsal stress fractures. *The American Journal of Sports Medicine*, 32(8), 1-6. doi:10.1177/0363546504265191
- Whyte, E., Burke, A., White, E., & Moran, K. (2015). A high-intensity, intermittent exercise protocol and dynamic postural control in men and women. *Journal of Athletic Training*, 50(4), 392-399. doi:10.4085/1062-6050-49.6.08

Supporting information

- S1 Fig. Non-dominant and dominant lunge to the net movement
- S2 Fig. Net and shuttlecock position during test
- S3 Fig. Pressure zone distribution and sensor configuration
- S1 Table. Peak pressure in kilopascals (Kpa) for lead foot
- S2 Table. 2 Mean pressure in kilopascals (Kpa) for lead foot
- S3 Table. Peak pressure in kilopascals (Kpa) for trail foot
- S4 Table. Mean pressure in kilopascals (Kpa) for foot trail foot

Table 1 Peak pressure in kilopascals (Kpa) for lead foot

Parameter	LD task		LND task		<i>p</i> -value Interaction	<i>p</i> -value Pre - Post	η _p ²	<i>p</i> -value Lunge Dom	η _p ²
	Mean (Kpa) ± SD		Mean (Kpa) ± SD						
	Pre	Post	Pre	Post					
Hallux	2450.9±901.6	2520.6±1127.4	2702.4±570.4	2315.6±1117.1	0.170	0.460	0.05	0.913	0.00
2 nd 3 rd Phal	2503.1±818.5	2257.9±1328.3	2584.6±793.7	2269.5±1150.3	0.731	0.284	0.10	0.716	0.01
4 th 5 th Phal *	1698.5±1190.7	851.6± 1022.5	1548.3±993.5	1206.±1132.4	0.268	0.035	0.32	0.457	0.05
1 st Met	1887.9±911.9	1966.7±1006.0	2209.0±829.7	2135.30±841.1	0.475	0.991	0.000	0.058	0.27
2 nd 3 rd Mets ‡	1616.1±779.3	1858.1±919.5	1981.0±922.8	1435,6±742.2	0.011	0.595	0.02	0.799	0.01
4 th 5 th Mets	1053.7± 1076.9	1198.1±1283.7	1261.9±1028.6	1212.74±1215.7	0.301	0.857	0.00	0.317	0.08
Med-midfoot	1190.5±1034.4	1318.5± 1170.2	1058.8±852.4	1386.3±944.6	0.569	0.393	0.06	0.796	0.01
Lat-midfoot	1548.9±1034.9	1423.0±1159.5	1290.0±843.5	1660.2±1295.1	0.146	0.618	0.02	0.893	0.00
Calcaneus	1960.4±931.5	1770.9±916.8	2043.9±797.5	1827.2±968.6	0.893	0.306	0.09	0.309	0.09

* significant difference between Pre and Post † significant difference between Lunge Dominance

‡ significant interaction between Pre and post and Lunge Dominance

Table 2 Mean pressure in kilopascals (Kpa) for lead foot

Parameter	LD task		LND task		<i>p</i> -value Interaction	<i>p</i> -value Pre - Post	ηp ²	<i>p</i> -value Lunge Dom	ηp ²
	Mean (Kpa) ± SD		Mean (Kpa) ± SD						
	Pre	Post	Pre	Post					
Hallux	867.8± 507.2	790.8± 490.5	783.9± 297.1	719.6± 517.1	0.920	0.432	0.05	0.303	0.09
2 nd 3 rd Phal	636.1± 518.1	541.19± 397.2	615.0± 391.25	532.5± 374.6	0.883	0.380	0.07	0.736	0.01
4 th 5 th Phal *	966.4± 767.5	475.49± 613.9	956.9± 706.13	643.0± 609.1	0.496	0.031	0.33	0.394	0.06
1 st Met	670.8± 350.2	612.53± 355.6	750.2± 381.58	621.3± 310.8	0.354	0.331	0.08	0.374	0.07
2 nd 3 rd Mets	397.2± 206.4	361.56± 193.8	405.5± 159.04	305.8± 168.1	0.220	0.176	0.15	0.459	0.05
4 th 5 th Mets	227.8± 193.1	182.28± 168.7	256.5± 193.29	193.4± 185.3	0.667	0.225	0.12	0.419	0.06
Med-midfoot	298.1± 245.9	317.93± 281.9	258.9± 161.69	320.3± 198.3	0.645	0.479	0.04	0.516	0.04
Lat-midfoot	267.8± 180.2	233.94± 195.3	254.7± 185.42	266.6± 202.5	0.249	0.797	0.01	0.638	0.02
Calcaneus †	635.2± 389.9	538.08± 276.8	661.6± 386.04	596.2± 319.4	0.532	0.327	0.08	0.032	0.33

* significant difference between Pre and Post † significant difference between Lunge Dominance

‡ significant interaction between Pre and post and Lunge Dominance

Table 3 Peak pressure in kilopascals (Kpa) for trail foot

Parameter	LD task		LND task		<i>p</i> -value Interaction	<i>p</i> -value Pre - Post	η _p ²	<i>p</i> -value Lunge Dom	η _p ²
	Mean (Kpa) ± SD		Mean (Kpa) ± SD						
	Pre	Post	Pre	Post					
Hallux1 †	2347.9±806.9	1984.4±1161.8	2116.3±840.5	1619.1±1052.7	0.671	0.083	0.23	0.008	0.45
2 nd 3 rd Phal ‡	1749.7±965.0	1347.5±837.0	1250.5±837.0	1535.8±1051.3	0.017	0.460	0.05	0.615	0.02
4 th 5 th Phal †	1040.5±851.5	764.2± 745.7	673.2± 723.9	492.6± 588.0	0.463	0.198	0.13	0.026	0.35
1 st Met *	2382.0±915.2	1948.0±1036.1	2211.8±1070.1	1870.6±874.2	0.757	0.048	0.29	0.473	0.04
2 nd 3 rd Mets	1365.3±993.5	1123.2±653.8	1413.7±1034.5	1203.5±755.8	0.863	0.204	0.13	0.542	0.03
4 th 5 th Mets	1627.2±1038.0	1265.5±1082.4	1336.8±1179.0	1102.1±986.1	0.575	0.084	0.23	0.106	0.20
Med-midfoot *	576.3± 371.3	967.2± 608.7	595.6± 558.2	998.0± 726.4	0.966	0.022	0.37	0.837	0.00
Lat-midfoot	910.6± 916.8	875.0± 855.3	1146.9±1220.1	735.1± 692.4	0.191	0.288	0.09	0.702	0.01
Calcaneus	638.4± 415.3	456.1± 279.2	583.5± 332.7	563.4± 376.0	0.095	0.167	0.15	0.659	0.02

* significant difference between Pre and Post † significant difference between Lunge Dominance

‡ significant interaction between Pre and post and Lunge Dominance

Table 4 Mean pressure in kilopascals (Kpa) for foot trail foot

Parameter	LD task		LND task		<i>p</i> -value Interaction	<i>p</i> -value Pre - Post	η_p^2	<i>p</i> -value Lunge Dom	η_p^2
	Mean (Kpa) $\pm$ SD		Mean (Kpa) $\pm$ SD						
	Pre	Post	Pre	Post					
Hallux * †	649.8 $\pm$ 318.4	488.5 $\pm$ 302.2	560.85 $\pm$ 245.9	425.06 $\pm$ 329.0	0.811	0.020	0.38	0.038	0.31
2 nd 3 rd Phal	336.8 $\pm$ 219.9	221.7 $\pm$ 159.3	291.76 $\pm$ 239.0	254.67 $\pm$ 190.2	0.101	0.067	0.25	0.806	0.01
4 th 5 th Phal †	564.6 $\pm$ 453.3	408.7 $\pm$ 383.6	376.04 $\pm$ 395.08	280.43 $\pm$ 312.5	0.349	0.175	0.15	0.017	0.39
1 st Meta *	804.1 $\pm$ 439.2	645.1 $\pm$ 501.3	721.71 $\pm$ 353.45	576.00 $\pm$ 286.0	0.869	0.046	0.29	0.312	0.09
2 nd 3 rd Meta	354.8 $\pm$ 233.7	284.2 $\pm$ 161.2	406.64 $\pm$ 240.6	323.44 $\pm$ 172.7	0.711	0.113	0.20	0.120	0.19
4 th 5 th Meta	300.9 $\pm$ 192.7	271.3 $\pm$ 264.2	246.46 $\pm$ 163.32	223.26 $\pm$ 198.4	0.791	0.449	0.05	0.130	0.18
Med-midfoot *	155.3 $\pm$ 79.8	225.7 $\pm$ 122.5	138.82 $\pm$ 103.71	216.72 $\pm$ 147.7	0.889	0.046	0.29	0.623	0.02
Lat-midfoot	174.5 $\pm$ 135.5	162.9 $\pm$ 146.5	195.1 $\pm$ 170.17	128.49 $\pm$ 114.1	0.138	0.177	0.15	0.619	0.02
Calcaneus	229.0 $\pm$ 139.1	208.0 $\pm$ 154.1	236.27 $\pm$ 150.6	237.8 $\pm$ 182.8	0.378	0.813	0.01	0.372	0.07

* significant difference between Pre and Post † significant difference between Lunge Dominance

‡ Significant interaction between Pre and post and Lunge Dominance




Figure 1. Non-dominant and dominant lunge to the net movement



Figure 2. Net and shuttlecock position during test




Figure 3. Pressure zone distribution and sensor configuration