

Central Lancashire Online Knowledge (CLOK)

Title	Can orthotic wedges change the lower-extremity and multi-segment foot kinematics during gait in people with plantar fasciitis?
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/43361/
DOI	https://doi.org/10.1016/j.gaitpost.2022.07.259
Date	2022
Citation	Harutaichun, Pavinee, Vongsirinavarat, Mantana, Pakpakorn, Phrattaya, Sathianpantarit, Paiboon and Richards, James (2022) Can orthotic wedges change the lower-extremity and multi-segment foot kinematics during gait in people with plantar fasciitis? Gait & Posture.
Creators	Harutaichun, Pavinee, Vongsirinavarat, Mantana, Pakpakorn, Phrattaya, Sathianpantarit, Paiboon and Richards, James

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.gaitpost.2022.07.259>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

1 Can orthotic wedges change the lower-extremity and multi-segment foot kinematics during
2 gait in people with plantar fasciitis?

3
4 Pavinee Harutaichun, PhD¹

5 Mantana Vongsirinavarat, PhD¹

6 Phrattaya Pakpakorn, BsC²

7 Paiboon Sathianpantarit, BsC²

8 Jim Richards, PhD³

9
10 ¹Faculty of Physical Therapy, Mahidol University, Nakhon Pathom, Thailand

11 ²Physical Therapy Center, Mahidol University, Nakhon Pathom, Thailand

12 ³Faculty of Allied Health and Well-being, University of Central Lancashire, Preston, United
13 Kingdom

14
15 Address correspondence to Pavinee Harutaichun, 999 Phuttamonthon 4 Road, Salaya,
16 Nakhon Pathom, Thailand. E-mail: pavinee.har@mahidol.ac.th

17
18 The study protocol was approved by the center of Ethical Reinforcement for Human
19 Research of Mahidol University (COA No. MU-CIRB 2020/178.0511). The authors certify
20 that they have no affiliations with or financial involvement in any organization or entity with
21 a direct financial interest in the subject matter or materials discussed in the article. There is
22 no conflict of interest to declare.

23
24 Word count of the abstract portion: 297 words

25 Word count of the text portion: 3071 words

26
27 **AUTHOR CONTRIBUTIONS:** All authors agreed to be accountable for all aspects of the
28 work, critically revised the article for important intellectual content, and approved the final
29 manuscript to be published. Dr Harutaichun contributed to the conception and design of the
30 study, acquisition of data, analysis and interpretation of data as well as writing of manuscript.
31 Dr Vongsirinavarat contributed to the conception and design of the study, analysis and
32 interpretation of data as well as editing of manuscript. Mr Pakpakorn and Mr Sathianpantarit
33 contributed to the acquisition of data and review of manuscript. Dr Richards analyzed and
34 interpreted data as well as editing of manuscript.

35
36 **DATA SHARING:** All data relevant to the study are included in the article or are available as
37 supplementary files. Please ensure that no patient-identifiable data are available.

38
39 **PATIENT AND PUBLIC INVOLVEMENT:** Participants were not involved in the design of
40 the study or in the interpretation or translation of the study findings.

41
42 **ACKNOWLEDGEMENTS:** This research project was supported by Mahidol University. The
43 authors would like to thank Mr Preecha Romsaisiri for the materials of customized foot
44 orthoses and Mr Vasapol Teravanapanth for Mathlab analysis.

Can orthotic wedges change the lower-extremity and multi-segment foot kinematics during gait in people with plantar fasciitis?

ABSTRACT

Background: Orthotic wedges with medial posting of the forefoot and rearfoot have been shown to be effective in controlling excessive foot pronation in people with plantar fasciitis (PF), however the best prescription remains unclear.

Research question: The aim of this study was to determine the biomechanical effects of two designs of orthotic wedges with the shoe on the hip, knee, rearfoot, and forefoot kinematics in individuals with PF.

Methods: Thirty-five participants with PF were recruited. They were asked to walk under three randomized conditions; shod, shod with orthotic wedges with foot assessment technique 1 (W1), and shod with orthotic wedges from a new assessment technique (W2).

Biomechanical outcomes included lower limb and multi-segment foot kinematics in each subphase of the stance gait, including contact phase, midstance phase, and propulsive phase.

Results: Compared with shod, the W1 significantly increased shoe motion of rearfoot dorsiflexion, decreased shoe motions of peak forefoot dorsiflexion, and peak rearfoot eversion during the contact phase. In addition, W1 increased shoe motion of rearfoot inversion, decreased shoe motions of hallux dorsiflexion, and peak hallux dorsiflexion during the propulsive phase. For W2, the wedge significantly decreased peak knee internal rotation, decreased shoe motions of forefoot abduction, peak forefoot dorsiflexion, and peak rearfoot eversion during the contact phase. In addition, W2 increased rearfoot inversion, decreased hallux dorsiflexion, and decreased peak hallux dorsiflexion during the propulsive phase. When comparing W1 and W2, W1 showed greater shoe motion of rearfoot dorsiflexion during the contact phase.

Significance: These findings suggest that the use of forefoot varus wedges, and the combination of forefoot and rearfoot varus wedges, can change the lower limb kinematics, the shoe motions of multi-segment foot kinematics, and the relative length of the plantar fascia which would be associated with a reduction in pain and symptoms during walking.

Keywords: Plantar fasciitis, Kinematics, Gait, Prescription, Foot Orthotics

INTRODUCTION

Plantar fasciitis (PF) is an overuse syndrome that affects up to 15% of all adult foot complaints[1]. This condition affects the tissue under the medial longitudinal arch of the foot, causing a stabbing pain in the heel[1]. PF usually resolves within 6 to 18 months without treatment, however recovery from PF can be a slow process[2], and the nature of pain from PF can lead to a reduction in daily and sporting activities[3].

Different treatment modalities have been used in the management of PF from conservative treatments to surgery[3], with foot orthotics being the most common primary intervention[1, 3], with a view to reduce tension in the plantar fascia by decreasing over pronation during gait[4, 5]. A systematic review and meta-analysis studied different types of foot orthotics, including a medial rearfoot wedge, a medial forefoot wedge as well as a combination of medial rearfoot and forefoot wedges, in people with flexible pes planovalgus[6]. No significant differences in any outcomes were reported when comparing the medial rearfoot wedge condition and the control condition. In contrast, both the forefoot and the combination wedge decreased peak rearfoot eversion when compared with the control condition, which appears to be an effective control of excessive foot pronation during stance phase[6], however the prescription of foot orthotics in people with PF remains unclear, and

there is a lack of information on the kinematic changes in different types of foot orthotics used in the management of people with PF.

It has been suggested that the rearfoot and forefoot angles should be examined to prescribe the appropriate amount of correction needed[6-10]. Two different foot assessment techniques have been reported[9, 11]. The assessment following Root[11] is widely used by podiatrists to determine the rearfoot and forefoot angles before customizing foot orthotics[6, 12, 13], which considers an intrinsic reference frame that is relative to the proximal segment to determine the clinical forefoot and rearfoot angles. However, previous studies found poor correlation between the clinical rearfoot angle and rearfoot kinematics during the stance phase of gait[9, 14]. These findings indicate that the rearfoot and forefoot angles derived from this technique did not reflect the interaction between the foot and the ground[15], and showed poor inter-rater reliability which might be attributed to the technique of finding the subtalar neutral position[16]. The foot assessment proposed by Monaghan et al. was developed to reduce these limitations by using an extrinsic reference frame to determine the rearfoot and forefoot angles[9]. The clinical forefoot angle was defined as the angle between a line through the metatarsal head and the caudal edge of the table, which is parallel to a mediolateral axis of the foot and parallel to the ground when standing. Additionally, the clinical rearfoot angle was defined as the angle between a bisecting line of the calcaneus and a line perpendicular to the caudal edge of the table. It has been suggested that such an extrinsic clinical measure would better predict the rearfoot and forefoot angles and provide a better foot assessment to determine the amount of posting required[7, 9], however, there is a lack of information regarding the kinematic comparisons of foot orthotics in the management of people with PF.

This study considered two orthotic wedge designs 1 (W1) and 2 (W2), W1 following the foot assessment described by Root and W2 following the assessment by Monaghan et al.

Both designs used the same orthotic wedges but used different techniques for the foot assessment. To assign the orthotic wedges for each participant, those with rearfoot and forefoot angles between 3 and 6 degrees received the 3-degree wedge, between 6 and 8 degrees received the 6-degree wedge, and more than 8 degrees received the 8-degree wedge. Therefore, the aim of this study was to determine the biomechanical effects of two designs of orthotic wedges on the hip, knee, rearfoot, and forefoot kinematics in individuals with PF.

METHODS

This was a within-subject, randomized, cross-over design to determine the biomechanical effects between three conditions; shod, shod with W1, and shod with W2. The research protocols were approved by the center of Ethical Reinforcement for Human Research of Mahidol University (COA No. MU-CIRB 2020/178.0511).

The sample size was calculated using G*power version 3.1.9.4[17] for repeated-measures ANOVA. Twenty-seven participants were required to determine a 5% significance level to detect biomechanical differences between the three conditions with a medium effect (Cohen's $d = 0.5$) at a power of 80%.

The participants were diagnosed with PF by an experienced physical therapist using the following criteria: pain in the proximal attachment of the plantar fascia at the medial tubercle of the calcaneus, sharp or dull deep pain, first-step pain in the morning or during the day after prolong sitting which reduced after a few steps of walking, pain during the day after prolong walking or standing, and pain during barefoot walking and going upstairs[3]. Inclusion criteria were people between 18 and 60 years old who met the diagnosis of PF, with at least 6 weeks of symptoms[18], and an average pain intensity during the last week of at least 30-mm on a 100-mm visual analog scale (VAS). Exclusion criteria were: BMI more

than 30 kg/m², a leg length difference more than 1 cm, a positive sciatica test, history of lower-extremity fracture, or diagnosed with any systematic diseases. All participants provided written informed consent prior to data collection.

Physical assessments

Physical characteristics of the participants were assessed by a physical therapist with 7 years experience of using foot orthotics for the management of musculoskeletal problems. These included; femoral anteversion angle, tibial torsion angle, ankle inversion angle, ankle eversion angle, rearfoot angle, and forefoot angle.

Orthotic wedges

Rearfoot and forefoot angles were assessed using two techniques described by Root[11] and Monaghan et al[9] (Figure 1). Full length orthotics using a 3-mm soft foam layer (Figure 2A) were provided to each participant by the physical therapist. Orthotic wedges were made from solid rubber with a thin fabric cover, which were available in small, medium and large sizes, according to the foot length of participants (Figure 2B). Previous studies recommended the posting at 60% of the measured forefoot angle, up to a maximum of 8 degrees, for extrinsic forefoot varus wedge and the posting at 50% of the measured forefoot angle, up to a maximum of 6 degrees, for extrinsic rearfoot varus wedge, following the technique from Root[8, 10]. Regarding the amount of posting from the method introduced by Monaghan et al, the forefoot was posted at 50% of the measured forefoot angle, and the rearfoot was posted at 20% of the measured rearfoot angle[7].

Gait assessment

A 10 camera three-dimensional motion analysis system (Vicon, Vantage V5 series, Oxford, UK) was used to track the lower-extremity kinematics and the shoe motions of multi-segment foot kinematics during gait at a sampling rate of 100 Hz. The cameras were

synchronized with two force plates (AMTI, model OR6-7, USA), sampling at 1000 Hz positioned within an 8-m walkway. Forty-two retro-reflective markers were attached to the participants by the same physical therapist following the Plug-In-Gait (PIG) model and the Oxford Foot Model (OFM) with the markers applied to the shoes (Figure 3)[19]. Ten sizes of the shoe were available to ensure participants were assigned the correct size. No markers were removed from the shoes during testing. Participants were asked to walk under three conditions with the same shoe; shod, shod with W1, and shod with W2, the order of which was randomized. Before data collection in each condition, the therapist checked the location of the markers on the shoe and the shoelaces were tightened with a similar tension. The participants were asked to walk for approximately one minute to familiarize themselves with each condition. Data were collected for 3-5 successful gait trials per condition at a self-selected speed. A successful gait trial was defined as the foot making contact with the force plate with no part of the foot being over the edge of the plate. The comfort level was assessed after walking in each condition ranging from 0 to 10, with the higher score representing greater comfort. Intra-rater reliability from the gait assessment showed the ICC(3,1) values ranging from 0.75 to 0.96 and the SEM values ranged from 0.01 to 1.56.

Data processing

The kinematic and kinetic data were filtered using the 4th order zero-lag, low-pass Butterworth technique with cut off frequencies of 6 Hz and 30 Hz, respectively. Joint kinematics were tracked using Nexus (version 2.8.1) to determine the pelvis, hip, knee, rearfoot (hindfoot relative to tibia), forefoot (forefoot relative to hindfoot) in all three planes of motion, and the hallux motion in sagittal plane. Initial contact and toe-off events of each foot were identified using the vertical ground reaction force (GRF) data using a 10 N threshold. The stance phase of each foot was then normalized over a gait cycle by using the custom MATLAB software (R2017a). Peak angle and range of motion of each joint were

determined within each subphase of the stance, including contact phase, midstance phase, and propulsive phase. Contact phase was defined as the time from ipsilateral heel strike to contralateral heel off; midstance phase was defined as the time from contralateral heel off to contralateral heel contact; and propulsive phase was defined as the time from contralateral heel contact to ipsilateral toe off[20]. In addition, an approximately relative length of the plantar fascia which was distance from the 1st metatarsophalangeal joint marker to the medial calcaneus marker[21, 22], peak anteroposterior, mediolateral, and vertical GRFs were also investigated.

Statistical analysis

Shapiro–Wilk tests were used to determine if the data were normally distributed. For the normally distributed data the kinematic and kinetic characteristics from the symptomatic limbs were shown as mean $\pm$ standard deviation (SD). Repeated Measures ANOVA (RM ANOVA) were used to compare the peak angle and the range of motion of the lower extremity and multi-segment foot as well as the ground reaction force data during gait between the three conditions; shod, shod with W1, and shod with W2. Where a significant main effect was seen post-hoc pairwise comparison test with a Bonferroni correction were performed. For the non-normally distributed data kinematic and kinetic characteristics were shown as median (25th/75th Percentiles) and non-parametric Friedman tests and post hoc Wilcoxon signed-rank test were used. All statistical analyses were performed using SPSS software version 22.0 (IBM Statistics, USA), with a statistical significance level set at $P<0.05$. Effect sizes using Cohen's d were calculated for all variables[23].

RESULTS

Thirty-five participants with PF (26 females and 9 males), with a total of 41 symptomatic limbs, with an average age of 40.14 years (SD 10.53) and body mass index (BMI) of 26.35 kg/m² (SD 5.65) were included in the analysis. The symptomatic limbs showed higher forefoot varus angles when compared with previously reported normative values[8, 24, 25]. Regarding the comfort scores, there was significant difference among three conditions ($P=0.009$), the W1 showed great comfort than the shoe ($P=0.003$), but there were no significant differences between the W2 and shoe ($P=0.100$) as well as the W1 and W2 ($P=0.666$) (Table 1).

Significant differences were seen in the lower-extremity kinematics and the shoe motions of multi-segment foot kinematics between the three conditions. These included hip internal rotation ($P=0.037$), knee adduction ($P=0.039$), forefoot inversion ($P=0.035$), forefoot abduction ($P=0.011$), rearfoot dorsiflexion ($P=0.008$), hallux eversion ($P<0.001$), and relative length of the plantar fascia ($P=0.001$) during the contact phase, with the midstance phase showing differences in pelvis abduction ($P=0.033$) and rearfoot inversion ($P=0.023$). During the propulsive phase significant differences were seen in the rearfoot inversion and hallux dorsiflexion ($P=0.001$ and $P<0.001$), respectively (Table 2, Figure 4). No significant differences were seen in the GRF among three conditions ($P\geq 0.05$).

Further pairwise comparisons showed that compared with shod, the W1 significantly increased rearfoot dorsiflexion ($P=0.035$), decreased peak forefoot dorsiflexion ($P=0.011$), decreased peak rearfoot eversion ($P=0.035$), and decreased relative length of the plantar fascia ($P=0.029$) during the contact phase. In addition, W1 increased rearfoot inversion ($P=0.009$), decreased hallux dorsiflexion ($P=0.010$), and decreased peak hallux dorsiflexion ($P=0.006$) during the propulsive phase. For W2, the wedge significantly decreased forefoot abduction ($P=0.032$), decreased peak forefoot dorsiflexion ($P=0.001$), decreased peak rearfoot eversion ($P=0.001$), decreased peak knee internal rotation ($P=0.033$), and decreased

relative length of the plantar fascia ($P=0.009$) during the contact phase. In addition, W2 increased rearfoot inversion ($P=0.002$), decreased hallux dorsiflexion ($P<0.001$), and decreased peak hallux dorsiflexion ($P<0.001$) during the propulsive phase. When comparing W1 and W2, W1 showed greater rearfoot dorsiflexion ($P=0.005$) during contact phase, Table 3 and Table 4.

In the non-symptomatic sides, the orthotic wedge significantly decreased peak hallux dorsiflexion ($P<0.001$) during the propulsive phase when compared with the shod condition, which was similar to the response on the symptomatic sides. There was no significant difference between the orthotic wedges and the shod condition for the peak angle or the other segments and the relative length of the plantar fascia ($P\geq 0.05$), as shown in Table 5.

DISCUSSION

The aim of this study was to determine the biomechanical effects of two designs of orthotic wedges in people with PF. The results confirmed that both designs of orthotic wedges produced significant biomechanical changes when compared with shod walking during the contact and propulsive phases. Regarding the contact phase, both W1 and W2 decreased shoe motions of peak forefoot dorsiflexion and peak rearfoot eversion, as well as decreased relative length of the plantar fascia. These results imply that orthotic wedges reduce dynamic foot pronation during the contact phase, since greater rearfoot eversion and forefoot dorsiflexion are included in the components of over pronation during gait, which could then induce a greater strain in the plantar fascia [4, 5, 26]. The present findings, therefore, support the use of orthotic wedges to decrease excessive elongation in the plantar fascia during gait in individuals with PF. Such effect was found in only the symptomatic

sides, with no significant difference seen in the relative length of the plantar fascia in the non-symptomatic sides between the orthotic wedge condition and the shod condition.

In addition, both W1 and W2 produced more shoe motion of rearfoot inversion, less shoe motions of hallux dorsiflexion, and peak hallux dorsiflexion during the propulsive phase. It was thus assumed from the present findings that the use of orthotic wedges could produce earlier inversion of the foot than the shod only condition. Although there were no significant differences between conditions during the midstance phase, the greater rearfoot inversion and less hallux dorsiflexion when using the orthotic wedges indicates greater foot stability during the propulsive phase than the shod only condition. Such biomechanical changes could reduce the excessive tension of plantar fascia and improve propulsion in people with PF[27-29].

This study also considered the biomechanical effects between two orthotic wedges designed based on two different foot assessment techniques. The two designs of orthotic wedges provided different effects when compared with the shod, with the W1 increasing shoe motion of rearfoot dorsiflexion during the contact phase; whereas the W2 decreased shoe motion of forefoot abduction and decreased peak knee internal rotation during the contact phase. Regarding the comparisons between the two designs of orthotic wedges, the W1 produced more shoe motion of rearfoot dorsiflexion, which can be associated with the different posting used within the two types of orthotic wedges. The amount of posting for the two designs was calculated from two different techniques of foot assessment[7, 8], with the technique suggested by Monaghan et al[9] providing less rearfoot posting than that from the assessment suggested by Root[11]. From the assessments the majority of the participants received posting of the W1 (Root) at both the rearfoot and forefoot while W2 (Monaghan et al.) mostly posted at the forefoot only. Only the participants with rearfoot varus in relaxed position of more than 15 degrees were provided with a W2 with posting at rearfoot[9].

Since W1 mostly included both the forefoot and rearfoot postings, it produced more shoe motion of rearfoot dorsiflexion than W2. One possible mechanism for PF has been suggested as a lack of shank-calcaneus dorsiflexion during the contact phase. Such a limitation could induce more midfoot dorsiflexion, resulting in over foot pronation and more stretch of the plantar fascia[4, 30]. W1 could thus reduce relative length of the plantar fascia by producing more dorsiflexion of the shank-calcaneus found in the early stance. Whereas, W2, which mainly included forefoot posting, showed superior effects than the W1 in the reduction of forefoot abduction. It is possible that only the forefoot posting was sufficient to decrease foot pronation during the weight acceptance by shifting the weight-bearing line from the medial to the lateral side of the foot[31]. When considering the coupling mechanism between the foot, tibia, femur, and hip, a reduction of foot pronation has been suggested to decrease internal rotation of the limb[26]. Therefore, the W2 could reduce over stretch of the plantar fascia by reducing the deviation of foot movement in the transverse plane and subsequently decreasing the prolonged internal rotation of the knee which is supported by these present findings. No difference was seen in the relative length of the plantar fascia when comparing W1 and W2, as both provided significant changes in the kinematics on the symptomatic sides during gait. Therefore, either design of the orthotic wedge could be used for individuals with PF, however further studies are required to support longer term clinical outcomes.

To our knowledge, this is the first study to determine the biomechanical effects of orthotic wedges in people with PF, and these findings are supported by a previous meta-analysis which suggested the use of forefoot varus wedge and the combination of forefoot and rearfoot varus wedges to reduce over pronation during walking[6]. However, the results from the present study provide data only on the immediate effects of the orthotic wedges and further studies should be conducted to determine the longer term clinical effects alongside the

biomechanical changes, which should also consider muscle activity within the foot and lower limb.

CONCLUSION

Both techniques of orthotic prescription indicate biomechanical changes of the lower extremity and the shoe motions of the multi-segment foot during the contact and propulsive subphases of gait. However the two orthotic wedges provided different effects with the W1 increasing shoe motion of rearfoot dorsiflexion during the contact phase, whereas the W2 decreased shoe motion of forefoot abduction and decreased peak knee internal rotation during the contact phase. These findings support the use of forefoot varus wedge and the combination of forefoot and rearfoot varus wedges to reduce the relative length of the plantar fascia which might be associated with a reduction in pain and symptoms during walking.

REFERENCES

- [1] T.G. McPoil, R.L. Martin, M.W. Cornwall, D.K. Wukich, J.J. Irrgang, J.J. Godges, Heel pain--plantar fasciitis: clinical practice guidelines linked to the international classification of function, disability, and health from the orthopaedic section of the American Physical Therapy Association, *J Orthop Sports Phys Ther* 38(4) (2008) A1-A18.
- [2] T. Forcum, T. Hyde, D. Aspegren, G. Lawson, Chiropractic Round Table. Plantar Fasciitis and Heel Pain Syndrome, *Journal of the American Chiropractic Association* 47(7) (2010) 26-33.
- [3] R.L. Martin, T.E. Davenport, S.F. Reischl, T.G. McPoil, J.W. Matheson, D.K. Wukich, C.M. McDonough, A. American Physical Therapy, Heel pain-plantar fasciitis: revision 2014, *Journal of Orthopaedic & Sports Physical Therapy* 44(11) (2014) A1-33.

336 [4] R. Chang, P.A. Rodrigues, R.E. Van Emmerik, J. Hamill, Multi-segment foot kinematics
337 and ground reaction forces during gait of individuals with plantar fasciitis, *J Biomech* 47(11)
338 (2014) 2571-7.

339 [5] P. Harutaichun, S. Boonyong, P. Pensri, Differences in lower-extremity kinematics
340 between the male military personnel with and without plantar fasciitis, *Phys Ther Sport* 50
341 (2021) 130-137.

342 [6] G. Desmyttere, M. Hajizadeh, J. Bleau, M. Begon, Effect of foot orthosis design on lower
343 limb joint kinematics and kinetics during walking in flexible pes planovalgus: A systematic
344 review and meta-analysis, *Clin Biomech (Bristol, Avon)* 59 (2018) 117-129.

345 [7] W.H. Hsu, C.L. Lewis, G.M. Monaghan, E. Saltzman, J. Hamill, K.G. Holt, Orthoses
346 posted in both the forefoot and rearfoot reduce moments and angular impulses on lower
347 extremity joints during walking, *J Biomech* 47(11) (2014) 2618-25.

348 [8] G.P. Brown, R. Donatelli, P.A. Catlin, M.J. Wooden, The effect of two types of foot
349 orthoses on rearfoot mechanics, *J Orthop Sports Phys Ther* 21(5) (1995) 258-67.

350 [9] G.M. Monaghan, C.L. Lewis, W.H. Hsu, E. Saltzman, J. Hamill, K.G. Holt, Forefoot
351 angle determines duration and amplitude of pronation during walking, *Gait Posture* 38(1)
352 (2013) 8-13.

353 [10] R.A. Donatelli, C. Hurlburt, D. Conaway, R. St Pierre, Biomechanical foot orthotics: a
354 retrospective study, *J Orthop Sports Phys Ther* 10(6) (1988) 205-12.

355 [11] M.I. Root, Biomechanical examination of the foot, *Journal of the American Podiatry*
356 *Association* 63(1) (1973) 28-9.

357 [12] C.L. MacLean, I.S. Davis, J. Hamill, Short- and long-term influences of a custom foot
358 orthotic intervention on lower extremity dynamics, *Clin J Sport Med* 18(4) (2008) 338-43.

359 [13] J.M. Genova, M.T. Gross, Effect of foot orthotics on calcaneal eversion during standing
360 and treadmill walking for subjects with abnormal pronation, *J Orthop Sports Phys Ther*
361 30(11) (2000) 664-75.

362 [14] M.R. Pierrynowski, S.B. Smith, Rear foot inversion/eversion during gait relative to the
363 subtalar joint neutral position, *Foot Ankle Int* 17(7) (1996) 406-12.

364 [15] H.L. Jarvis, C.J. Nester, P.D. Bowden, R.K. Jones, Challenging the foundations of the
365 clinical model of foot function: further evidence that the root model assessments fail to
366 appropriately classify foot function, *J Foot Ankle Res* 10 (2017) 7.

367 [16] B. Van Gheluwe, K.A. Kirby, P. Roosen, R.D. Phillips, Reliability and accuracy of
368 biomechanical measurements of the lower extremities, *J Am Podiatr Med Assoc* 92(6) (2002)
369 317-26.

370 [17] F. Faul, E. Erdfelder, A.G. Lang, A. Buchner, G*Power 3: a flexible statistical power
371 analysis program for the social, behavioral, and biomedical sciences, *Behav Res Methods*
372 39(2) (2007) 175-91.

373 [18] S.C. Wearing, J.E. Smeathers, B. Yates, P.M. Sullivan, S.R. Urry, P. Dubois, Sagittal
374 movement of the medial longitudinal arch is unchanged in plantar fasciitis, *Med Sci Sports*
375 *Exerc* 36(10) (2004) 1761-7.

376 [19] J. Stebbins, M. Harrington, N. Thompson, A. Zavatsky, T. Theologis, Repeatability of a
377 model for measuring multi-segment foot kinematics in children, *Gait Posture* 23(4) (2006)
378 401-10.

379 [20] A. De Cock, D. De Clercq, T. Willems, E. Witvrouw, Temporal characteristics of foot
380 roll-over during barefoot jogging: reference data for young adults, *Gait Posture* 21(4) (2005)
381 432-9.

382 [21] R. Ferber, B. Benson, Changes in multi-segment foot biomechanics with a heat-
383 mouldable semi-custom foot orthotic device, *J Foot Ankle Res* 4(1) (2011) 18.

384 [22] J. Sinclair, J. Isherwood, P.J. Taylor, The effects of orthotic intervention on
385 multisegment foot kinematics and plantar fascia strain in recreational runners, *J Appl*
386 *Biomech* 31(1) (2015) 28-34.

387 [23] J. Cohen, *Statistical power analysis for the behavioral sciences*, L. Erlbaum Associates,
388 Hillsdale, N.J., 1988.

389 [24] K.R. Buchanan, I. Davis, The relationship between forefoot, midfoot, and rearfoot static
390 alignment in pain-free individuals, *J Orthop Sports Phys Ther* 35(9) (2005) 559-66.

391 [25] R.A. Donatelli, Abnormal biomechanics of the foot and ankle, *J Orthop Sports Phys*
392 *Ther* 9(1) (1987) 11-6.

393 [26] V.H. Chuter, X.A. Janse de Jonge, Proximal and distal contributions to lower extremity
394 injury: a review of the literature, *Gait Posture* 36(1) (2012) 7-15.

395 [27] B.L. Costa, F.A. Magalhaes, V.L. Araujo, J. Richards, F.M. Vieira, T.R. Souza, R.
396 Trede, Is there a dose-response of medial wedge insoles on lower limb biomechanics in
397 people with pronated feet during walking and running?, *Gait Posture* 90 (2021) 190-196.

398 [28] G. Desmyttere, M. Hajizadeh, J. Bleau, S. Leteneur, M. Begon, Anti-pronator
399 components are essential to effectively alter lower-limb kinematics and kinetics in
400 individuals with flexible flatfeet, *Clin Biomech (Bristol, Avon)* 86 (2021) 105390.

401 [29] S.C. Lin, C.P. Chen, S.F. Tang, A.M. Wong, J.H. Hsieh, W.P. Chen, Changes in
402 windlass effect in response to different shoe and insole designs during walking, *Gait Posture*
403 37(2) (2013) 235-41.

404 [30] W.B. Kibler, C. Goldberg, T.J. Chandler, Functional biomechanical deficits in running
405 athletes with plantar fasciitis, *Am J Sports Med* 19(1) (1991) 66-71.

406 [31] C. MacLean, I.M. Davis, J. Hamill, Influence of a custom foot orthotic intervention on
407 lower extremity dynamics in healthy runners, *Clin Biomech (Bristol, Avon)* 21(6) (2006)
408 623-30

409 **Table 1** Participant characteristics (n=41). Data are shown as mean (SD), minimum and maximum

Characteristics	Mean (SD)	Minimum	Maximum
Femoral anteversion angle, degrees	14.67 (2.81)	9.0	21.0
Tibial torsion angle, degrees	22.51 (4.13)	13.0	35.0
Ankle inversion angle, degrees	14.66 (5.97)	7.0	31.0
Ankle eversion angle, degrees	6.32 (2.71)	3.0	15.0
Foot assessment for the orthotic wedge design 1 (W1)			
- Rearfoot varus angle, degrees	0.98 (2.34)	-4.0	7.0
- Rearfoot varus wedge, degrees	4.63 (1.80)	0	8
- Forefoot angle, degrees	11.46 (3.63)	4.0	23.0
- Forefoot varus wedge, degrees	5.32 (1.84)	0	8
Foot assessment for the orthotic wedge design 2 (W2)			
- Rearfoot angle, degrees	5.24 (3.07)	0.0	15.0
- Forefoot angle, degrees	18.05 (7.30)	6.0	33.0
- Forefoot varus wedge, degrees	6.20 (1.99)	0	8
Comfort with foot orthoses (FOs)			
- Shoe, points	5.72 (1.94)	1.65	9.80
- W1, points	6.54 (1.68)	2.65	9.05
- W2, points	6.31 (2.04)	1.45	9.68

410

411

412

413

414

415

416

417

418

419 **Table 2** Comparisons of the mean (SD) or median (IQR) of the lower-extremity range of motion among the shod condition, W1 condition, and
 420 W2 condition in each subphase of stance gait (n=41)

	Contact phase					Midstance phase					Propulsive phase				
	Mean (SD) / Median (IQR)			P	Effect size	Mean (SD) / Median (IQR)			P	Effect size	Mean (SD) / Median (IQR)			P	Effect size
	Shod	W1	W2			Shod	W1	W2			Shod	W1	W2		
Pelvis															
Sagittal (°)	1.8 (0.8)	1.9 (0.8)	1.8 (0.7)	0.469	0.293	2.7 [1.8,3.8]	2.6 [1.7,3.5]	2.6 [1.8,3.1]	0.247	0.556	2.0 (0.9)	1.9 (1.0)	2.0 (0.9)	0.703	0.191
Frontal (°)	4.0 (1.1)	3.9 (1.1)	3.7 (1.1)	0.064	0.574	6.0 (2.0)	5.9 (1.8)	5.7 (1.7)	0.033*	0.614	4.0 [3.1,4.6]	3.8 [3.1,4.5]	3.7 [2.7,4.4]	0.540	0.366
Transverse (°)	2.2 (1.2)	2.4 (1.2)	2.4 (1.3)	0.364	0.340	7.6 (2.2)	7.5 (1.8)	7.5 (1.7)	0.824	0.155	2.2 (1.3)	2.4 (1.3)	2.3 (1.2)	0.616	0.230
Hip															
Sagittal (°)	12.2 (2.6)	12.6 (3.3)	12.7 (2.4)	0.235	0.398	29.2 (4.4)	28.8 (4.7)	28.6 (4.5)	0.239	0.392	8.8 (2.6)	6.3 (2.6)	9.3 (2.3)	0.115	0.496
Frontal (°)	7.0 (2.2)	6.8 (2.1)	6.7 (2.2)	0.273	0.381	5.5 (1.8)	5.2 (1.9)	5.2 (1.7)	0.231	0.398	10.4 (2.4)	10.1 (2.6)	10.1 (2.8)	0.151	0.449
Transverse (°)	26.5 (9.8)	27.5 (10.6)	25.6 (10.3)	0.037*	0.602	14.6 (5.7)	14.4 (5.3)	14.0 (5.4)	0.475	0.286	18.0 (9.6)	18.4 (10.6)	18.6 (10.5)	0.465	0.286
Knee															
Sagittal (°)	8.5 [6.1,11.7]	8.4 [6.3,10.8]	8.6 [5.4,11.1]	0.140	0.690	8.5 (3.3)	8.3 (3.5)	8.1 (3.4)	0.388	0.333	25.4 (8.9)	25.8 (9.5)	25.6 (9.9)	0.863	0.127
Frontal (°)	5.7 [3.4,7.4]	5.7 [3.6,7.4]	5.3 [3.3,7.4]	0.039*	0.907	5.0 (2.8)	4.9 (2.5)	5.1 (3.1)	0.723	0.191	21.6 (9.6)	22.5 (10.5)	22.5 (10.3)	0.142	0.464
Transverse (°)	15.7 (4.5)	14.9 (3.8)	14.4 (3.6)	0.118	0.510	7.9 (2.7)	7.5 (2.4)	7.5 (2.3)	0.493	0.278	9.3 (4.3)	8.9 (5.2)	8.5 (4.2)	0.359	0.333
HF-TB															
Sagittal (°)	16.1 [13.4,17.8]	17.3 [14.7,20.0]	15.6 [14.0,18.9]	0.008*	1.190	16.8 (5.0)	17.4 (4.3)	17.0 (5.2)	0.746	0.180	30.8 (7.3)	31.1 (7.5)	29.8 (6.7)	0.282	0.375
Frontal (°)	25.4 [19.6,32.9]	26.1 [20.0,31.6]	23.3 [19.5,30.7]	0.506	0.386	19.2 (7.5)	17.7 (6.6)	17.9 (6.7)	0.023*	0.648	13.3 (6.8)	15.2 (7.4)	15.3 (6.7)	0.001*	0.912
Transverse (°)	10.0 (3.8)	10.7 (3.9)	9.7 (3.6)	0.054	0.590	7.3 (3.3)	7.6 (4.2)	7.6 (3.1)	0.777	0.168	15.3 (9.3)	15.6 (10.7)	14.7 (8.1)	0.631	0.220
FF-HF															
Sagittal (°)	4.8 (1.2)	4.8 (1.3)	4.5 (1.4)	0.286	0.381	2.1 (0.8)	2.1 (0.9)	2.1 (1.0)	0.844	0.142	2.8 (1.1)	3.2 (1.6)	2.9 (1.2)	0.293	0.392
Frontal (°)	1.9 (0.6)	2.1 (0.5)	1.9 (0.5)	0.035*	0.625	1.6 (0.6)	1.5 (0.4)	1.5 (0.4)	0.801	0.168	1.6 (0.6)	1.4 (0.7)	1.4 (0.5)	0.068	0.582
Transverse (°)	4.0 (1.3)	3.6 (1.2)	3.6 (1.1)	0.011*	0.728	1.9 [1.4,2.6]	1.8 [1.1,2.4]	1.9 [1.3,2.5]	0.238	0.580	3.5 (1.6)	3.6 (1.7)	3.5 (1.3)	0.831	0.142
Hallux															
Sagittal (°)	5.2 (2.9)	5.1 (2.6)	5.2 (2.7)	0.868	0.127	6.2 (3.1)	6.5 (3.6)	6.7 (3.1)	0.311	0.381	24.9 (8.3)	20.8 (7.9)	19.1 (6.5)	<0.001*	1.223

421
 422 * Significant difference of the main effect
 423

423

424

425

426

427

428 **Table 3** Pairwise comparisons of the mean or median of the lower-extremity range of motion among the shod condition, W1 condition, and W2
 429 condition in each subphase of stance gait (n=41)

	Contact phase						Midstance phase						Propulsive phase					
	Mean difference (SE) / Z score			P *	P **	P ***	Mean difference (SE) / Z score			P *	P **	P ***	Mean difference (SE) / Z score			P *	P **	P ***
	Shod VS W1	Shod VS W2	W1 VS W2				Shod VS W1	Shod VS W2	W1 VS W2				Shod VS W1	Shod VS W2	W1 VS W2			
Pelvis																		
Sagittal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Frontal (°)	N/A	N/A	N/A	N/A	N/A	N/A	0.2 (0.1)	0.3 (0.1)	0.1 (0.1)	0.345	0.055	0.717	N/A	N/A	N/A	N/A	N/A	N/A
Transverse (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Hip																		
Sagittal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Frontal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Transverse (°)	-1.0 (0.6)	0.9 (0.8)	1.9 (0.8)	0.311	0.730	0.062	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Knee																		
Sagittal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Frontal (°)	-0.335	-1.704	-1.835	0.738	0.088	0.067	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Transverse (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
HF-TB																		
Sagittal (°)	-2.110	-0.566	-2.814	0.035	0.572	0.005	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Frontal (°)	N/A	N/A	N/A	N/A	N/A	N/A	1.5 (0.6)	1.3 (0.7)	-0.3 (0.5)	0.058	0.163	1.000	-1.9 (0.6)	-2.0 (0.5)	-0.1 (0.6)	0.009	0.002	1.000
Transverse (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
FF-HF																		
Sagittal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Frontal (°)	-0.2 (0.1)	-0.01 (0.1)	0.1 (0.1)	0.080	1.000	0.092	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Transverse (°)	0.4 (0.2)	0.5 (0.2)	0.1 (0.2)	0.077	0.032	1.000	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Hallux																		
Sagittal (°)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	4.0 (1.3)	5.8 (1.1)	1.7 (1.1)	0.010	< 0.001	0.326

430
 431 * Comparison between shod and W1 ** Comparison between shod and W2 *** Comparison between W1 and W2
 432

432

433

434

435

436

437 **Table 4** Comparisons of the mean (SD) or median (IQR) of the peak angle, arch height index, relative length of the plantar fascia, and ground
438 reaction force among the shod condition, W1 condition, and W2 condition (n=41)

	Condition			P	Effect size	P *	P **	P ***
	Shod	W1	W2					
Hip								
Peak adduction (°)	6.88 (2.99)	6.48 (3.27)	6.40 (3.18)	0.107	0.492	N/A	N/A	N/A
Peak internal rotation (°)	25.00 [11.02, 35.50]	28.16 [7.45, 37.22]	26.76 [9.09, 38.88]	0.298	0.522	N/A	N/A	N/A
Knee								
Peak adduction (°)	3.41 [-0.86, 7.16]	3.42 [-1.25, 6.35]	2.94 [-1.98, 6.13]	0.332	0.490	N/A	N/A	N/A
Peak internal rotation (°)	0.60 [-17.88, 13.11]	0.96 [-22.69, 8.73]	-0.43 [-25.09, 8.01]	0.040	0.904	0.078	0.033	0.115
HF-TB								
Peak plantarflexion (°)	-16.50 (6.99)	-15.97 (7.35)	-15.81 (7.50)	0.465	0.313	N/A	N/A	N/A
Peak eversion (°)	-28.96 [-48.37, -5.55]	-18.93 [-46.82, -6.15]	-25.19 [-43.28, -1.52]	0.005	1.211	0.035	0.001	0.068
Peak abduction (°)	-5.46 [-8.59, -2.46]	-5.58 [-9.09, -3.41]	-6.24 [-9.37, -3.02]	0.165	0.713	N/A	N/A	N/A
FF-HF								
Peak dorsiflexion (°)	10.16 (7.03)	9.69 (6.97)	9.27 (7.00)	0.001	1.553	0.011	0.001	0.101
Peak inversion (°)	0.04 [-1.50, 2.88]	0.75 [-1.38, 2.93]	0.11 [-1.96, 2.79]	0.101	0.752	N/A	N/A	N/A
Peak adduction (°)	4.00 [0.16, 8.12]	5.11 [0.64, 7.71]	5.10 [1.08, 7.62]	0.225	0.592	N/A	N/A	N/A
Hallux								
Peak dorsiflexion (°)	38.21 (13.57)	33.97 (12.80)	32.14 (11.49)	< 0.001	2.548	0.006	< 0.001	0.189
Peak inversion (°)	-15.84 (9.43)	-15.29 (9.44)	-15.60 (9.25)	0.534	0.017	N/A	N/A	N/A
Peak abduction (°)	-3.67 [-5.42, -1.23]	-3.81 [-4.87, -0.95]	-3.27 [-4.21, -0.99]	0.021	1.131	0.194	0.045	0.005
Arch								
Arch height index	1.97 (0.92)	1.80 (0.86)	1.66 (0.88)	0.156	0.483	N/A	N/A	N/A
Relative length of the plantar fascia	4.71 (1.81)	4.03 (1.47)	3.90 (1.52)	0.001	0.956	0.029	0.009	1.000
Anteroposterior GRF								
First peak	16.13 (3.99)	15.51 (3.88)	15.59 (3.78)	0.324	0.352	N/A	N/A	N/A
Second peak	21.15 (3.21)	20.62 (3.32)	20.55 (3.41)	0.159	0.444	N/A	N/A	N/A
Mediolateral GRF								
First peak	5.60 (1.38)	5.71 (1.45)	5.54 (1.53)	0.729	0.191	N/A	N/A	N/A
Second peak	4.38 (1.92)	4.47 (2.07)	4.03 (1.90)	0.057	0.565	N/A	N/A	N/A
Vertical GRF								
First peak	108.87 (8.85)	108.46 (9.53)	108.41 (9.30)	0.885	0.110	N/A	N/A	N/A
Second peak	105.24 (7.89)	105.24 (8.03)	104.65 (7.89)	0.808	0.155	N/A	N/A	N/A

439
440 * Comparison between shod and W1 ** Comparison between shod and W2 *** Comparison between W1 and W2
441

442

443

444 **Table 5** Comparisons of the mean (SD) or median (IQR) of the peak angle, arch height index, relative length of the plantar fascia, and ground
445 reaction force among the shod condition, W1 condition, and W2 condition in the non symptomatic sides (n=29)


	Condition			P	Effect size	p *	p **	p ***
	Shod	W1	W2					
Hip								
Peak adduction (°)	7.70 (4.97)	7.65 (5.41)	7.78 (5.27)	0.919	0.127	N/A	N/A	N/A
Peak internal rotation (°)	19.16 [5.73, 29.25]	16.66 [5.56, 29.12]	18.01 [5.56, 32.51]	0.584	0.443	N/A	N/A	N/A
Knee								
Peak adduction (°)	3.50 [-0.38, 5.64]	2.86 [-0.32, 5.99]	3.61 [-0.55, 5.93]	0.535	0.480	N/A	N/A	N/A
Peak internal rotation (°)	1.93 [-6.54, 16.54]	-0.91 [-11.70, 13.08]	0.73 [-12.63, 11.42]	0.153	0.884	N/A	N/A	N/A
HF-TB								
Peak plantarflexion (°)	-15.59 (7.24)	-15.52 (8.77)	-14.82 (8.68)	0.723	0.247	N/A	N/A	N/A
Peak eversion (°)	-20.32 [-47.88, -8.89]	-18.27 [-36.80, -5.43]	-21.14 [-34.40, -6.83]	0.125	0.939	N/A	N/A	N/A
Peak abduction (°)	-6.98 [-10.62, 0.55]	-6.74 [-10.32, -0.57]	-6.44 [-11.33, -0.32]	0.866	0.226	N/A	N/A	N/A
FF-HF								
Peak dorsiflexion (°)	8.52 (4.17)	8.24 (6.30)	7.63 (6.34)	0.657	0.271	N/A	N/A	N/A
Peak inversion (°)	1.45 [-0.60, 4.09]	1.28 [-0.66, 3.28]	0.81 [-0.14, 3.31]	0.247	0.727	N/A	N/A	N/A
Peak adduction (°)	3.02 [-0.12, 8.00]	3.99 [0.99, 8.13]	4.53 [1.01, 8.15]	0.257	0.714	N/A	N/A	N/A
Hallux								
Peak dorsiflexion (°)	48.75 [29.85, 59.24]	43.10 [25.74, 54.32]	32.14 [30.28, 51.72]	< 0.001	2.990	0.003	0.007	0.819
Peak inversion (°)	-13.35 [-27.56, -7.11]	-14.48 [-26.69, -3.25]	-12.92 [-28.09, -6.85]	0.926	0.164	N/A	N/A	N/A
Peak abduction (°)	-4.00 [-5.40, -1.02]	-2.40 [-4.85, 0.25]	-2.76 [-5.10, -0.99]	0.123	0.920	N/A	N/A	N/A
Arch								
Arch height index	2.28 (1.10)	2.17 (1.19)	1.87 (0.95)	0.071	0.696	N/A	N/A	N/A
Relative length of the plantar fascia	4.28 (1.54)	4.10 (1.63)	3.89 (1.62)	0.519	0.387	N/A	N/A	N/A
Anteroposterior GRF								
First peak	14.85 [11.82, 18.59]	15.28 [12.21, 17.38]	15.59 [12.05, 18.31]	0.867	0.224	N/A	N/A	N/A
Second peak	20.44 (3.18)	20.13 (3.24)	19.62 (3.35)	0.020	0.883	0.686	0.065	0.194
Mediolateral GRF								
First peak	6.16 (1.98)	6.14 (1.96)	6.05 (1.99)	0.928	0.110	N/A	N/A	N/A
Second peak	4.86 (2.46)	4.83 (2.58)	4.99 (2.45)	0.864	0.168	N/A	N/A	N/A
Vertical GRF								
First peak	106.08 [101.31, 115.18]	107.38 [100.35, 113.68]	106.10 [101.63, 114.44]	0.108	0.953	N/A	N/A	N/A
Second peak	104.57 [98.43, 110.97]	106.58 [97.13, 111.78]	103.21 [99.29, 111.76]	0.872	0.215	N/A	N/A	N/A

446
447 * Comparison between shod and W1 ** Comparison between shod and W2 *** Comparison between W1 and W2
448

449

450


Figure 1 The foot assessment technique 1 (W1: Right side) and the foot assessment technique 2 (W2: Left side). A represents the rearfoot angle formed between a bisection line at distal one third of lower leg and a bisection line at calcaneus in subtalar neutral position. B represents the forefoot angle formed between a bisection line at calcaneus and a parallel line through the metatarsal heads in subtalar neutral position. C represents the rearfoot angle formed between a bisection line at calcaneus and a line perpendicular to the caudal edge of the table. D represents the angle formed between a line through the metatarsal head and a line parallel to the caudal edge of the table.



465

466 **Figure 2** Orthotic wedges in the present study (A: orthotic wedges with a full length of soft foam layer, B: medial forefoot and rearfoot varus
467 wedge with three different sizes i.e. small (S), medium (M), large (L). Blue color is the 3-degree wedge, Yellow color is the 6-degree wedge,
468 Red color is the 8-degree wedge)

469



478

479

480

481

482

483


484

485


Figure 3 Marker placement of the lower extremity and multi-segment foot (A: lateral view, B: anterior view, C: posterior view, D: anterior view of barefoot)




Figure 4 Comparisons of the lower-extremity kinematics and the vertical GRF among the shod condition, W1 condition, and W2 condition in each subphase of stance gait (n=41) (Shod represented by a dashed line, W1 represented by a straighted line, and W2 represented by a dotted line)




525



526




527




529


530



531




532




533

534

535



536



537

538

539

540

541