

Central Lancashire Online Knowledge (CLoK)

Title	The associations between digit ratio (2D:4D and right – left 2D:4D), maximal oxygen consumption and ventilatory thresholds in professional male football players
Type	Article
URL	https://clok.uclan.ac.uk/id/eprint/50580/
DOI	https://doi.org/10.1002/ajhb.24047
Date	2024
Citation	Parpa, Koulla, Manning, John T., Kobus, Magdalena, Mason, Laura and Michaelides, Marcos (2024) The associations between digit ratio (2D:4D and right – left 2D:4D), maximal oxygen consumption and ventilatory thresholds in professional male football players. American Journal of Human Biology, 36 (6). ISSN 1042-0533
Creators	Parpa, Koulla, Manning, John T., Kobus, Magdalena, Mason, Laura and Michaelides, Marcos

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1002/ajhb.24047

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the http://clok.uclan.ac.uk/policies/

1	Title: The associations between digit ratio (2D:4D and right - left 2D:4D), maximal oxygen
2	consumption and ventilatory thresholds in professional male football players
3	
4 5	Authors: Koulla Parpa ^{1*} , John T. Manning ² , Magdalena Kobus ^{3,4} , Laura Mason ² , Marcos Michaelides ¹ .
6 7	 University of Central Lancashire - Cyprus Campus, University Avenue 12-14, 7080, Pyla, Cyprus
8 9 10	2. Applied Sports, Technology, Exercise, and Medicine (A-STEM), Swansea University, Swansea, UK
11 12	3. Department of Anthropology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, 90-237, Poland
13 14 15 16	4. Institute of Biological Sciences, Faculty of Biology and Environmental Sciences, Cardinal Stefan Wyszynski University in Warsaw, 01-938 Warsaw, Poland
17	Assistant Professor *Koulla Parpa (corresponding author)
18	kparpa@uclan.ac.uk
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	

Abstract

29

Introduction: Digit ratio (2D:4D: the relative length of the 2nd and 4th digit) is thought to be a 30 negative correlate of prenatal testosterone. The 2D:4D is related to oxygen metabolism, but 31 the precise nature of this relationship is unclear. The purpose of the present study was to 32 consider associations between digit ratios (right 2D:4D, left 2D:4D, right-left 2D:4D [Dr-1]) 33 and VO_{2max} and ventilatory thresholds (VT1 and VT2). Methods: One hundred and thirty-34 35 three Caucasian (n=133) professional football players competing in Cyprus participated in the study. Players underwent anthropometric measurements, and digit lengths were measured 36 from hand scans. They also completed an incremental cardiopulmonary test to exhaustion on 37 a treadmill. Results: There were negative correlations between digit ratios and VO_{2max} (right 38 2D:4D, r = -.65; left 2D:4D r = -.37, both p < .0001; Dr-1 r = -.30, p = .0005). There were no 39 relationships between digit ratios and VT1. For VT2, there were negative relationships with 40 digit ratios (right 2D:4D, r = -.43, p < .0001; left 2D:4D, r = -.21 and Dr-l, r = -.21, both p = -.2141 .02). Digit ratios are negatively related to VO_{2max} with large (right 2D:4D) and medium (left 42 2D:4D, Dr-l) effect sizes. For VT2, there were also negative correlations, which were 43 medium (right 2D:4D) and small (left 2D:4D, Dr-l). Conclusion: Our findings may help 44 clarify the relationships between digit ratios and high-intensity actions for extended periods, 45 46 which are dependent on efficient oxygen metabolism.

Key Words: Prenatal testosterone, Aerobic fitness, Digit ratios, Soccer

48

47

49

50

51

52

53

Introduction

55

The relative lengths of the 2nd and 4th digits (2D:4D) and the side difference in 2D:4D (Dr-1: 56 right-left 2D:4D) are thought to be negative correlates of 1st-trimester testosterone and 57 positive correlates of 1st-trimester oestrogen (Manning et al., 1998; Manning et al., 2002; 58 Breedlove, 2010; Swift-Gallant et al., 2020). The 2D:4D and Dr-l show sexual dimorphism 59 (males<females); the sex difference appears in the 1st trimester and shows little change in 60 children, juveniles and adults (Malas et al., 2006; Trivers et al., 2006; Manning et al., 2022). 61 62 In contrast to the links between digit ratios and prenatal sex steroids, there is little evidence of associations between 2D:4D and background levels of testosterone or oestrogen in adults 63 64 (Hönekopp et al., 2007). Manning and Taylor (2001) were the first to report that 2D:4D was negatively associated with 65 performance among male participants from a range of sports, including >300 elite footballers 66 competing in the English Leagues. In addition, meta-analyses have found negative 67 relationships between 2D:4D and performance in a number of sports with mean right-hand 68 effect sizes of r = -0.28 (Hönekopp and Schuster, 2010) and weak negative relationships with 69 hand grip strength (Pasanen et al., 2022). With regard to endurance disciplines, Manning et 70 al., (2007) have reported strong correlations between 2D:4D and running speed in middle-71 and long-distance races(r^2 values of approximately 25% for males and females). They 72 suggested that 2D:4D may be a strong correlate of vascular health. However, reports of 73 74 associations between 2D:4D, VO_{2max} and ventilatory threshold (VT) employing objective labbased measures of VO_{2max} and VT have yielded mixed results from samples that were small 75 and were recruited from a range of backgrounds in sports(Hill et al., 2012; Holzapfel et al., 76 2016; Lombardo et al., 2020). In this regard, it is important to examine the relationships 77 between digit ratios (2D:4D and Dr-l) and oxygen metabolism in a larger sample of athletes 78 who participate in the same sport. The latter includes VO_{2max}(maximal oxygen consumption; 79

Hill and Lupton, 1923) and ventilator thresholds [(VT1 the point during exercise at which 80 pulmonary ventilation and carbon dioxide output begin to increase exponentially; Cerezuela-81 Espejo et al., 2018), and VT2 or RC (the point associated with hyperventilation at which 82 lactate is rapidly increasing with intensity; Meyer et al., 2004). 83 Evidence for links between digit ratios and oxygen metabolism may be indicated by the types 84 of sport linked to 2D:4D or Dr-l. Low values of digit ratios have been reported to be 85 associated with high performance in a range of sports. For males, these include football 86 (soccer; Manning and Taylor, 2001), rugby (Bennett et al., 2010), skiing (Manning, 2002), 87 rowing(Longman et al., 2011), surfing (Kilduff et al., 2011), wrestling (Keshavarz et al., 88 2017), basketball (Klapprodt et al., 2018) and for females, rowing (Hull et al., 2015), skiing 89 (Manning 2002), and Olympic athletes participating in power, endurance and technical sports 90 91 (Eklund et al., 2020). Therefore, low digit ratios may be linked to both strength and endurance. However, a consideration of associations between 2D:4D and running speed 92 suggests that the latter shows greater effect sizes than the former. In this regard, Manning et 93 al., (2007) and Longman et al., (2015) have reported correlations between 2D:4D and running 94 speed in long-distance races ranging in strength from r = .40 to r = .60 in males and r = .20 to 95 r = .30 in females. In contrast, 2D:4D was indicated to be weakly related to sprinting speed, 96 with correlations averaging about r = .10 (Hönekopp and Schuster, 2010; Manning and Hill, 97 98 2009). Physiological variables (VO_{2max}, velocity at maximal oxygen uptake, and changes in lactate levels), training load and fat mass are considered the main factors determining 99 performance in long-distance races (Alvero-Cruz et al, 2020). The strong relationship 100 between 2D:4D and speed in long-distance races suggests that 2D:4D may be a negative 101 correlate of maximal aerobic performance, and in particular, it is likely to be predictive of 102 103 maximal oxygen uptake (VO_{2max}) and/or Ventilatory Thresholds (VT1 and VT2).

However, attempts to quantify relationships between digit ratio and VO_{2max} and VT1 and VT2 have met with mixed results. Hill et al., (2012) considered relationships between digit ratios and oxygen metabolism in 41 boys (mean age 13.9 [SD1.3] years). They found no significant relationships for right or left 2D:4D, but there were negative correlations of medium strength for Dr-l and VO_{2max}. In contrast, Holzapfel et al., (2016) reported no significant correlations between 2D:4D (Dr-l was not considered) and VO_{2max} in a sample of 26 men and 28 women, but strong negative relationships were demonstrated for 2D:4D and VT. Furthermore, Lombardo and Otieno (2020) reported on digit ratio and aerobic fitness variables in 11 boys and 15 girls, aged between 11 and 19 years, who were the top five finishers in 10 or more races of 10 km. In their study, boys (but not girls) with lower right 2D:4D had significantly greater VO_{2max}. Girls (but not boys) with lower right 2D:4D had significantly greater VT. Thus, it appears that digit ratios are related to maximal aerobic performance, but the strength of the relationship and the relative importance of VO_{2max} and VT need to be clarified. In general, sample sizes thus far were small, and participants varied in their participation in sports. Therefore, we consider relationships between digit ratios (2D:4D and Dr-l) and VO_{2max}, and VT1 and VT2 in a large sample of male professional football players.

Materials and Methods

122 Participants

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

123

124

125

126

127

128

An initial sample of 143 professional male football players (age: 25.21 ± 5.47 years, height: 180.15 ± 6.12 cm, weight: 76.40 ± 7.12 kg) participating in Division 1 and 2 in the Eastern Mediterranean was recruited. The sample included 133 Caucasian and 10 Black participants. Due to significant differences in the anthropometric characteristics and digit ratios between the Caucasian and Black players, our statistical analyses were mainly focused on the Caucasian players (n=133).

Testing was undertaken during the months of June and July before the pre-season period. Exclusion criteria included injuries within the last two months before the testing. Anthropometric measurements (age, stature, body weight, body fat and hand scans) were recorded before the physical tests. Players' characteristics are given in Table 1. The players completed an incremental cardiopulmonary test to exhaustion on a treadmill. All players were familiar with the testing protocol as this was part of their annual testing. They were instructed to avoid heavy physical activity the day prior to the testing. All participants completed an informed consent after being briefed about the procedures, and the technical director of the team approved all the testing protocols. The research complied with the relevant national regulations, was conducted in accordance with the Declaration of Helsinki, and was approved by the National Committee of Bioethics (EEBK EP 2022.01.290).

Procedure

- 141 Anthropometric measurements
- Anthropometric measurements were conducted using a wall stadiometer (Leicester; Tanita,

 Tokyo, Japan) to determine the players' stature and a leg-to-leg bioelectrical impedance

 analyser (BC418MA; Tanita) to assess body composition (% body fat). The players were

 instructed to follow the standard guidelines prior to the bioelectrical impedance testing (Kyle

Hand scans

et al., 2004).

Players were asked to place their hand on the surface of the photocopier (EPSON scanner, DS-50000) with the palm facing downwards and fingers as straight as possible according to the methodology described by previous investigators (Manning, 2002). They were instructed not to exert too much pressure but lightly place their fingers on the photocopier and wait until the scan was completed. The scan was evaluated by a single examiner, and in cases where it

was not clear, it was repeated. The finger length was measured twice by the same investigator, blind to the oxygen data, and the 2D:4D ratio was calculated from each set of scans. Digit length was measured to an accuracy of 0.05 mm using Vernier callipers (Mitutoyo, D15, Japan).

Incremental cardiopulmonary testing on a treadmill

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

The players completed an incremental cardiopulmonary test to exhaustion (CPET) on a treadmill (HP Cosmos Quasar med, HP Cosmos Sports, and Medical GmbH, Nussdorf-Traunstein, Germany). Gas exchange measurements were collected with reusable masks, a turbine flow meter, and a two-way nonrebreathing valve (model 7940, Hans Rudolph, Kansas City, MO). Heart rate (COSMED wireless HR monitor, Rome, Italy), VO2, carbon dioxide (VCO2) production and expired minute volume (VE) were continuously monitored throughout the test, and a breath-by-breath analysis was performed on a computerised (Cosmed Quark CPET, Rome, Italy) system. Before each test, the air VO2 flowmeter and oxygen-carbon dioxide meters were calibrated with a three-litre air syringe and a gas of known oxygen (16.5%) based on the manufacturer's recommendations. Throughout the testing, laboratory conditions were kept constant, with the temperature being around 21-22 degrees (C) and the relative humidity around 50%. During the test, the inclination was kept constant at 1%. The players started the test at a speed of 8km/hr, and the speed was increased every 3.15 minutes by 2km/hr until they reached volitional exhaustion or could no longer continue. The recovery speed was 5 km/h for 2-3 minutes. The VO_{2max} was identified after filtering the results by indicating the highest value

for an average of 10 seconds and was expressed relative to body mass (ml/kg/min). The ventilatory threshold (VT1) was identified through the V-slope method (the point at which the increase in the rate of elimination of carbon dioxide is greater than the increase in VO2)

and was verified at the nadir of the VE/VO2 curve. The respiratory compensation point (VT2 or RC) was determined at the nadir of the VE/VCO2 curve (Beaver et al., 1986).

Statistical Analysis

Means and standard deviations (mean \pm SD) were calculated for all the parameters. The homogeneity of variance was tested using the Brown-Forsythe test, and the normality assumption was verified using the Shapiro–Wilk test. Interclass correlation coefficients (ICC) (absolute agreement) between the first and second 2D:4D's of the right and left digits were calculated. Pearson-product moment correlation coefficients were used to determine the association between 2D:4D, VO_{2max} and its associated ventilatory thresholds. Correlations were referred to as trivial (0–0.1), small (0.1–0.3), moderate (0.3–0.5), large (0.5–0.7), very large (0.7–0.9), nearly perfect (>0.9) and perfect (1.0) (Hopkins et al., 2009). Three multiple regression analyses with independent variables age, right 2D:4D and left 2D:4D and dependent variables VO_{2max} or VT1 or VT2 were performed. All statistical analyses were performed in IBM® SPSS® Statistics, version 26.0, for Windows (SPSS Inc., Chicago, IL, USA), and the statistical significance was set at p <0.05.

Results

Two values of digit ratios were calculated. Intra-class correlations coefficients (r_I , used for the assessment of the consistency of the measurements) were high and significant for right 2D:4D (n = 142, $r_I = .976$, F = 82.79, p < .0001), left 2D:4D (n = 140, $r_I = .960$, F = 48.48, p < .0001) and Dr-1 (n = 139, $r_I = .954$, F = 42.29, p < .0001). The average of the two measurements was used to obtain the final values for right and left 2D:4D and Dr-1 ratios.

Descriptive statistics for the total sample and the sample split by ethnicity are given in Table

1. In comparison to Caucasians, Black players had greater mass, BMI, % body fat, and VO2

at VT as well as lower right and left 2D:4D. Therefore, we removed the Black players from the sample and reported relationships for Caucasians (n = 133) only for the following

analyses.

201

202

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

There were no significant relationships between digit ratios (right and left and Dr-l) and age or body size variables (r=0.04 between age and right 2D:4D, r= 0.06 between height and right 2D:4D, r=0.03 between weight and right 2D:4D, r=0.04 between age and left 2D:4D, r=0.06 between height and left 2D:4D, r=0.07 between weight and left 2D:4D, all p>.05). The correlations between digit ratios (right and left and Dr-1), and VO_{2max} and ventilatory thresholds VT1 and VT2 are given in Table 2. Correlations were strongest between digit ratios and VO_{2max}, effect sizes were greatest for right 2D:4D, and all correlations were negative. With regard to VO_{2max} , there was a large correlation with right 2D:4D (r = -0.65; Figure 1) and medium correlations with left 2D:4D (r = -0.37) and Dr-1 (r = -0.30). There were no significant relationships between digit ratios and VT1 (r varying from -0.02 to -0.12). Considering VT2, right 2D:4D showed a moderate correlation (r = -0.43), and there were small correlation coefficients for left 2D:4D and Dr-1 (both r = -0.21). VO_{2max}, V1 and V2 were interrelated with varying strengths (very large, VO_{2max} and VT2, r = .73; large, VT1and VT2, r = .59; moderate, VO_{2max} and VT1, r = .34: all p < .0001). In addition to the correlations (r) for the Caucasian participants, we also considered the total sample (i.e. Caucasian and Black players, n = 143) together with the total sample after

sample (i.e. Caucasian and Black players, n=143) together with the total sample after ethnicity effects were removed (standardised regression coefficient, b)—the values of b are presented in parenthesis in Table 2. There was one notable change in r and p values, i.e. for the total sample of Caucasian plus Black players; right 2D:4D was now negatively and significantly related to VT1 (r=-.21, p=.01). There were no substantial differences in effect

sizes and p values between the Caucasian sample and the total sample when ethnicity effects

- were removed.
- Age may influence VO_{2max}, VT1 and VT2. Therefore, we performed three multiple regression
- analyses with independent variables age, right 2D:4D and left 2D:4D and dependent variables
- VO_{2max} or VT1 or VT2. With regard to VO_{2max}, the overall relationship was r = 0.67 ($r^2 =$
- 229 0.45, age b = -.08, SE = .06, p = .25, right 2D:4D b = -0.61, SE = 9.77, t = -8.35, p < .0001,
- 230 left 2D:4D b = -0.13, SE = 10.81, t = -1.80, p = 0.08). For VT1 the overall relationship was r
- = 0.21 (r^2 = 0.04). There was a small negative relationship for age but no relationships for
- digit ratios (age b = -0.17, SE = .06, t = -1.98, p = 0.049). Considering VT2, the overall
- relationship was r = 0.48 ($r^2 = 0.23$). There was a small negative relationship with age and a
- 234 moderate negative association for right 2D:4D (age b = -0.22, SE = .06, t = -2.82, p = 0.006,
- right 2D:4D b = -0.41, SE = 10.40, t = -4.84, p < .0001). There was no relationship for left
- 236 2D:4D.

237

Discussion

- Football is an intermittent sport with repeated high-intensity phases. As a result of
- improvements in training techniques, football players today are much more similar to
- endurance athletes than 50 years ago (Edwards et a., 2003). Therefore, comparisons between
- our results and those from endurance athletes are appropriate.
- Our finding of a mean VO_{2max} of 56.05 ± 4.53 was close to large sample measures of elite
- football players (range, goalkeeper 50.42±4.2 to winger-sides back 60.53±5.02, median
- 58.25: Manari et al., 2016). In our total sample of 143 participants, there were 133
- Caucasians and 10 Black football players. The latter differed from the former in their 2D:4D
- (Caucasian>Black) and in mass, BMI, % body fat, and VO2 at VT1. High 2D:4D in
- Caucasians and low 2D:4D in Black populations have been reported in a number of studies
- 248 (Manning, 2002; Butovskaya et al., 2021). Such differences can obscure relationships.

Therefore, the less numerous group was removed, and subsequent analyses focused on

250 Caucasians.

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

With regard to our Caucasian sample, we have found significant negative relationships between all three-digit ratio variables (right 2D:4D, left 2D:4D and Dr-l) and VO_{2max}. The large correlation between right 2D:4D and VO_{2max} was the strongest of the three associations, such that right 2D:4D explained 42% of the variance in VO_{2max}. Associations for left 2D:4D and Dr-l with VO_{2max} were medium in strength. There were no significant relationships between digit ratios and VT1. For VT2, all digit ratio correlations were negative and significant, with a moderate (and strongest) relationship for right 2D:4D and small correlations for left 2D:4D and Dr-l. Our study is one of the larger studies to consider relationships between digit ratios and VO_{2max} and VTs in males. The sample was relatively homogeneous in that the participants were all male Caucasian professional football players competing in Leagues 1 and 2, Eastern Mediterranean. Moreover, they can be regarded as being relatively homogenous in terms of their exercise regime. A similar study by Hill et al., (2012) indicated no association between 2D:4D (right or left) and VO_{2max} but reported a significant negative correlation for Dr-l in young athletic teenage boys of Middle East origin (age: 13.9 ± 1.3 years) during an incremental treadmill test. We have replicated this latter association in our larger adult male sample. Hill et al., (2012) participants were drawn from a wide range of sports with different training regimes (soccer, squash, table tennis and athletics). This may have masked the relationship between right and left 2D:4D and VO_{2max}. Importantly, both our present sample and that of Hill et al., (2012) controlled for ethnicity by considering a single ethnic group. Holzapfel et al., (2016)reported little or no relationship between 2D:4D (Dr-l was not

considered) and VO_{2max} in a sample of 26 men (13 sedentary and 13 distance runners).

However, they found large negative correlations between 2D:4D and VT. On the contrary, in our sample, there were no relationships between digit ratios and VT1. The distance runners in the Holzapfel et al., (2016)study had higher mean VO_{2max} (62.6±11.2) than our sample of football players (55.91 \pm 4.51, Cohen's d=.78). However, this was unlikely to account for the differences as there were large correlations between digit ratios and VT in both their sedentary and runner samples. Their sample was recruited from the student population of a South-Eastern US University, and the authors did not report any controls for ethnicity. Thus, the discrepancies between the Holzapfel et al., (2016) study and the Hill et al., (2012) and the present study may have arisen as the result of differences in sample size and controls for ethnicity. In this regard, the removal of ethnicity controls in our present study resulted in a significant relationship between right 2D:4D and VT1. A similar study by Lombardo and Otieno (2020)reported correlations between right 2D:4D and VO_{2peak}, VT and Point of Equivalent Change (PEC) in 11 boys who were elite distance runners. All three variables were negatively related to right 2D:4D with two ($VO_{2peak}r = -.62$; PEC r = -.66) showing significance at p<.05. However, significance was lost for both when adjusted for mass. The strength of the correlation with right 2D:4D was similar to that of our finding for right 2D:4D and VO_{2max.} These findings suffer from small sample sizes. However, we judge them to be not incompatible with our findings. With regard to the value of 2D:4D to coaches and scouts. We suggest that 2D:4D may be of predictive value in sports that are performance-dependent on high values of VO_{2max} (e.g. distance running, tennis, rowing & football). Values of 2D:4D appear to be more or less stable across puberty, thus, 2D:4D may yield predictive information in adolescents. An explanation for the links between low 2D:4D and high values of Vo_{2max} and VT2 may lie in the relationship between 2D:4D and prenatal testosterone. Manning (2002) has suggested

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

that 2D:4D is a highly conserved trait that is linked to the early emergence of tetrapods from an aquatic to a terrestrial existence. Since this suggestion, there have been some 30 reports concerning 2D:4D sexual dimorphism in amphibians, reptiles, birds, artiodactyls, rodents, and primates (Lupu et al., 2023). The findings suggest a pattern of sex differences of moderate effect, with some species showing male 2D:4D lower than female 2D:4D and other species having the reverse pattern. However, not all species have significant sexual dimorphism in 2D:4D. For example, Lombardo & Thorpe (2008) did not report evidence of sexual dimorphism in 2D:4D green anolis lizards (Anolis carolinensis), and Lombardo et al., (2008) did not find evidence of sexual dimorphism in 2D:4D in four species of birds (house sparrows, tree swallows, budgerigars, chickens). When the four species were pooled, male 2D:4D was greater than female 2D:4D. With regard to more direct evidence that sex differences in 2D:4D are testosterone dependent, there are nine studies that included manipulations of testosterone or its receptor; six of these report a masculinisation effect, one a feminisation effect, and two a null effect (Manning and Fink, 2023). The emergence from an aquatic existence is associated with a suite of traits, including the ability to process gaseous oxygen (Manning, 2002; Manning and Fink, 2023). Low testosterone compromises mitochondrial function (Yan et al., 2017), and in human males, it is linked to cardiovascular disease (Harada, 2018). High 2D:4D is associated with elevated fibrinogen levels and early myocardial infarction (Manning et al., 2019). Thus, our expectation is that low 2D:4D is related to efficient oxygen metabolism. Our study has a number of limitations. We have not considered non-Caucasian and female football players as it was not possible to recruit sufficient numbers. Moreover, we suggest that associations between 2D:4D and oxygen metabolism should be considered in a variety of sports. These could range from those that require a very high level of aerobic fitness (e.g.

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

321	professional cyclists participating in the Girod d'Italia, Tour de France and Vuelta de Espana)
322	to those in which fitness is somewhat less important (e.g. table tennis).
323	Conclusions
324	In conclusion, we have found significant negative correlations between digit ratios and
325	VO _{2max} in 133 professional male football players. They were large (right 2D:4D) and medium
326	(left 2D:4D, Dr-l) in effect size. For VT2, there were also significant negative correlations,
327	which were medium (right 2D:4D) and small (left 2D:4D, Dr-l) in effect size. There were no
328	associations between digit ratios and VT1. All associations were controlled for ethnicity. We
329	hope these findings help to clarify associations between digit ratios and oxygen metabolism
330	in men. Further work is necessary to quantify these associations in women.
331	Figure legends
332	Figure 1 . VO2max and right 2D:4D ($r =65$, $R^2 = 0.425$)

- 334 References
- 335 Alvero-Cruz JR, Carnero EA, García MAG, Alacid F, Correas-Gómez L, Rosemann T,
- Nikolaidis PT, Knechtle B. (2020). Predictive Performance Models in Long-Distance
- Runners: A Narrative Review. *Int J Environ Res Public Health*, 17(21), 8289.
- Beaver, W. L., Wasserman, K., & Whipp, B. J. (1986). A new method for detecting anaerobic
- threshold by gas exchange. *Journal of applied physiology*, 60(6), 2020-2027.
- Bennett, M., Manning, J. T., Cook, C. J., & Kilduff, L. P. (2010). Digit ratio (2D: 4D) and
- performance in elite rugby players. *Journal of sports sciences*, 28(13), 1415-1421.
- 342 Breedlove, S. M. (2010). Minireview: organisational hypothesis: instances of the
- 343 fingerpost. *Endocrinology*, *151*(9), 4116-4122.
- Butovskaya, M., Burkova, V., Apalkova, Y., Dronova, D., Rostovtseva, V., Karelin, D., ...
- &Batsevich, V. (2021). Sex, population origin, age and average digit length as predictors of
- digit ratio in three large world populations. *Scientific reports*, 11(1), 8157.
- 347 Cerezuela-Espejo, V., Courel-Ibáñez, J., Morán-Navarro, R., Martínez-Cava, A., & Pallarés,
- 348 J. G. (2018). The relationship between lactate and ventilatory thresholds in runners: validity
- and reliability of exercise test performance parameters. Frontiers in physiology, 9, 1320.
- Edwards, A. M., Clark, N., & Macfadyen, A. M. (2003). Lactate and ventilatory thresholds
- reflect the training status of professional soccer players where maximum aerobic power is
- unchanged. *Journal of sports science & medicine*, 2(1), 23.
- Eklund, E., Ekström, L., Thörngren, J. O., Ericsson, M., Berglund, B., & Hirschberg, A. L.
- 354 (2020). Digit ratio (2D: 4D) and physical performance in female Olympic athletes. Frontiers
- *in Endocrinology*, *11*, 292.
- Harada, N. (2018) Role of androgens in energy metabolism affecting on body composition,
- metabolic syndrome, type 2 diabetes, cardiovascular disease, and longevity: lessons from a
- meta-analysis and rodent studies, *Bioscience, Biotechnology, and Biochemistry*, 82:10, 1667.
- Hill, A. V., & Lupton, H. (1923). Muscular exercise, lactic acid, and the supply and
- utilisation of oxygen. *QJM: Quarterly Journal of Medicine*, (62), 135-171.
- 361 Hill, R., Simpson, B., Millet, G., Manning, J., & Kilduff, L. (2012). Right–left digit ratio
- 362 (2D: 4D) and maximal oxygen uptake. *Journal of sports sciences*, 30(2), 129-134.
- Holzapfel, S. D., Chomentowski III, P. J., Summers, L. A. M., & Sabin, M. J. (2016).
- Running head: 2D: 4D and Aerobic Fitness in Young Adults: The relationship between digit
- ratio (2D:4D), VO2max, ventilatory threshold and running performance. *International*
- *Journal of Sports Sciences & Fitness*, *6*(1).

- Hönekopp, J., & Schuster, M. (2010). A meta-analysis on 2D: 4D and athletic prowess:
- 368 Substantial relationships but neither hand out-predicts the other. *Personality and Individual*
- 369 *Differences*, 48(1), 4-10.
- Hönekopp, J., Bartholdt, L., Beier, L., & Liebert, A. (2007). Second to fourth digit length
- ratio (2D: 4D) and adult sex hormone levels: new data and a meta-analytic
- 372 review. Psychoneuroendocrinology, 32(4), 313-321.
- Hopkins, W., Marshall, S., Batterham, A., & Hanin, J. (2009). Progressive statistics for
- 374 studies in sports medicine and exercise science. *Medicine+ Science in Sports+*
- 375 Exercise, 41(1), 3.
- Hull, M. J., Schranz, N. K., Manning, J. T., & Tomkinson, G. R. (2015). Relationships
- between digit ratio (2D: 4D) and female competitive rowing performance. *American Journal*
- *of Human Biology*, *27*(2), 157-163.
- Keshavarz, M., Bayati, M., Farzad, B., Dakhili, A., & Agha-Alinejad, H. (2017). The second
- to fourth digit ratio in elite and non-elite Greco-Roman wrestlers. *Journal of human*
- 381 *kinetics*, 60, 145.
- Kilduff, L. P., Cook, C. J., & Manning, J. T. (2011). Digit ratio (2D: 4D) and performance in
- male surfers. *The Journal of Strength & Conditioning Research*, 25(11), 3175-3180.
- Klapprodt, K. L., Fitzgerald, J. S., Short, S. E., Manning, J. T., & Tomkinson, G. R. (2018).
- Relationships between the digit ratio (2D: 4D) and game-related statistics in professional and
- semi-professional male basketball players. American Journal of Human Biology, 30(6),
- 387 e23182.
- Kyle, U. G., Bosaeus, I., De Lorenzo, A., Deurenberg, P., Elia, M., Gómez, J. M., et al.
- 389 (2004). Composition of the ESPEN Working Group. Bioelectrical impedance analysis--part I:
- review of principles and methods. *Clinical Nutrition*, 23(5):1226–43.
- Lombardo, M. P., & Otieno, S. (2021). The associations between digit ratio, aerobic fitness,
- 392 physical skills, and overall physical fitness of elite youth distance runners. American Journal
- *of Human Biology*, *33*(1), e23448.
- 394 Longman, D., Stock, J. T., & Wells, J. C. K. (2011). Digit ratio (2D: 4D) and rowing
- ergometer performance in males and females. American journal of physical
- 396 anthropology, 144(3), 337-341.
- Lombardo, M. P., & Thorpe, P. A. (2008). Digit ratios in green anolis lizards (Anolis
- 398 *carolinensis*). *Anat Rec (Hoboken), 291*(4), 433-440.

- Lombardo, M. P., Thorpe, P. A., Brown, B. M., & Sian, K. (2008). Digit ratio in birds. Anat
- 401 Rec (Hoboken), 291(12), 1611-1618.
- 402 Longman, D., Wells, J. C., & Stock, J. T. (2015). Can persistence hunting signal male
- 403 quality? A test considering digit ratio in endurance athletes. *PLoS One*, 10(4), e0121560.
- Lupu DC, Monedero I, Rodriguez-Ruiz C, Pita M, Turiegano E (2023). In support of 2D:4D:
- 405 More data exploring its conflicting results on handedness, sexual orientation and sex
- 406 differences. PLoS ONE 18(8): e0280514.
- Malas, M. A., Dogan, S., Evcil, E. H., &Desdicioglu, K. (2006). Fetal development of the
- 408 hand, digits and digit ratio (2D: 4D). Early human development, 82(7), 469-475.
- 409 Manari, D., Manara, M., Zurini, A., Tortorella, G., Vaccarezza, M., Prandelli, N., ... & Galli,
- D. (2016). VO 2Max and VO 2AT: athletic performance and field role of elite soccer
- players. Sport Sciences for Health, 12, 221-226.
- Manning, J. T. (2002). The ratio of 2nd to 4th digit length and performance in skiing. *Journal*
- of sports medicine and physical fitness, 42(4), 446.
- 414 Manning, J. T. (2002). Digit ratio: A pointer to fertility, behavior, and health. Rutgers
- 415 University Press.
- 416 Manning, J.T., Bundred, P.E., Kasielska-Trojan, A., Smith-Straney, T. & Mason, L. (2019).
- Digit ratio (2D:4D), myocardial infarction and fibrinogen in men. Early Human
- 418 *Development*, 133, 18-22.
- Manning, J.T. & Fink, B. (2023). Digit ratio (2D:4D) and its relationship to foetal and
- 420 maternal sex steroids: A mini-review. Early Human Development, 183, 105799.
- 421 Manning, J. T., & Hill, M. R. (2009). Digit ratio (2D: 4D) and sprinting speed in
- boys. *American Journal of Human Biology: The Official Journal of the Human Biology*
- 423 Association, 21(2), 210-213.
- Manning, J. T., & Taylor, R. P. (2001). Second to fourth digit ratio and male ability in sport:
- implications for sexual selection in humans. *Evolution and human behavior*, 22(1), 61-69.
- 426 Manning, J. T., Fink, B., Mason, L., Kasielska-Trojan, A., & Trivers, R. (2023). The effects
- of sex, nation, ethnicity, age and self-reported pubertal development on participant-measured
- right-left 2D: 4D (Dr-l) in the BBC internet study. Journal of biosocial science, 55(2), 383-
- 429 395.
- 430 Manning, J. T., Morris, L., & Caswell, N. (2007). Endurance running and digit ratio (2D:
- 431 4D): implications for fetal testosterone effects on running speed and vascular
- health. *American Journal of Human Biology: 19*(3), 416-421.

- Manning, J. T., Scutt, D., Wilson, J., & Lewis-Jones, D. I. (1998). The ratio of 2nd to 4th
- digit length: a predictor of sperm numbers and concentrations of testosterone, luteinising
- hormone and oestrogen. *Human Reproduction (Oxford, England)*, 13(11), 3000-3004.
- 436 Meyer, T., Faude, O., Scharhag, J., Urhausen, A., & Kindermann, W. (2004). Is lactic
- acidosis a cause of exercise induced hyperventilation at the respiratory compensation
- point?. British journal of sports medicine, 38(5), 622-625.
- Pasanen, B. E., Tomkinson, J. M., Dufner, T. J., Park, C. W., Fitzgerald, J. S., & Tomkinson,
- 440 G. R. (2022). The relationship between digit ratio (2D: 4D) and muscular fitness: A
- systematic review and meta-analysis. *American journal of human biology*, 34(3), e23657.
- Swift-Gallant, A., Johnson, B. A., Di Rita, V., & Breedlove, S. M. (2020). Through a glass,
- darkly: Human digit ratios reflect prenatal androgens, imperfectly. *Hormones and*
- 444 Behavior, 120, 104686.

451

- Trivers, R., Manning, J., & Jacobson, A. (2006). A longitudinal study of digit ratio (2D: 4D)
- and other finger ratios in Jamaican children. Hormones and behavior, 49(2), 150-156.
- Testosterone Upregulates the Expression of Mitochondrial ND1 and ND4 and Alleviates the
- 448 Yan, W, Kang, Y, Ji, X, Li, S, Li, Y, Zhang, G, Cui, H, & Shi, G. (2017). Oxidative Damage to
- the Nigrostriatal Dopaminergic System in Orchiectomized Rat. Oxidative Medicine and
- 450 *Cellular Longevity*, doi.org/10.1155/2017/1202459