

Central Lancashire Online Knowledge (CLOK)

Title	Increasing cadence with a metronome and running barefoot changes the sagittal kinematics of the lower limbs and trunk
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/50890/
DOI	https://doi.org/10.1080/02640414.2024.2330818
Date	2024
Citation	Molina-Molina, Alejandro, Delgado-García, Gabriel, Richards, James, Mercado-Palomino, Elia, Manuel Soto-Hermoso, Víctor and Ángel Latorre-Román, Pedro (2024) Increasing cadence with a metronome and running barefoot changes the sagittal kinematics of the lower limbs and trunk. <i>Journal of Sports Sciences</i> , 42 (5). pp. 404-414. ISSN 0264-0414
Creators	Molina-Molina, Alejandro, Delgado-García, Gabriel, Richards, James, Mercado-Palomino, Elia, Manuel Soto-Hermoso, Víctor and Ángel Latorre-Román, Pedro

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1080/02640414.2024.2330818>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

**Increasing cadence with a metronome and running barefoot changes the sagittal
kinematics of the lower limbs and trunk.**

Alejandro MOLINA-MOLINA ^{1-2 *}, Gabriel DELGADO-GARCÍA ³, Jim RICHARDS ⁴, Elia
MERCADO-PALOMINO ², Víctor Manuel SOTO-HERMOSO ² and Pedro Ángel LATORRE-
ROMÁN ⁵

¹ Universidad San Jorge. Campus Universitario, Autov A23 km 299, 50830. Villanueva de
Gállego, Zaragoza, Spain.

² Sport and Health University Research Institute (iMUDS), Department of Physical Education
and Sports, Faculty of Sport Sciences, University of Granada, 18071, Granada, Spain.

³ SER Research Group, CESAG, Pontifical University of Comillas. 07013 Palma, Spain

⁴ Allied Health Research Unit, University of Central Lancashire, Victoria St., Preston, PR1 2HE,
UK.

⁵ Universidad de Jaén, Department of Didactics of Corporal Expression. Campus de Las
Lagunillas s/n. D2 Building, Dep. 142.23071, Jaén, Spain.

*Correspondence: author: Alejandro Molina-Molina, Universidad San Jorge. Campus
Universitario, Autov A23 km 299, 50830. Villanueva de Gállego Zaragoza. Spain.

E-mail: amolinam@usj.es

Phone: 0034 625 538 520

Twitter: @amolina_m

WORD COUNT: 4204 words

**Increasing cadence with a metronome and running barefoot changes the sagittal
kinematics of the lower limbs and trunk.**

ABSTRACT

The purpose was to compare two non-laboratory based running retraining programs on lower limb and trunk kinematics in recreational runners. Seventy recreational runners (30 ± 7.3 years old, 40% female) were randomised to a barefoot running group (BAR), a group wearing a digital metronome with their basal cadence increased by 10% (CAD), and a control group (CON). BAR and CAD groups included intervals from 15 to 40 minutes over 10 weeks and 3 days/week. 3D sagittal kinematics of the ankle, knee, hip, pelvis, and trunk were measured before and after the retraining program, at comfortable and high speeds. A 3×2 mixed ANOVA revealed that BAR and CAD groups increased knee and hip flexion at footstrike, increased peak hip flexion during stance and flight phase, decreased peak hip extension during flight phase, and increased anterior pelvic tilt at both speeds after retraining. In addition, BAR increased ankle plantar flexion at footstrike and increased anterior trunk tilt. Both retraining programs demonstrated significant moderate to large effect size changes in parameters that could reduce the mechanical risks of injury associated with excessive knee stress, which is of interest to coaches, runners and those prescribing rehabilitation and injury prevention programs.

KEYWORDS: mobile app; long-term usage; step rate; unshod; running form.

Increasing cadence with a metronome and running barefoot changes the sagittal kinematics of the lower limbs and trunk.

INTRODUCTION

Endurance running is an activity that has become widespread among the recreational population with the aim of improving health and personal performance (1,2). However, the annual incidence of running-related injuries can rise to 74% among long-distance runners (3–5). Excessive knee loading, and high vertical and horizontal impact forces have been associated with running-related injuries such as knee injuries (6,7), which are common among recreational long-distance runners (8). Running retraining is a process of learning a new running motor pattern, which has been associated with improved performance and injury prevention and is often included in rehabilitation programmes (9). A fundamental purpose of running retraining is to modify a motor pattern associated with a potential risk of running-related injury due to overuse, which is often considered to be a faulty motor pattern.

Increasing cadence has been proposed as a method for running retraining that could be integrated into a non-laboratory setting through the use of a mobile metronome application (10,11), and has been suggested to reduce impact forces and knee loading (12–14). Furthermore, this reduction in impact forces and knee loading has been associated with changes in lower limb kinematics, such as an increase in knee flexion at footstrike or a decrease in knee flexion during stance phase and ankle dorsiflexion at footstrike (10–12,14). For example, a two-week retraining program that increased cadence by 10% was shown to reduce foot and ankle angles at footstrike, impact forces and anterior trunk tilt, but no changes were found in knee and hip flexion (14). However, reduced hip flexion at footstrike was found to be an immediate effect of a 10% increase in cadence under laboratory conditions (15). After 12 weeks of field-based retraining using 10% increased cadence feedback, participants reduced impact forces, foot angle at footstrike, and peak knee flexion during stance phase (11). A reduction in peak knee flexion has been shown to be a useful predictor of a reduction in knee loading (12).

Barefoot running has also been considered as a non-laboratory retraining alternative that promotes impact attenuation, which influences changes in lower limb kinematics (16–21) and has been hypothesised to reduce the risk of injury (16,22,23). Some of the kinematic indicators associated with a reduced risk of injury were reduced ankle dorsiflexion at footstrike, increased knee flexion at footstrike and reduced peak knee flexion at stance (16–21). An 8-week barefoot running program found a subgroup of responders who reduced impact forces and ankle dorsiflexion (18). However, two other fully immersive 8-week barefoot running programs showed no significant changes in ankle dorsiflexion, cadence or stride length (24,25), nor in foot or knee kinematics (24). In contrast, a recent more conservative 10-week programme of barefoot running intervals and habitual footwear training showed a reduction in foot angle at initial contact and some changes in spatiotemporal parameters when evaluating runners running in their natural footwear condition, but lower limb kinematics were not assessed (26).

While several studies of barefoot running and studies of increasing basal cadence by 10% separately provide strong evidence of reduced impact (12–14,17,18) and changes in foot kinematics and spatiotemporal parameters, such as a reduction of ankle dorsiflexion and foot at footstrike or increase in cadence (10,11,23,25,26,12,14,16–21), the effects of long-term and out-of-laboratory retraining programs on lower limb kinematics remain inconsistent. Although the effects of barefoot running and a 10% increase in cadence on running kinematics are similar, these effects have not been compared in a homogeneous sample. Except for a recent study that evaluated the effect of increasing cadence versus barefoot running on spatiotemporal parameters, footstrike angle and prevalence of rearfoot strike in a homogeneous population over 10 weeks (26). Furthermore, as mentioned above, there is a lack of evidence on the effects of these programs on the kinematics of proximal body regions (i.e., pelvis and trunk). A more flexed trunk during running demands more hip extensor work and less extensor work during running reducing patellofemoral pain (27,28).

Therefore, the aim of this study was to compare the effects of two non-laboratory based 10-week running retraining programs on lower limb and trunk running kinematics in recreational runners

at comfortable and high speeds. The groups considered were; a barefoot group (BAR) and a cadence group (CAD) who were compared to a control group (CON). The hypothesis of the present study was that running barefoot and increasing cadence by 10% might induce certain similar changes in ankle, knee, and hip kinematics.

METHODS AND MATERIALS

Experimental design

The effect of the two running retraining programs were explored through changes from baseline in the angles and joint excursions of ankle, knee, hip, pelvis and trunk, and angular velocities of ankle, knee and hip. These variables were measured at two different running speeds: i) comfortable speed (CS) which was self-selected by the runners in the warm-up, and ii) high speed (HS), which was self-selected by considering the best 5 km time in the current season of the participant, adapted from Latorre et al. (23). The participants visited the laboratory twice, at baseline (pretest) and at the end of the running retraining program (post-test). Participants were asked to not perform any heavy physical exertion for 72 hours prior to data collection.

Participants

A total of 103 runners, with no previous experience of running retraining programs, were initially recruited from local running clubs, and randomly assigned to one of three groups; BAR, CAD and CON, see Figure 1. Inclusion criteria were: all the participants were healthy, had regularly participated in running training with a minimum frequency of three sessions per week over the last two years, and had no history of injury in the previous six months that had limited their training. Exclusion criteria were, participants with cardiorespiratory diseases such as asthma, allergies, diabetes, or other cardiac pathologies were not included. A simple blind randomisation method was used to establish the experimental groups. This study conformed to the Declaration of Helsinki (2013) and was approved by the Ethics Committee at the University of XXXX (No. XXXXX). Participants were informed about the study and signed a consent form. Those

participants who discontinued the study were due to injuries unrelated to the retraining programs or personal reasons.

*** FIGURE 1 ABOUT HERE ***

Instruments and procedures

Body height and weight were measured to the nearest 0.1 kg and 0.1 cm respectively (SECA Instruments, Germany), and body mass index (kg/m^2) was calculated. An 8-camera motion capture system (model mvBluecoughar-XD104C, Matrix Vision GmbH, Germany) operating at 100 Hz with a resolution of 2048 x 1088 pixels and the Simi Motion software v.9.2.2. (Simi Reality Motion Systems GmbH, Germany) were used to collect kinematic data. A total of 28 markers were placed on the participants according to the International Society of Biomechanics (ISB) standard (29,30). Retroreflective markers were placed bilaterally on the acromioclavicular joint, posterior superior iliac spine, anterior superior iliac spine, greater trochanter, thigh, lateral epicondyle of the femur, medial epicondyle of the femur, shank, lateral malleolus, medial malleolus, second and third metatarsal heads and the posterior surface of the calcaneus. Two additional markers were placed on the spinous process of the 7th cervical and 8th thoracic vertebra. Following the placement of all anatomical markers, the subject was asked to stand on the treadmill for a static trial which was performed to define the anatomical coordinate systems for the foot, shank, thigh, pelvis and trunk segments (29,30).

Before the running test, the participants were asked to run consistently on a professional treadmill (Woodway Pro XL, Waukesha, WI, USA) for an 8 minute warm-up at their self-selected comfortable speed (CS) and wearing their own running shoes (26). The indications for the CS condition were: “Run comfortably and non-stop at a speed that allows you to speak and breathe easily”. Once the CS was selected and the warm-up completed, participants were instructed to run consistently for 120 s and the data collection was carried out during the last 15 s. HS was defined by participants’ best 5 km pace in the current season (26). To control for the potential effect of fatigue during the running test, intensity was measured using the Borg’s 0–10 scale (31),

and were reported as: very light (<2), light (2-3.9), moderate (4-6.9), vigorous (7-8.9), very hard (9-9.9) and maximum effort (10). For each participant, there was no more than one week between data collection and the intervention period (start and end).

Kinematics data processing

Visual 3D software (v6; C-Motion, Rockville, MD) was used to compute the 3D running kinematic variables. The raw data were filtered with a cut-off frequency of 8 Hz using a fourth-order low-pass Butterworth filter (32). Footstrike was defined using the technique described by Handsaker et al. (33), which was cross checked against the video recordings. This algorithm involves using the peak vertical acceleration of the marker placed on the posterior surface of the calcaneus for rearfoot strike runners, and the marker placed on the second and third metatarsal heads for forefoot strike runners. Prior to the application of the algorithm, the runners were classified according to their footstrike pattern (23). An 8-segment model of the lower limb and trunk was then constructed (29,30). Angles, velocities and excursions of the joints and segments (ankle, knee, hip, pelvis and trunk) were calculated for the sagittal plane, see Figure 2. Peak and footstrike values were obtained for joint and segment angles and velocities. Using the peak values and the footstrike values, the excursions were calculated (e.g., from footstrike to peak knee extension, or from peak knee flexion to peak knee extension, ...).

*** FIGURE 2 ABOUT HERE ***

Retraining program

The retraining programs were implemented by coaches and supervised by the principal researcher with regular random visits to the track each week to check compliance. The retraining programs consisted of three retraining sessions per week with the same amount of time and similar workload, adapted from Latorre et al. (23), Table 1. The BAR group performed a progression of barefoot runs on a soft, flat, non-slip grass surface (i.e., a football pitch). The CAD group was asked to land to the beat of a metronome which increased by 10% of their baseline cadence (at pre-test) at a CS. This speed was controlled on a treadmill, or using a GPS, or by lap time running

on a 400-metre track, depending on the runners' preference or ability to maintain their CS. Both groups received a weekly training diary, setting a minimum of 85% compliance, and monitoring the intensity of the sessions using the Borg scale from 0 to 10 (31). The CON group was allowed to run at a comfortable speed and perform running drills (i.e. skipping, high knees, etc.) while the BAR and CAD groups performed their retraining sessions for the same amount of time. All three groups were asked to keep their weekly volume unchanged. All participants were advised to decrease the intensity or even abandon the retraining program if pain or injury occurred. Training loads and habits were maintained by all groups.

*** TABLE 1 ABOUT HERE ***

Statistical Analysis

The distribution and homogeneity of all data were tested before analysis using the Kolmogorov-Smirnov and Levene's tests respectively, and all data were found to be suitable for parametric testing. Descriptive data were reported using the mean and standard deviations (SD). A 3 x 2 mixed model ANOVA was conducted to examine the effects of time (pre-test and post-test) and groups (BAR, CAD and CON) for each variable. Paired t-tests were used as post-hoc Bonferroni tests when a significant interaction between groups and time was detected. The significance level was set at $p < 0.05$. Additionally, effect sizes for group differences were expressed using Cohen's d (34) and were reported as: trivial (<0.2), small ($0.2-0.49$), medium ($0.5-0.79$), and large (≥ 0.8) (34). All data were analysed using SPSS, v.25.0 for Windows (SPSS Inc, Chicago, USA). To verify that there were no differences between groups at baseline, a one-way ANOVA was used for continuous variables and a chi-square test (χ^2) was used for the sex variable (dichotomous).

RESULTS

Seventy out of the 103 participants who were randomly assigned underwent analysis. The retraining program was successfully completed by 79 individuals with a minimal attendance rate of 85% (BAR group = 86% and CAD group = 86%). Technical issues resulted in 9 participants

being excluded from the 3D tracking analysis. During each acquisition period, none of the participants indicated a score higher than 6 out of 10 on the Borg scale while performing the running protocol. Average scores on the Borg scale over the ten weeks were 3.6 for the BAR group, 3.5 for the CAD group and 3.5 for the CON group. There were no differences for any of the demographic and training characteristics between the three groups (Table 2). Supplementary Tables 1 and 2 show pre- and post-test means and standard deviations, and mean difference 95% confidence intervals for joint and segment angles, velocities and excursions at both speeds.

*** TABLE 2 ABOUT HERE ***

A significant Time x Group interaction effect was seen for joint and segment angles at both speeds, see Table 3. Further post-hoc paired t-tests showed that the BAR and CAD groups increased knee flexion and hip flexion at footstrike, peak hip flexion during stance and flight phase, and pelvic anterior tilt (maximum and minimum) at both speeds; and decreased peak hip extension during flight phase (at both speeds) and peak knee extension during flight phase (at comfortable speed) after retraining, see Figure 3, Figure 4, and Table 4. Additionally, the BAR group increased anterior trunk tilt (maximum and minimum, at both speeds), decreased ankle dorsiflexion at footstrike (at both speeds) and peak ankle dorsiflexion during flight phase (at high speed) after retraining. In contrast, the CON group decreased hip flexion at footstrike (at high speed), peak hip extension during the flight phase (at both speeds), pelvic anterior tilt (maximum and minimum, at comfortable speed), maximum trunk anterior tilt (at comfortable speed), and minimum trunk anterior tilt (at both speeds) after the two time points.

*** TABLE 3 ABOUT HERE ***

*** FIGURE 3 ABOUT HERE ***

*** FIGURE 4 ABOUT HERE ***

*** TABLE 4 ABOUT HERE ***

A significant Time x Group interaction effect was seen for joint angular velocities at both speeds see Table 3. Further post-hoc paired t-tests showed that the BAR and CAD groups increased peak ankle dorsiflexion velocity during stance phase (at comfortable speed and only for the BAR group at high speed); the CAD group decreased peak hip extension velocity during stance phase (at comfortable speed); and finally, the CON group decreased peak ankle dorsiflexion velocity during stance phase (at both speeds), and increased peak ankle plantarflexion velocity during stance phase (at high speed), after the two time points, see Table 4.

A significant Time x Group interaction effect was observed for joint excursions at both speeds, see Table 3. Further post-hoc paired t-tests, showed that the BAR and CAD groups decreased knee flexion excursion in stance phase (at both speed); the BAR group decreased hip extension excursion from stance to flight phase (at comfortable speed) and increased ankle dorsiflexion excursion during stance phase (at high speed); and finally, the CON group decreased ankle dorsiflexion excursion during stance phase (at high speed), after the two time points, see Table 4.

DISCUSSION

The main findings of this study were: i) the BAR and CAD groups increased knee and hip flexion at footstrike and stance phase, decreased knee and hip extension in flight phase, and increased anterior pelvic tilt with moderate to large effect sizes at both speeds; and ii) the BAR group significantly reduced ankle dorsiflexion and increased trunk anterior tilt with a small or moderate effect sizes at both speeds. Both retraining programs produced significant kinematic changes in lower limbs and trunk running kinematics with large effect sizes. No running-related injuries were reported during the running retraining programs. The average intensity over the 10 weeks of the retraining programs was light to moderate, between 3 and 4 on the Borg scale for both the BAR and CAD groups (31). The results of the study confirmed the hypothesis that barefoot running and a 10% increase in cadence produced similar changes in ankle, knee and hip kinematics at comfortable and high speeds, although only barefoot running retraining produced changes for the

trunk. To our knowledge, this is the first study to investigate the effects of two retraining programs on lower limb and trunk running kinematics in a similar cohort of recreational endurance runners. Previous findings from immersive programs for barefoot running or with habitual barefoot runners showed an increase in knee flexion at footstrike and at 10% of the stance, and a decrease in peak knee flexion at mid-stance, as well as a decrease in foot angle at footstrike (17,18,20,26), and a decrease in hip flexion at footstrike (35). The knee and ankle appear to be the two joints most sensitive to changes after barefoot running. Compared to previous transitions to barefoot running, our results are consistent as runners increased knee flexion at footstrike, and decreased ankle dorsiflexion. However, peak knee flexion did not decrease during the stance phase for the BAR group. The decrease in knee joint excursion from initial contact to peak flexion found at both speeds may indicate less eccentric knee movement during the first half of the contact phase due to the reduced range of motion. In addition, our results showed an increase in hip flexion, anterior pelvic tilt and anterior trunk tilt during the running cycle. This is a finding that has not been mentioned in several barefoot running programs, mainly because the kinematics of the pelvis and trunk have not been studied, focusing on the foot, ankle and knee (17,18,25). Nevertheless, and similar to our findings, a greater hip flexion, greater knee flexion and less dorsal ankle flexion at footstrike has been observed in habitual barefoot runners (16), or running with a forefoot strike (14). It should be noted that during our intervention the participants were not completely immersed in barefoot running, but combined periods of barefoot running (i.e., warm-up and cool-down) with their running training in shoes and were also assessed in shoes during data collection. Therefore, the BAR group, who completed a program of barefoot running periods with shod running, showed certain similar running performance to habitual barefoot runners when assessed with shod running.

The 10-week program based on increasing the basal cadence using a mobile metronome (CAD group) showed an increase in knee flexion at footstrike, and an increase in hip flexion and anterior pelvic tilt throughout the running cycle. These results are in agreement with those of Heiderscheid et al. (13) who found an increase in knee flexion at footstrike. And also with Lenhart et al. (12)

who also found an increase in knee flexion at footstrike, in addition to a decrease a decrease in knee flexion and ankle flexion (decrease in dorsiflexion) at mid-phase. However, we found a reduction in knee flexion excursion during stance at both speeds for the CAD group, which may help to reduce the eccentric load on the knee as this range of motion was reduced. None of these studies found kinematic changes at the hip, and they did not measure pelvic kinematics. However, Lenhart et al. (12) observed a reduction in the muscle activity of the hip extensors (e.g. gluteus maximus), and our results showed a reduction in the peak velocity of hip extension, which could be related to a lower energy generation during the propulsion phase (13). Our results also showed an increase in hip flexion during running and an anterior trunk limb, both kinematic changes were previously associated by Huang et. al. (15). after a short-term approach. They also reported less awkwardness and effort when running on non-heel strike than on heel strike, which may be related to less ankle dorsiflexion at foot strike. Comparing our results with a non-laboratory setting using a mobile metronome over 12 weeks (11), knee flexion at mid-stance was reduced, but no changes in knee flexion at footstrike or hip kinematics were found. Our findings suggest that although there are some common kinematic changes between studies. There is no predominant pattern when comparing our findings with those of other studies and further research is needed.

As mentioned above, the two 10-week retraining programs, i.e., the 10% cadence increase and barefoot running, share several kinematic changes at the knee, hip and pelvis. These common kinematic changes have been associated with a 14% reduction in peak knee force and reduced peak muscle extensor forces at the hip, knee and ankle joints (12), as well as reduced mechanical energy absorption at the knee and hip (13). Greater anterior trunk tilt, observed only after the barefoot running retraining program, has been associated with less patellofemoral joint pain (27,28) and may reduce lumbopelvic loading (36). The authors of the present study believe that pelvic anteversion can be a movement associated with anterior trunk tilt. Teng H. and Power C. observed a reduced dependence on knee extensor moments during stance by increasing hip extensor moments using a more anterior trunk tilt (28). Therefore, these changes have been suggested to be strong predictors of a reduction in patellofemoral joint loading (12,13,17,18), and

the prevention of high impact injuries, such as, patellofemoral pain syndrome (37), iliotibial band syndrome (38) and tibial stress fractures (39). However, caution should be exercised in this regard as the current study did not measure kinetic parameters.

In the CON group, changes were found in the opposite direction to the two running retraining programmes. A more extended hip, posterior pelvic tilt and posterior trunk tilt with a small to moderate effect size. These kinematic changes have been observed in habitual shod runners (16). As the runners in the CON group had a short running experience (3.0 ± 1.2 years), they did not maintain their running pattern, but slightly changed some parameters towards the pattern observed in habitual shod runners, with a small effect size.

It is noteworthy that considering that both running retraining programs were performed at a comfortable speed, our results showed very similar changes in lower limb kinematics in the high speed condition. This confirms that not only was the new motor pattern acquired at the same speed as the retraining sessions, but also that the learning was transferred to higher running speeds for both retraining programs. Two studies have examined the effects of similar retraining programs performed at a comfortable and high speeds on the foot kinematics and spatiotemporal parameters (23,26). To our knowledge, this is the first study to examine changes in ankle, knee, hip, pelvis and trunk kinematics using a three-dimensional motion capture system for these two retraining programs. These retraining programs may induce adaptations in musculoskeletal structures that could affect running effectiveness in real race conditions, in the same way that short periods of barefoot running at a comfortable speed have induced changes in the shod running at a comfortable and high-speed running in these recreational runners, but further research is needed.

Finally, there are some limitations to consider. First, the effects of two 10-week retraining programs were studied in healthy recreational runners a low training load. We should be cautious about their effects on injured runners or on experienced long-distance runners with a high training load. Second, although the main focus was on sagittal plane kinematics, information on other planes of motion, ground reaction forces, joint moments or muscle activation remains unknown.

327 Third, the effects of two 10-week retraining programs were evaluated, but their long-term effects
328 (e.g. a 12-month follow-up) were not assessed.

329 From a practical approach, both retraining programs could be useful in non-laboratory settings,
330 using clinically feasible and simple methods to induce changes in lower limb and trunk kinematics
331 in recreational runners towards a running pattern that could reduce the mechanical risk of injury
332 associated with excessive knee loading. These two programs differ in certain characteristics that
333 may lead us to choose one option over the other. Barefoot running does not require the use of a
334 mobile phone and a digital metronome. Nevertheless, it may not be recommended for runners
335 who need plantar insoles, and barefoot running may not be recommended for runners with certain
336 pathologies. As for the cadence increase program, it can be used with the runner's footwear and
337 possibly with their insoles, although this use was not evaluated in this study. Another slight
338 difference is that the barefoot running programme included higher intensities for short periods in
339 addition to comfortable speeds. However, the cadence based intervention does not allow for these
340 speed changes as cadence is dependent on running speed.

341 **CONCLUSIONS**

342 A progression of periods of barefoot running and periods of 10% increase in cadence, both
343 performed at a comfortable running speed, showed an increase in knee flexion at footstrike and
344 hip angle flexion and pelvic anterior tilt during the running cycle with a moderate to large effect
345 sizes at both comfortable and high running speeds after 10 weeks. In addition, the progression
346 of barefoot running periods showed a significant reduction in ankle dorsiflexion at footstrike
347 and an increase in anterior trunk tilt with a small to moderate effect size at both speeds after the
348 10-week retraining program. Barefoot running was slightly more effective as a running
349 retraining program than increasing cadence with a digital metronome. This may be useful for
350 reducing knee risk factors and increasing running efficiency, but further research is needed to
351 explore the long-term effects.

Acknowledgments: The authors would like to acknowledge the study participants involved in recruitment.

Author Contributions: A.M.-M., V.M.S.-H. and P.A.L.-R. defined the experimental design and conceptualized the approach. A.M.-M., G.D.-G and E.M.-P. collected the data. A.M.-M., P.A.L.-R. and J.R. carried out the statistical analysis. A.M.-M. wrote the paper. All authors reviewed the manuscript for scientific content. All authors have read and agreed to the published version of the manuscript.

Funding: The authors declare no funding has been received for this research.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the University of XXXXX (No. XXXXXX)

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Conflicts of interest: The authors declare no conflict of interest.

368 **REFERENCES**

- 369 1. Carmack MA, Martens R. Measuring Commitment to Running: A Survey of Runners'
370 Attitudes and Mental States. *J Sport Psychol.* 2016;
- 371 2. Urbaneja JS, Farias EI. Trail running in Spain. Origin, evolution and current situation;
372 natural areas. *Retos.* 2018;2041(33):123–8.
- 373 3. Buist I, Bredeweg SW, Lemmink K a PM, van Mechelen W, Diercks RL. Predictors of
374 Running-Related Injuries in Novice Runners Enrolled in a Systematic Training Program.
375 *Am J Sports Med.* 2010 Feb 4;38(2):273–80.
- 376 4. Daoud AI, Geissler GJ, Wang F, Saretsky J, Daoud Y a., Lieberman DE. Foot strike and
377 injury rates in endurance runners: A retrospective study. *Med Sci Sports Exerc.*
378 2012;44(7):1325–34.
- 379 5. Hollander K, Baumann A, Zech A, Verhagen E. Prospective monitoring of health
380 problems among recreational runners preparing for a half marathon. *BMJ Open Sport*
381 *Exerc Med.* 2018 Jan;4(1):e000308.
- 382 6. Gaitonde DY, Ericksen A, Robbins RC. Patellofemoral Pain Syndrome. *Am Fam*
383 *Physician.* 2019;99(2):88–94.
- 384 7. Lankhorst NE, Bierma-Zeinstra SMA, van Middelkoop M. Risk Factors for
385 Patellofemoral Pain Syndrome: A Systematic Review. *J Orthop Sport Phys Ther.* 2012
386 Feb;42(2):81-A12.
- 387 8. Van Gent RN, Siem D, Van Middelkoop M, Van Os AG, Bierma-Zeinstra SMA, Koes
388 BW. Incidence and determinants of lower extremity running injuries in long distance
389 runners: A systematic review. *Br J Sports Med.* 2007;41(8):469–80.
- 390 9. Winstein CJ. Movement science series knowledge of results and motor implications for
391 physical therapy learning. *Mov Sci Ser.* 1991;71(2):140–9.

- 392 10. Bramah C, Preece SJ, Gill N, Herrington L. A 10% Increase in Step Rate Improves
393 Running Kinematics and Clinical Outcomes in Runners With Patellofemoral Pain at 4
394 Weeks and 3 Months. *Am J Sports Med.* 2019;47(14):3406–13.
- 395 11. Wang J, Luo Z, Dai B, Fu W. Effects of 12-week cadence retraining on impact peak,
396 load rates and lower extremity biomechanics in running. *PeerJ.* 2020 Aug 24;8:e9813.
- 397 12. Lenhart RL, Thelen DG, Wille CM, Chumanov ES, Heiderscheit BC. Increasing
398 Running Step Rate Reduces Patellofemoral Joint Forces. *Med Sci Sport Exerc.* 2014
399 Mar;46(3):557–64.
- 400 13. Heiderscheit BC, Chumanov ES, Michalski MP, Wille CM, Ryan MB. Effects of step
401 rate manipulation on joint mechanics during running. *Med Sci Sports Exerc.*
402 2011;43(2):296–302.
- 403 14. dos Santos AF, Nakagawa TH, Lessi GC, Luz BC, Matsuo HTM, Nakashima GY, et al.
404 Effects of three gait retraining techniques in runners with patellofemoral pain. *Phys Ther*
405 *Sport.* 2019;36:92–100.
- 406 15. Huang Y, Xia H, Chen G, Cheng S, Cheung RTH, Shull PB. Foot strike pattern, step
407 rate, and trunk posture combined gait modifications to reduce impact loading during
408 running. *J Biomech.* 2019;86:102–9.
- 409 16. Lieberman DE, Venkadesan M, Werbel WA, Daoud AI, D’Andrea S, Davis IS, et al.
410 Foot strike patterns and collision forces in habitually barefoot versus shod runners.
411 *Nature.* 2010 Jan 27;463(7280):531–5.
- 412 17. Azevedo AP da S, Mezêncio B, Amadio AC, Serrão JC. 16 Weeks of Progressive
413 Barefoot Running Training Changes Impact Force and Muscle Activation in Habitual
414 Shod Runners. Fisher G, editor. *PLoS One.* 2016 Dec 1;11(12):e0167234.
- 415 18. Tam N, Tucker R, Astephen Wilson JL. Individual Responses to a Barefoot Running
416 Program. *Am J Sports Med.* 2016 Mar 7;44(3):777–84.

- 417 19. Hollander K, Liebl D, Meining S, Mattes K, Willwacher S, Zech A. Adaptation of
418 Running Biomechanics to Repeated Barefoot Running: A Randomized Controlled Study.
419 Am J Sports Med. 2019 Jul 5;47(8):1975–83.
- 420 20. Bonacci J, Saunders PU, Hicks A, Rantalainen T, Vicenzino BT, Spratford W. Running
421 in a minimalist and lightweight shoe is not the same as running barefoot: A
422 biomechanical study. Br J Sports Med. 2013;47(6):387–92.
- 423 21. Hall JPL, Barton C, Jones PR, Morrissey D. The biomechanical differences between
424 barefoot and shod distance running: A systematic review and preliminary meta-analysis.
425 Sport Med. 2013;43(12):1335–53.
- 426 22. Murphy K, Curry EJ, Matzkin EG. Barefoot running: Does it prevent injuries? Sport
427 Med. 2013;43(11):1131–8.
- 428 23. Latorre-Román PA, García-Pinillos F, Soto-Hermoso VM, Muñoz-Jiménez M. Effects of
429 12 weeks of barefoot running on foot strike patterns, inversion–eversion and foot
430 rotation in long-distance runners. J Sport Heal Sci. 2019 Nov;8(6):579–84.
- 431 24. Hollander K, Liebl D, Meining S, Mattes K, Willwacher S, Zech A. Adaptation of
432 Running Biomechanics to Repeated Barefoot Running: A Randomized Controlled Study.
433 Am J Sports Med. 2019;47(8):1975–83.
- 434 25. Tam N, Tucker R, Astephen Wilson J, Santos-Concejero J. Effect on Oxygen Cost of
435 Transport from 8-Weeks of Progressive Training with Barefoot Running. Int J Sports
436 Med. 2015 Sep 2;36(13):1100–5.
- 437 26. Molina-Molina A, Latorre-Román PÁ, Mercado-Palomino E, Delgado-García G,
438 Richards J, Soto-Hermoso VM. The effect of two retraining programs, barefoot running
439 vs increasing cadence, on kinematic parameters: A randomized controlled trial. Scand J
440 Med Sci Sports. 2022 Mar 9;32(3):533–42.
- 441 27. Teng HL, Powers CM. Sagittal plane trunk posture influences patellofemoral joint stress

442 during running. *J Orthop Sports Phys Ther.* 2014;44(10):785–92.

443 28. Teng HL, Powers CM. Hip-extensor strength, trunk posture, and use of the knee-
 444 extensor muscles during running. *J Athl Train.* 2016;51(7):519–24.

445 29. Wu G, Siegler S, Allard P, Kirtley C, Leardini A, Rosenbaum D, et al. ISB
 446 recommendation on definitions of joint coordinate system of various joints for the
 447 reporting of human joint motion--part I: ankle, hip, and spine. International Society of
 448 Biomechanics. *J Biomech.* 2002 Apr;35(4):543–8.

449 30. Wu G, Van Der Helm FCT, Veeger HEJ, Makhsous M, Van Roy P, Anglin C, et al. ISB
 450 recommendation on definitions of joint coordinate systems of various joints for the
 451 reporting of human joint motion - Part II: Shoulder, elbow, wrist and hand. *J Biomech.*
 452 2005;38(5):981–92.

453 31. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sport Exerc.* 1982;

454 32. Chan ZYS, Zhang JH, Ferber R, Shum G, Cheung RTH. The effects of midfoot strike
 455 gait retraining on impact loading and joint stiffness. *Phys Ther Sport.* 2020;42:139–45.

456 33. Handsaker JC, Forrester SE, Folland JP, Black MI, Allen SJ. A kinematic algorithm to
 457 identify gait events during running at different speeds and with different footstrike types.
 458 *J Biomech.* 2016;49(16):4128–33.

459 34. Cohen J. Statistical power analysis for the behavioral sciences. Vol. 2nd, Statistical
 460 Power Analysis for the Behavioral Sciences. 1988. 567 p.

461 35. McCarthy C, Fleming N, Donne B, Blanksby B. Barefoot running and hip kinematics:
 462 Good news for the knee? *Med Sci Sports Exerc.* 2015;47(5):1009–16.

463 36. Seay J, Selbie WS, Hamill J. In vivo lumbo-sacral forces and moments during constant
 464 speed running at different stride lengths. *J Sports Sci.* 2008 Dec 15;26(14):1519–29.

465 37. Willson JD, Davis IS. Lower extremity mechanics of females with and without
 466 patellofemoral pain across activities with progressively greater task demands. *Clin*

467 Biomech. 2008 Feb;23(2):203–11.

468 38. Hamill J, Miller R, Noehren B, Davis I. A prospective study of iliotibial band strain in
469 runners. Clin Biomech. 2008 Oct;23(8):1018–25.

470 39. Pohl MB, Mullineaux DR, Milner CE, Hamill J, Davis IS. Biomechanical predictors of
471 retrospective tibial stress fractures in runners. J Biomech. 2008;41(6):1160–5.

472

473

TABLES

Table 1: Running volumes and speeds of the 10-week running retraining programs.							
	Weeks						
	1 - 2	3 - 4	5 - 6	7	8	9	10
Barefoot retraining group	15' CS	20' CS	25' CS	30' CS	10' CS 10' MS 10' CS	35' CS	10' CS 10' MS 10' CS 5' MS 5' CS
Cadence retraining group	15' CS	20' CS	25' CS	30' CS	30' CS	35' CS	40' CS
Weekly volume by group	45'	60'	75'	90'	90'	105'	120'

Comfortable speed (CS); Medium speed (MS). In weeks 4, 6, and 9, the BAR group completed five sets of 80-meter progressive sprints.

Table 2: Demographic, anthropometric and training characteristics of runners expressed as mean (SD)				
Variable	BAR (n = 23)	CAD (n = 23)	CON (n = 24)	P value
Gender (percentage)	43.5% females, 56.5% males	47.8% females, 52.2% males	29.1% females, 70.9% males	0.432
Age (year old)	31.4 (7.4)	29.4 (7.4)	29.2 (7.1)	0.543
Height (cm)	175.3 (10.4)	173.1 (7.2)	173.1 (6.7)	0.593
Body mass (kg)	73.0 (12.2)	69.8 (11.3)	70.2 (11.5)	0.598
Body mass index (kg/m ²)	23.7 (3.0)	23.2 (2.5)	23.3 (2.8)	0.795
Comfortable speed (km/h)	9.9 (1.1)	10.2 (1.0)	10.5 (0.9)	0.150
High speed (km/h)	14.0 (2.0)	14.2 (1.9)	15.1 (1.4)	0.095
Running experience (years)	3.4 (1.3)	3.4 (1.7)	3.0 (1.2)	0.457
Running's sessions (n/wk.)	3.8 (1.0)	3.9 (1.1)	3.0 (1.0)	0.625
Workload per week (kms)	27.0 (14.2)	21.9 (11.7)	26.4 (9.3)	0.287
Competitions per year (n)	10.6 (7.9)	8.0 (6.6)	7.3 (4.3)	0.202

Barefoot retraining group (BAR), cadence retraining group (CAD) and control group (CON). P < 0.05

Table 3. Interaction effects Time (pre-test and post-test) vs. Groups (BAR, CAD and CON) of the mixed model 2 x 3 ANOVA for each variable at both speeds.

Speed	Phase	Variable	<i>P</i> value Group x time	F-test (df ₁ , df ₂)	η ²
Comfortable speed	Stance Phase	Ankle dorsiflexion at footstrike (°)	0.049	3.163 (2, 65)	0.089 M
		Knee flexion at footstrike (°)	<0.001	8.899 (2, 65)	0.215 L
		Hip flexion at footstrike (°)	<0.001	12.507 (2, 63)	0.284 L
		Peak hip flexion (°)	<0.001	11.341 (2, 62)	0.268 L
		Knee flexion excursion (°)	0.004	5.955 (2, 64)	0.157 L
		Peak ankle dorsiflexion velocity (°/s)	0.001	8.230 (2, 62)	0.210 L
		Peak hip extension velocity (°/s)	0.043	3.304 (2, 64)	0.094 M
	Flight Phase	Peak knee extension (°)	0.003	3.609 (2, 64)	0.101 M
		Peak hip extension (°)	<0.001	16.665 (2, 63)	0.346 L
		Peak hip flexion (°)	<0.001	10.327 (2, 63)	0.247 L
	Full cycle	Maximum pelvic anterior tilt (°)	<0.001	14.347 (2, 64)	0.310 L
		Minimum pelvic anterior tilt (°)	<0.001	16.960 (2, 64)	0.346 L
		Maximum trunk anterior tilt (°)	<0.001	11.457 (2, 63)	0.267 L
		Minimum trunk anterior tilt (°)	<0.001	9.149 (2, 63)	0.225 L
		Hip extension excursion (°)	0.010	5.006 (2, 63)	0.139 M
High speed	Stance Phase	Ankle dorsiflexion at footstrike (°)	0.004	5.912 (2, 63)	0.158 L
		Knee flexion at footstrike (°)	0.001	7.963 (2, 62)	0.204 L
		Hip flexion at footstrike (°)	<0.001	9.280 (2, 62)	0.230 L
		Peak hip flexion (°)	0.001	8.212 (2, 62)	0.209 L
		Ankle dorsiflexion excursion (°)	<0.001	9.236 (2, 63)	0.227 L
		Knee flexion excursion (°)	0.012	4.740 (2, 62)	0.133 M
		Peak ankle dorsiflexion velocity (°/s)	<0.001	9.416 (2, 62)	0.233 L
		Peak ankle plantarflexion velocity (°/s)	0.021	4.140 (2, 62)	0.120 M
	Flight Phase	Peak ankle dorsiflexion (°)	0.005	5.772 (2, 63)	0.155 L
		Peak hip extension (°)	<0.001	13.999 (2, 62)	0.311 L
		Peak hip flexion (°)	0.001	8.249 (2, 62)	0.210 L
	Full cycle	Maximum pelvic anterior tilt (°)	<0.001	9.317 (2, 61)	0.234 L
		Minimum pelvic anterior tilt (°)	0.001	8.193 (2, 62)	0.209 L
		Maximum trunk anterior tilt (°)	0.001	7.542 (2, 59)	0.204 L
		Minimum trunk anterior tilt (°)	<0.001	9.218 (2, 59)	0.228 L
Barefoot group (BAR); Cadence group (CAD); Control group (CON); Degrees of freedom (df); Partial eta squared (η ²); Small effect size = 0.01 (S); Medium effect size between = 0.06 (M); Large effect size = 0.14 (L)					

479

480

Table 4. Post-hoc paired t-tests for joint and segment angles, velocities and excursions before (pre-test) and after (post-test) the 10-week programs at comfortable and high speed.

Speed	Phase	Variable	Barefoot retraining group		Cadence retraining group		Control group	
			Mean difference	P value (Cohen's d)	Mean difference	P value (Cohen's d)	Mean difference	P value (Cohen's d)
Comfortable speed	Stance Phase	Ankle dorsiflexion at footstrike (°)	-2.09 ± 0.91	0.024 (0.40)	-1.16 ± 0.87	0.186 (0.26)	0.93 ± 0.85	0.159 (0.28)
		Knee flexion at footstrike (°)	5.10 ± 1.09	<0.001 (1.01)	4.70 ± 1.04	<0.001 (0.91)	-0.44 ± 1.02	0.524 (0.13)
		Hip flexion at footstrike (°)	6.53 ± 1.65	<0.001 (0.96)	6.76 ± 1.54	<0.001 (1.06)	-2.90 ± 1.54	0.064 (0.46)
		Peak hip flexion (°)	6.28 ± 1.70	<0.001 (0.88)	6.21 ± 1.62	<0.001 (1.00)	-3.16 ± 1.59	0.051 (0.50)
		Knee flexion excursion (°)	-2.98 ± 1.08	0.009 (0.67)	-3.50 ± 1.06	0.002 (0.64)	1.10 ± 1.01	0.282 (0.20)
		Peak ankle dorsiflexion velocity (°/s)	30.68 ± 9.87	0.003 (0.50)	20.64 ± 9.87	0.041 (0.39)	-21.19 ± 9.43	0.028 (0.57)
		Peak hip extension velocity (°/s)	8.90 ± 7.13	0.117 (0.86)	-14.53 ± 6.65	0.033 (0.30)	3.79 ± 6.51	0.563 (0.49)
	Flight Phase	Peak knee extension (°)	3.19 ± 0.84	<0.001 (0.86)	2.07 ± 0.82	0.014 (0.60)	0.16 ± 0.78	0.844 (0.03)
		Peak hip extension (°)	8.01 ± 1.66	<0.001 (1.27)	6.63 ± 1.59	<0.001 (1.08)	-3.65 ± 1.52	0.019 (0.48)
		Peak hip flexion (°)	8.25 ± 1.92	<0.001 (1.11)	6.92 ± 1.83	<0.001 (0.92)	-2.39 ± 1.75	0.178 (0.33)
	Full cycle	Maximum pelvic anterior tilt (°)	5.58 ± 1.40	<0.001 (1.01)	5.56 ± 1.36	<0.001 (1.10)	-3.16 ± 0.12	0.018 (0.53)
		Minimum pelvic anterior tilt (°)	5.82 ± 1.40	<0.001 (1.04)	6.00 ± 1.36	<0.001 (1.15)	-3.58 ± 0.08	0.008 (0.55)
		Maximum trunk anterior tilt (°)	2.05 ± 0.56	0.001 (0.59)	0.91 ± 0.54	0.094 (0.27)	-1.48 ± -0.42	0.005 (0.41)
		Minimum trunk anterior tilt (°)	1.57 ± 0.54	0.005 (0.39)	0.38 ± 0.51	0.458 (0.11)	-1.50 ± -0.26	0.003 (0.50)
		Hip extension excursion (°)	-1.47 ± 0.61	0.018 (0.33)	-0.17 ± 0.58	0.748 (0.02)	1.15 ± 0.57	0.470 (0.02)
High speed	Stance Phase	Ankle dorsiflexion at footstrike (°)	-3.19 ± 0.95	0.001 (0.58)	-1.41 ± 0.93	0.134 (0.29)	1.30 ± 0.91	0.140 (0.28)
		Knee flexion at footstrike (°)	4.81 ± 1.08	<0.001 (0.85)	3.77 ± 1.03	0.0001 (0.71)	-0.64 ± 1.00	0.524 (0.13)
		Hip flexion at footstrike (°)	6.55 ± 1.83	0.001 (0.85)	4.73 ± 1.75	0.009 (0.66)	-3.43 ± 0.77	0.049 (0.42)
		Peak hip flexion (°)	5.71 ± 1.80	0.002 (0.70)	4.66 ± 1.71	0.008 (0.65)	-3.24 ± 1.66	0.058 (0.40)
		Ankle dorsiflexion excursion (°)	1.64 ± 0.62	0.009 (0.43)	0.83 ± 0.61	0.164 (0.30)	-1.83 ± 0.59	0.003 (0.53)
		Knee flexion excursion (°)	-2.12 ± 1.01	0.040 (0.38)	-2.15 ± 0.96	0.029 (0.35)	1.47 ± 0.92	0.123 (0.24)
		Peak ankle dorsiflexion velocity (°/s)	52.44 ± 14.24	< 0.001 (0.78)	28.28 ± 14.24	0.051 (0.29)	-30.22 ± 13.60	0.030 (0.45)
		Peak ankle plantarflexion velocity (°/s)	-27.03 ± 15.60	0.088 (0.64)	-18.11 ± 15.24	0.240 (0.36)	31.75 ± 15.60	0.046 (0.59)
	Flight Phase	Peak ankle dorsiflexion (°)	-4.14 ± 1.06	<0.001 (0.63)	-1.13 ± 1.04	0.278 (0.24)	0.82 ± 1.01	0.421 (0.17)
		Peak hip extension (°)	7.66 ± 1.75	<0.001 (1.07)	6.13 ± 1.67	<0.001 (1.00)	-3.81 ± 1.63	0.023 (0.49)
		Peak hip flexion (°)	8.28 ± 1.87	<0.001 (0.86)	6.76 ± 1.78	<0.001 (0.85)	-1.23 ± 1.74	0.484 (0.14)
	Full cycle	Maximum pelvic anterior tilt (°)	6.69 ± 1.67	<0.001 (1.01)	4.16 ± 1.59	0.011 (0.76)	-2.82 ± 1.59	0.081 (0.46)
		Minimum pelvic anterior tilt (°)	6.98 ± 1.79	<0.001 (0.93)	4.48 ± 1.71	0.011 (0.75)	-2.46 ± 1.67	0.142 (0.32)
		Maximum trunk anterior tilt (°)	2.13 ± 0.66	0.002 (0.51)	1.178 ± 0.62	0.064 (0.32)	-1.21 ± 0.61	0.052 (0.30)
		Minimum trunk anterior tilt (°)	1.81 ± 0.58	0.003 (0.36)	0.61 ± 0.56	0.277 (0.15)	-1.54 ± 0.19	0.006 (0.47)

Post-hoc paired tests were used when a significant interaction effect between Time × Group was seen. Bold denotes $p < 0.05$. A positive value in the mean difference indicates an increase from the pre-test to the post-test and vice versa.

Supplementary table 1. Means (SDs) and mean difference 95% Confidence Intervals for joint and segment angles, velocities and excursions before (pre-test) and after (post-test) the 10-week programs at comfortable speed.

Phase	Variable	Barefoot retraining group			Cadence retraining group			Control group		
		Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% CI [low, high]	Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% C [low, high]	Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% CI [low, high]
Stance Phase	Ankle dorsiflexion at footstrike (°)	3.7 (5.2)	1.61 (5.21)	[0.28, 3.90]	4.31 (4.41)	3.16 (4.56)	[-0.57, 2.89]	3.13 (4.15)	4.06 (4.24)	[-2.63, 0.76]
	Knee flexion at footstrike (°)	11.69 (4.8)	16.79 (5.25)	[-7.27, -2.92]	11.49 (4.79)	16.18 (5.48)	[-6.77, -2.62]	14.93 (4.55)	14.48 (4.94)	[-1.59, 2.48]
	Hip flexion at footstrike (°)	31.73 (7.47)	38.26 (6.06)	[-9.84, -3.23]	30.94 (6.40)	37.70 (6.40)	[-9.84, -3.68]	41.06 (5.38)	38.16 (7.08)	[-0.18, 5.99]
	Peak hip flexion (°)	32.19 (7.38)	38.47 (6.88)	[-9.68, -2.88]	31.01 (6.29)	37.21 (6.07)	[-9.45, -2.96]	41.58 (5.33)	38.42 (7.19)	[-0.01, 6.33]
	Knee flexion excursion (°)	23.66 (4.14)	20.68 (4.76)	[-5.14, -0.81]	23.47 (4.60)	19.98 (6.19)	[-5.61, -0.84]	22.89 (5.76)	23.99 (5.37)	[-0.93, 3.13]
	Peak ankle dorsiflexion velocity (°/s)	208.24 (46.01)	238.92 (73.14)	[-50.42, -10.95]	208.51 (56.5)	229.15 (48.38)	[-40.38, -0.91]	220.05 (42.27)	198.86 (30.60)	[2.34, 40.05]
Flight Phase	Peak hip extension velocity (°/s)	-277.40 (52.34)	-268.50 (39.00)	[-23.15, 5.35]	-259.37 (42.80)	-273.90 (54.19)	[1.25, 27.82]	-275.60 (42.54)	-271.81 (47.70)	[-16.8, 9.22]
	Peak knee extension (°)	8.73 (3.46)	11.92 (3.93)	[-4.86, -1.51]	9.88 (3.03)	11.94 (3.81)	[-3.7, -0.43]	13.00 (4.87)	13.16 (4.70)	[-1.72, 1.41]
	Peak hip extension (°)	-6.36 (5.92)	1.64 (7.06)	[-11.33, -4.68]	-6.73 (6.35)	-0.1 (5.95)	[-9.8, -3.46]	2.09 (8.11)	-1.56 (7.06)	[0.62, 6.69]
	Peak hip flexion (°)	43.67 (5.16)	51.92 (9.14)	[-12.09, -4.4]	42.4 (7.25)	49.32 (7.78)	[-10.59, -3.25]	49.86 (6.92)	47.47 (7.40)	[-1.12, 5.9]
Full cycle	Maximum pelvic anterior tilt (°)	20.79 (5.30)	26.37 (5.76)	[2.79, 8.37]	19.43 (5.44)	24.98 (4.66)	[2.83, 8.28]	26.88 (5.89)	23.72 (6.01)	[-5.77, -0.55]
	Minimum pelvic anterior tilt (°)	12.92 (6.07)	18.73 (5.12)	[3.03, 8.61]	12.73 (5.38)	18.73 (5.03)	[3.27, 8.73]	20.36 (6.51)	16.77 (6.59)	[-6.19, -0.97]
	Maximum trunk anterior tilt (°)	24.72 (3.94)	26.77 (2.99)	[0.92, 3.17]	23.88 (3.33)	24.79 (3.46)	[-0.16, 1.98]	26.23 (3.76)	24.76 (3.34)	[-2.5, -0.45]
	Minimum trunk anterior tilt (°)	19.81 (4.62)	21.38 (3.34)	[0.49, 2.64]	19.18 (3.55)	19.56 (3.47)	[-0.64, 1.41]	21.19 (3.11)	-19.68 (2.85)	[-2.49, -0.52]
	Hip extension excursion (°)	38.09 (4.60)	36.62 (4.22)	[-2.69, -0.26]	37.49 (4.47)	37.31 (3.92)	[-1.34, 0.97]	37.89 (4.59)	37.78 (4.47)	[0.02, 2.28]


CI denotes Confidence Intervals; Bold in Dif. 95% CI denotes post-hoc paired tests by Time (pre-test vs. post-test) with $p < 0.05$.


Supplementary table 2. Means (SDs) and mean difference 95% Confidence Intervals for joint and segment angles, velocities and excursions before (pre-test) and after (post-test) the 10-week programs at high speed.

Phase	Variable	Barefoot retraining group			Cadence retraining group			Control group		
		Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% CI [low, high]	Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% C [low, high]	Pre-test Mean (SD)	Post-test Mean (SD)	Dif. 95% CI [low, high]
Stance Phase	Ankle dorsiflexion at footstrike (°)	4.06 (5.13)	0.87 (5.86)	[1.28, 5.10]	4.31 (4.58)	2.89 (5.19)	[-0.45, 3.28]	2.34 (4.74)	3.63 (4.43)	[-3.12, 0.52]
	Knee flexion at footstrike (°)	12.72 (5.23)	17.53 (6.11)	[-6.97, -2.66]	12.47 (4.75)	16.24 (5.78)	[-5.82, -1.71]	16.16 (4.40)	15.52 (5.23)	[-1.36, 2.65]
	Hip flexion at footstrike (°)	34.92 (8.75)	41.47 (6.44)	[-10.22, -2.89]	35.01 (6.94)	39.74 (7.51)	[-8.23, -1.24]	45.04 (7.78)	41.60 (8.55)	[0.02, 6.85]
	Peak hip flexion (°)	35.26 (8.90)	40.97 (7.27)	[-9.31, -2.12]	34.30 (6.77)	38.96 (7.48)	[-8.08, -1.23]	44.46 (7.71)	41.22 (8.68)	[-0.11, 6.59]
	Ankle dorsiflexion excursion (°)	16.03 (3.71)	17.70 (4.13)	[0.43, 2.91]	16.34 (2.91)	17.19 (2.67)	[-0.36, 2.07]	16.43 (2.65)	15.05 (2.60)	[-3.01, -0.64]
	Knee flexion excursion (°)	21.99 (5.51)	19.87 (5.52)	[-4.13, -0.10]	22.34 (5.72)	20.19 (6.57)	[-4.08, -0.23]	21.65 (6.47)	23.12 (5.90)	[-0.41, 3.35]
	Peak ankle dorsiflexion velocity (°/s)	224.94 (41.13)	277.38 (86.16)	[-80.90, -23.98]	233.78 (42.61)	262.06 (58.18)	[-56.74, 0.18]	271.91 (78.49)	241.69 (54.19)	[3.03, 57.42]
	Peak ankle plantarflexion velocity (°/s)	-547.10 (119.51)	-574.13 (66.16)	[-4.17, 58.23]	-571.18 (77.51)	-589.28 (60.59)	[-12.38, 48.58]	-620.68 (56.07)	-588.93 (51.22)	[-62.94, -0.55]
Flight Phase	Peak ankle dorsiflexion (°)	7.22 (6.31)	3.08 (6.76)	[2.02, 6.27]	6.87 (4.73)	5.74 (4.85)	[-0.94, 3.21]	5.22 (4.54)	6.04 (4.91)	[-2.85, 1.21]
	Peak hip extension (°)	-9.74 (7.64)	-2.09 (6.62)	[-11.15, -4.16]	-10.55 (6.39)	-4.43 (5.84)	[-9.47, -2.79]	-2.17 (7.96)	-5.98 (7.50)	[0.55, 7.08]
	Peak hip flexion (°)	53.14 (8.43)	61.41 (10.70)	[-12.01, -4.54]	52.29 (7.70)	59.05 (8.19)	[-10.32, -3.2]	59.28 (9.37)	58.05 (7.85)	[-2.26, 4.71]
Full cycle	Maximum pelvic anterior tilt (°)	20.89 (7.58)	27.57 (5.48)	[3.36, 10.02]	21.09 (5.57)	25.25 (5.34)	[0.99, 7.34]	28.44 (5.87)	25.62 (6.41)	[-5.99, 0.36]
	Minimum pelvic anterior tilt (°)	12.69 (9.19)	19.66 (5.25)	[3.39, 10.56]	14.17 (6.08)	18.65 (5.80)	[1.06, 7.90]	20.87 (8.44)	18.39 (7.10)	[-5.83, 0.86]
	Maximum trunk anterior tilt (°)	25.64 (4.91)	27.76 (3.30)	[0.81, 3.44]	24.79 (3.77)	25.97 (3.51)	[-0.07, 2.43]	26.84 (4.06)	25.63 (4.03)	[-2.43, 0.01]
	Minimum trunk anterior tilt (°)	19.22 (5.78)	21.03 (4.05)	[0.64, 2.98]	19.00 (4.12)	19.61 (3.80)	[-0.50, 1.72]	20.44 (3.19)	18.9 (3.38)	[-2.63, -0.46]

CI denotes Confidence Intervals; Bold in Dif. 95% CI denotes post-hoc paired tests by Time (pre-test vs. post-test) with $p < 0.05$.


FIGURES






486

487



488



490

491 **FIGURES CAPTIONS**

492 Figure 1. Flow diagram of participant recruitment, follow-up and inclusion for analysis.

493 Figure 2. Calculated joint and segmental angles.

494 Figure 3. Changes during the stride cycle for the joint and segmental angles after the 10-week
495 retraining programs at comfortable speed (*denotes significant difference between pre-test and
496 post-test when an interaction effect between Time $\times$ Group was seen).

497 Figure 4. Changes during the stride cycle for the joint and segmental angles after the 10-week
498 retraining programs at high speed (*denotes significant difference between pre-test and post-test
499 when and interaction effect between Time $\times$ Group was seen).