

Central Lancashire Online Knowledge (CLOK)

Title	Wilderness Medical Society Clinical Practice Guidelines for the Treatment of Acute Pain in Austere Environments: 2024 Update
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/51334/
DOI	https://doi.org/10.1177/10806032241248422
Date	2024
Citation	Fink, Patrick B., Wheeler, Albert R., Smith, William R., Brant-Zawadzki, Graham, Lieberman, James R., McIntosh, Scott E., Van Tilburg, Christopher, Wedmore, Ian S., Windsor, Jeremy et al (2024) Wilderness Medical Society Clinical Practice Guidelines for the Treatment of Acute Pain in Austere Environments: 2024 Update. Wilderness & Environmental Medicine, 35 (2). ISSN 1080-6032
Creators	Fink, Patrick B., Wheeler, Albert R., Smith, William R., Brant-Zawadzki, Graham, Lieberman, James R., McIntosh, Scott E., Van Tilburg, Christopher, Wedmore, Ian S., Windsor, Jeremy, Hofmeyr, Ross and Weber, David

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1177/10806032241248422>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

WILDERNESS MEDICAL SOCIETY CLINICAL PRACTICE GUIDELINES

Title: Wilderness Medical Society Clinical Practice Guidelines for the Treatment of Acute Pain in Austere Environments: 2024 Update

Authors/Affiliations: Patrick B Fink, MD, University of Utah Department of Emergency Medicine, Salt Lake City, UT; Albert R. Wheeler III, MD, St. John's Health Department of Emergency Medicine, Jackson, WY; William R. Smith, MD, St. John's Health Department of Emergency Medicine, Jackson, WY; Graham Brant-Zawadzki, MD, MPH, University of Utah Department of Emergency Medicine, Salt Lake City, UT; James R. Lieberman, MD, unaffiliated; Scott E. McIntosh, MD, MPH, University of Utah Department of Emergency Medicine, Salt Lake City, UT; Christopher Van Tilburg, MD, Providence Hood River Memorial Hospital, Hood River, OR; Ian S Wedmore, MD, Uniformed Services University of the Health Sciences, Bethesda, MD; Jeremy S. Windsor, MD, University of Central Lancashire, Preston, United Kingdom; Ross Hofmeyr, MMed, University of Cape Town Department of Anaesthesia and Perioperative Medicine, Cape Town, South Africa; David Weber, FP-C, Mountain Rescue Collective, LLC, Park City, UT.

Corresponding Author: Dr. Patrick Fink, University of Utah Department of Emergency Medicine, 50 N Medical Dr., Salt Lake City, UT 84132. Email: patrickfinkmd@gmail.com

Summary Tallies

Word Count (abstract): 75

Word Count (manuscript with references): 15957

Reference Count: 213

Figure Count: 1

Table Count: 1

29 Abstract:

30 The Wilderness Medical Society (WMS) convened an expert panel to develop evidence-based

31 guidelines for the management of pain in austere environments. Recommendations are graded

32 based on the quality of supporting evidence as defined by criteria put forth by the American

33 College of Chest Physicians. This is an update of the 2014 version of the WMS Practice

34 Guidelines for the Treatment of Acute Pain in Remote Environments published in Wilderness &

35 Environmental Medicine 2014; 25:41–49.

36

37 Key words: Acute Pain / Therapy; Pain Management / Methods; Wilderness Medicine /

38 Methods; Wilderness Medicine / Standards; Societies, Medical

39

40

41

Introduction

The treatment of pain is important in the management of illness and injury in any context. Beyond the ethical considerations of compassion and management of suffering, effective management of acute pain in austere settings can facilitate the evaluation, packaging, and transportation of patients whose management would otherwise be challenging or impossible without pain control.

Management of acute pain is a multifaceted challenge which can involve physical stabilization of an acute musculoskeletal injury or wound, medical management of peripherally and centrally mediated pain responses, and management of psychological distress and anxiety. Any treatment of pain carries with it risks and benefits which may be amplified when these treatments are used in an austere environment. Limited patient access, difficulty of reevaluation during transport, need to protect the patient from environmental insults, lack of monitoring equipment, and limited means to respond to adverse effects of treatments can amplify risks. Consequently, treatment of pain in an austere environment requires a thoughtful and progressive approach that minimizes harms while managing the needs of the patient and treating team.

For the purposes of this guideline, austere environments are defined as wilderness, remote, or highly resource-limited settings where access to typical hospital or EMS resources is unavailable and evacuation may be challenging or prolonged. We include the setting of wilderness, and include expedition settings and other travel to regions where parties must be self-sufficient in their medical care. We exclude from this guideline pain management in the military setting, where resources and availability of evacuation are generally greater and for which guidelines already exist. We also exclude management of acute pain in space travel, where physiology and resource limitations are truly unique. The term ‘austere’ has been chosen to

replace ‘remote’ in this guideline as it better describes a setting of limited resources and environmental challenges. When terrain, weather, or altitude are challenges to care, resources do not need to be geographically remote to be functionally unavailable.

The legal specifics of the acquisition, storage, and administration of controlled substances is beyond the scope of this guideline. Variation between and even within nations and jurisdictions is too great to facilitate recommendations specific to all settings. Further, we do not limit the recommendations in this guideline to a particular class, certification, or training level of responder. Whether an amateur recreational backcountry user or trained medical responder, users of these guidelines must employ judgment as to which treatment modalities are within the scope of their training and appropriate to their treatment setting and available resources. In each of the sections to follow, we present the best available evidence identified by the panel as to the safety, efficacy, and austere-specific considerations for available pain treatment modalities.

Methods

An expert panel was convened during the 2013 annual winter meeting of the Wilderness Medical Society (WMS) in Park City, UT, and an original guideline was published in Wilderness and Environmental Medicine in 2014. A subsequent expert panel was convened in 2019 and a partial unpublished revision to the guideline was completed with integration of updated literature sources. To complete this updated literature review, a panel was convened in July of 2022, participation in which was based on the individual’s relevant clinical, research, and/or field experience. The panel included representatives from emergency medicine, anesthesiology, surgery, and the field of prehospital emergency medical services (EMS).

A review of relevant research articles was completed by a keyword search of the MEDLINE and PubMed Central databases via the PubMed search interface in addition to queries of SCOPUS and Google Scholar using keywords targeted to each subject area, with filters applied to remove non-English language sources. Searches were completed during June-August 2022. Keywords searched were generated by section authors. Abstracts were screened by section authors for relevance to austere settings, fundamental relevance to the understanding and treatment of pain, or the pharmacology and pharmacodynamics of medications discussed. Additional sources were identified by reviewing citations from identified sources and review articles. Where evidence directly pertinent to austere settings was not available, the best available evidence was included from EMS, tactical medicine, disaster medicine, emergency medicine, and global health literature sources. Pain control modalities were included based on expert consensus, and any modalities suggested by any member of the panel were subjected to a literature search for relevant evidence and applications.

Articles were excluded for lack of relevance as determined by expert review. Additional relevant review articles obtained for background as well as relevant guidelines on the management of acute pain and traumatic injuries were also selected for inclusion.

Recommendations were developed by section authors and revised by consensus approach. Where evidence relevant to austere environments was limited or of low quality, recommendations were made by the consensus of the group based on the clinical experience of panel members and relevant literature from hospital and pre-hospital medicine.

Recommendations were graded according to the criteria developed by the American College of Chest Physicians (Supplemental Table) (1). Final recommendations included were approved unanimously by all members of the panel.

Terms used in the guideline to discuss pain such as ‘pain’, ‘nociception’, and ‘sensitization’ are used in a manner consistent with the International Association for the Study of Pain terms and definitions (2).

General Approach to Pain Management in an Austere Environment

Musculoskeletal injuries, including strains, sprains, dislocations, and fractures, are the most common cause of pain treated in austere environments. Other circumstances that may require pain management include acute medical ailments and environmental injuries such as cold injury, bites, stings, and burns (3-5).

In all cases, the management of pain is a secondary consideration to the management of immediate threats to life, including compromise of airway, breathing, or circulation, and secondary to scene safety for both the patient and the treating party. After scene safety and a primary survey have been addressed, a secondary survey should be undertaken to evaluate the cause and severity of pain (6).

Traumatic tissue injury, heat, cold, and ischemia can cause pain by different mechanisms, and understanding the mechanism of injury or illness informs appropriate treatment. In an alert patient, a description of mechanism, location, quality, and severity of pain can guide treatment. In an uncooperative or unresponsive patient, evaluation of patient setting and the completion of a thorough secondary survey can identify traumatic as well as environmental injuries on the basis of deformity and tissue appearance (7). Pain control should not be withheld out of concern for masking injury or limiting examination quality, and myths to the contrary have been addressed by quality evidence and expert recommendation (8-11).

If possible, prior to treatment of pain, providers should evaluate pain severity, so as to be able to evaluate response to treatment. A numerical rating scale (e.g. Numerical Rating Scale [NRS-11] or Visual Analog Scale [VAS]) or the patient's subjective description of their pain severity should be used to evaluate response to treatments (12). However, protocols should not compel obligatory use of medications or other interventions in response to specific scores, as a rigid approach requiring treatment in response to a specific threshold has been shown to increase medication-related adverse events (13, 14).

Except for some psychosocial interventions, all pain treatment modalities discussed henceforth are accompanied by both benefits and risks to the patient. In general, centrally acting pharmacologic agents (e.g. opioid medications, benzodiazepines) carry greater risk than peripherally acting agents (non-steroidal medications, local anesthetics) due to their potential to affect hemodynamics and respiration. Parenteral routes of administration (intranasal [IN]/intramuscular [IM]/intravenous [IV]/intraosseous [IO]) carry more risk of adverse effects than enteral or topical routes on account of accelerated absorption and shorter time to peak serum concentration. Importantly, risk and benefit are not directly correlated in all cases, and some low-risk interventions like psychological support or injury immobilization can be substantially effective in the treatment of pain.

While previous WMS guidelines have recommended a stepwise and escalating approach to pain control, we update this approach to consider situations in which the treating clinician recognizes an inherently severe cause of pain and immediately initiates more aggressive treatment measures (15). Accordingly, we recommend an approach to pain treatment that selects and simultaneously applies pain treatment modalities based on the clinician's experience, the cause of pain, and appraisal of reported and anticipated pain severity. When possible, we still

recommend the use of lower-risk interventions before or in concert with higher-risk interventions. Figure 1 provides a conceptual schematic of how different modalities might be selected based on pain severity and available resources.

Figure 1

After application of a given treatment, the patient's pain level should be reevaluated, and the treating provider should determine if pain control is adequate to meet the needs of the patient and the treating provider or team. Given that different modalities of pain control are appropriate for different mechanisms of injury and illness, there can be no clear prescriptive approach or linear road map for the treatment of pain. In some patients, anxiety and psychological distress may significantly amplify a response to a minor injury, while in others, an isolated orthopedic injury may be amenable to a specific intervention (e.g. hematoma block) which is inappropriate in other cases. Experience and judgment, paired with frequent reevaluation of the patient and environment, are critical.

Medical responses in an austere environment are inherently limited, so one must evaluate and select pain treatments to minimize packed weight and bulk. An ideal medication for austere settings is compact, lightweight, durable against crush, abrasion, heat, and cold, nonsedating, has a large therapeutic index, works across multiple routes of administration, and has no potential for abuse or dependence (16-18). Since no such medication exists, providers must weigh the advantages and disadvantages of each medication carried considering the above constraints.

A survey of providers at the WMS summer conference in 2016 demonstrated that among practicing medical provider respondents, 90% carried oral (PO) nonsteroidal anti-inflammatory drugs (NSAIDs), 81% carried acetaminophen (APAP), and 47% carried PO opioids when selecting medications for organized treks (19). These medications, in addition to training in

nonpharmacologic pain management, are likely to suffice in many settings, as minor orthopedic injuries and soft tissue wounds comprise the majority of cases (5). In the management of severe acute pain, however, these medications may be inadequate.

In the following sections we provide further discussion of pain control modalities. In each section we discuss the risks and benefits of these interventions as well as the applications to which they are best suited. Additionally, we discuss treatment considerations such as the need for monitoring for adverse effects. Some of these treatments are advanced and require experience that can only be developed in high-volume clinical environments such as hospitals and EMS services. These treatments provide opportunities for organized rescue teams and advanced medical providers to further refine their treatment of pain, but they are not necessary *per se*, and in some settings may be inappropriate or excessive.

Recommendation: We recommend that providers should manage pain to a level that provides adequate relief, that meets the needs of the treating team, and that recognizes the limitations imposed on the team by the austere environment (Strong recommendation, moderate-quality evidence).

Recommendation: We recommend that, when appropriate to the clinical setting, providers should evaluate patient response to low-risk pain treatments before progressing to higher-risk treatment modalities or routes of administration so as to minimize harm in the austere environment (Strong recommendation, low-quality evidence).

Recommendation: We recommend that, when appropriate to the clinical setting, providers may initiate multiple simultaneous treatments to control pain of greater reported or anticipated severity (Strong recommendation, low-quality evidence).

Nonpharmacologic Pain Management

Psychosocial Interventions

The experience of pain is a product of nociceptive nerve signals from the peripheral nervous system that are interpreted and given emotional valence in the central nervous system.

Practically, this means that the psychological state of the patient can affect their experience of pain (20, 21). Laboratory studies have demonstrated that empathetic statements, caring touch, and the presence of a partner perceived as caring can modulate the experience of pain, reduce pain scores, and affect the autonomic response to pain (22-24).

Anxiety, a threat-detection function mediated by the amygdala, is also a potent mediator of pain. High-anxiety situations induced in the laboratory setting are associated with higher pain scores for a given pain stimulus (25). Additionally, greater acute pain intensity in the emergency department has been associated with elevated anxiety states, and postoperative pain has been correlated with preoperative anxiety levels (26, 27). Indeed, perioperative behavioral interventions have been demonstrated to reduce postoperative pain and pain-related disability (28). Attention can also be manipulated to treat pain; distraction from a noxious stimulus is associated with lower acute pain scores, while attention focused on the source is associated with higher pain scores (20, 25, 29, 30).

As empathy, caring, and distraction can mitigate pain, and conversely, coincident anxiety can increase pain, psychosocial interventions are potentially useful in the treatment of pain. Further, they are well-suited to an austere context, as they are weightless, durable, portable, and cost nothing. The most robust literature targeting the acute reaction to stress and its mitigation comes from work around post-traumatic stress disorder (PTSD) stemming from combat. Treatment modalities targeting reactions to stress in survivors of trauma have been codified into a toolset referred to collectively as Psychological First Aid (PFA) (31). PFA, while developed to treat survivors of combat, has more recently been directed towards wilderness responders and expedition members and has been integrated into wilderness medical texts and curricula (32-34).

The full extent of PFA is beyond the scope of this guideline. However, its principal components provide a framework and shared vocabulary for psychosocial interventions that are likely to aid in the treatment of pain through mitigation of negative emotional states, limitation of stress response, and facilitation of adaptive behaviors (21). These components have been adapted by the authors into a useful mnemonic for the application of PFA in austere environments (Table 1) (35).

Table 1: The ABCDE mnemonic for the psychosocial treatment of pain.

Recommendation: We recommend that providers should train in and apply relevant psychological first aid techniques for treatment of pain in austere environments (Strong recommendation, low-quality evidence).

Splinting, Bandaging, Immobilization, Cryotherapy

Treatment of acute musculoskeletal injuries has traditionally involved variations of the ‘protection, relative rest, ice, compression, and elevation’ (PRICE) treatment protocol. Limited evidence supports this practice. However, PRICE remains in widespread use based on moderate quality evidence and expert opinion (36, 37).

The physiological basis underlying this approach is to reduce edema formation. Extensive edema can be both painful and detrimental to mobilization and tissue healing (38, 39). Currently, the PRICE protocol is considered the best initial treatment regimen for suspected soft-tissue injury and fractures. An alternative strategy called the ‘movement, exercise, analgesics, and treatment/therapy’ (MEAT) protocol has been proposed for managing isolated ligament and tendon injury when fractures have been excluded. The MEAT protocol embraces inflammation, as the inflammatory process promotes healing and quicker return to normal function. One randomized trial showed that early therapeutic exercise resulted in quicker return to normal function vs PRICE therapy (40). If an isolated ligament or tendon injury is highly suspected and fracture is unlikely, MEAT therapy can be substituted. In an austere environment, however, providers may be unable to differentiate between an acute fracture or isolated soft tissue injury, and thus PRICE remains the recommended therapy for most acute musculoskeletal injuries.

The following treatment strategies are recommended for optimal outcomes when utilizing the PRICE protocol:

1. Protection from further injury and providing additional stability with taping, bracing, and/or splinting (41).
2. Relative rest will reduce inflammation and pain. If necessary for evacuation, patients can ambulate.

3. Ice or snow application, when available, will decrease skin temperature below 15°C which is the temperature at which nerve conduction is inhibited and pain decreases (42). Lower tissue temperature also reduces edema formation (39). Cooling should be performed with cycles of 10 minutes of cooling followed by 10 minutes of passive rewarming, whenever practical, for the first 24 to 48 hours post-injury (43). This schedule may not be logistically feasible during an evacuation. The tissues should not be cooled to the degree that might result in frostbite or hypothermia (7, 44). When ice or snow is not available, cold water may be substituted for cooling therapy. Evidence for cryotherapy is limited, so the risks of exposing the extremity and patient to potential frostbite and hypothermia must be weighed against the modest benefits of cryotherapy (37).
4. Compression with an elastic bandage or compression stocking can be used with the intent to reduce swelling. Some evidence exists to support this practice, though only a small number of trials have been published and results are mixed (45). The compression should be form fitting while still allowing adequate muscle expansion and sufficient blood flow. Capillary refill, sensation, and movement of the distal extremities should be checked regularly to monitor for excessive compression.
5. Elevation of the injured area above the level of the heart will reduce potential swelling.

Recommendation: We recommend that applicable elements of the PRICE treatment protocol should be used for acute musculoskeletal injury as first-line pain therapy in austere environments. Injuries in which fracture is thought unlikely should use the MEAT treatment protocol (Strong recommendation, low-quality evidence).

Acetaminophen, Anti-Inflammatories, and NSAIDs.

Acetaminophen (paracetamol)

Acetaminophen (APAP), also known as paracetamol, is a widely used analgesic and antipyretic agent. It acts by inhibiting cyclooxygenase (COX) pathways in the CNS, enzyme pathways required for the production of prostaglandins and prostacyclins (46). Both the analgesic and antipyretic effects of APAP are attributed to reduction in prostaglandin production (47). More recently, research suggests that APAP may also act as a cannabinoid system modulator (48). The analgesic and antipyretic actions of APAP appear to be limited to the CNS, as the medication has not been shown to act on peripheral COX pathways. For this reason, APAP is ineffective as an anti-inflammatory (48).

The most common reported side effect of APAP administration, though rare, is GI discomfort (49-51). The most common severe effects of APAP ingestion are seen with hepatotoxicity from significant overdose. APAP has been associated with hepatocellular damage and death. This is usually seen in high doses, typically a single administration >4 g/day in healthy adults or in chronic use of high doses (though toxicity from a single administration <150 mg/kg for adults or 200 mg/kg for children is rare). Care should be taken in chronic alcoholics and other patients with hepatic dysfunction, where the maximum safe daily dose of APAP may not be known (50, 52, 53).

APAP has the advantage of being lightweight and low bulk. APAP is available in PO formulations (tablets, capsules, syrup, and suspension or solution), as a rectal suppository (PR), and as an IV infusion. Multiple studies have evaluated the efficacy of IV APAP against PO and

rectal (PR) administrations and have found clinical equipoise amongst all formulations (54, 55).

The analgesic activity of all formulations of APAP diminishes over 6 hours (56).

PO administration of APAP is generally the preferred route in most environments if the patient can swallow. The IV route may be utilized with good effect, but no benefit has been shown with this route over others. If the patient cannot take medications PO, PR administration is likely faster, easier, and equally effective. No evidence has suggested teratogenic effects from APAP use in pregnancy (57).

Recommendation: We recommend that APAP, in any formulation or route, should be limited to a maximum daily dose of 60 mg/kg or 4 g in four divided doses (Strong recommendation, moderate-quality evidence).

Recommendation: We recommend that in the austere setting, APAP should be administered PO if possible. If a patient cannot tolerate PO medications, we recommend rectal administration of APAP as second line with suppository (preferred) or PO tablet given rectally (second line) (Strong recommendation, moderate-quality evidence).

Non-Steroidal Anti-Inflammatories

Non-steroidal anti-inflammatory drugs (NSAIDs) are a class of medications which act by inhibiting COX to produce anti-inflammatory as well as antipyretic and analgesic effects (58, 59). They are further classified into salicylates (e.g. aspirin), propionic acids (e.g. naproxen, ibuprofen), acetic acids (e.g. diclofenac, indomethacin), enolic acids (e.g. meloxicam), and selective COX-2 inhibitors (e.g. celecoxib). Except for the selective COX-2 inhibitors, NSAIDS

are non-selective inhibitors of both COX enzymes: COX-1 and COX-2. COX-1 expression is involved in renal function, platelet aggregation, and maintenance of the gastrointestinal (GI) lining. COX-2 expression is generally quiescent until induced during an inflammatory response (60). NSAIDs have been demonstrated to be effective analgesic agents that avoid or reduce the requirement for opiate medications (61, 62).

NSAIDs have numerous potential adverse effects, mainly related to their non-selective inhibition of COX-1 activity. While appropriately dosed short-term use of over-the-counter NSAIDs has been shown to be safe and well tolerated, prolonged use and/or high doses of NSAIDs may lead to gastritis and peptic ulcer disease (63-65). COX inhibition may lead to diminished platelet aggregation that may be compounded by concurrent NSAID-induced gastritis and/or underlying coagulopathies (66). However, multiple studies have shown that no increase in postoperative bleeding occurs with short-term use of NSAIDs (67, 68). Patients with reduced renal function or altered hemodynamics due to acute volume depletion or underlying disease may be at risk for renal injury from renal papillary necrosis leading to further renal dysfunction, electrolyte disturbances, and nephrotic or nephritic syndromes (69). While generally well-tolerated in those with normal renal function, recent data suggest that ibuprofen may have a nephrotoxic effect even in those with no prior renal disease (70).

Studies of multiple NSAIDs have suggested an analgesic ceiling at submaximal doses, e.g. 400 mg PO for ibuprofen (71, 72). The question of whether an additional anti-inflammatory effect is seen at higher doses is unclear. A 2015 Cochrane review could not find evidence of “clinically important differences in analgesic efficacy between NSAIDs and other PO analgesics”, suggesting that any additional anti-inflammatory effect for doses above the

analgesic ceiling has no significant effect on pain, though this is based on low- or very-low quality evidence (73).

Ibuprofen and naproxen are the two most common PO NSAIDs used for analgesia. Aspirin, though a common medication, is usually eschewed for analgesic use, particularly in traumatic injury, due to its anticoagulant properties. Recommended analgesic dosing of ibuprofen is 200-400 mg as needed every 6 hours PO for adults, or 5-10 mg/kg for children. In short term applications, an ibuprofen dose of 1200 mg/day or less has been shown to lead to no further adverse GI effects than placebo in otherwise healthy patients (74). Naproxen is also administered PO and is available in either immediate and extended-release tablets or suspension forms. Recommended analgesic dosing is generally a 550 mg loading dose, followed by 250 mg every 12 hours PO for adults, 5 mg/kg every 12 hours for children (maximum 10 mg/kg/d). Meloxicam is another PO NSAID notable for austere use in that its recommended analgesic dosing for adults is 7.5-15 mg PO as needed only once daily.

Ketorolac is an NSAID that is available PO and in a parenteral form, given IV or IM. Studies show no difference in reduction of pain scores when IM ketorolac is compared with PO ibuprofen, though the IM form may be useful when PO medications cannot be administered (75, 76). Doses for adults are 15 mg IV or 30 mg IM every 6 hours (77).

Topical NSAIDs, though proven to be effective analgesics, have a limited role for treatment of significant injury in the austere environments given the increased weight, bulk, and limited stability of these preparations (78).

Recommendation: We recommend NSAIDs as an effective class of analgesic agents that should be employed as a first-line treatment for mild to severe pain in the austere environment in the absence of contraindications (Strong recommendation, moderate-quality evidence).

Recommendation: We recommend that treatment strategies with NSAIDs should employ the lowest effective dose for the shortest possible duration to minimize adverse effects (Strong recommendation, high-quality evidence).

Recommendation: We recommend that NSAIDs should not be used with known renal dysfunction, pregnancy, history of bariatric surgery, and/or history of GI bleeding (Strong recommendation, high-quality evidence).

Synergistic Effects of Non-Opioid Analgesics

Combinations of NSAIDs with APAP have been demonstrated to provide superior pain control to either drug alone or in combination with a PO opioid. The side effects are fewer and patient satisfaction is higher. These benefits have been seen in a variety of injury patterns and in postoperative patients (79-81). A systematic review of 21 studies over 10 years found a combination of APAP and an NSAID was superior to either drug alone (82).

Military applications of NSAIDs/APAP combinations have been successful (82, 83). US military units give combat-deployed soldiers a ‘pill pack’ to be taken immediately after a penetrating extremity wound of any type. These packs contain APAP, meloxicam, and an antibiotic (84).

Acute traumatic pain can be treated with a combination of APAP and an NSAID effectively, and in many cases, without opioids. For severe pain, the initial doses of APAP and an NSAID should be simultaneous, after which the medications should be administered according to their appropriate dosing intervals. As the foundation of medical therapy for pain due to acute injury, these medications should be regularly administered throughout the acute injury period.

Recommendation: We recommend that a combination of APAP/NSAID should be considered in the treatment of moderate to severe acute pain in the austere environment (Strong recommendation, moderate-quality evidence).

Opioid Analgesia

If psychological first aid, PRICE therapy, and non-opioid therapy are inadequate, or pain is sufficiently severe, an escalation of analgesic agents may be considered. Providing opioid analgesia must be undertaken with care in the austere environment as opioid use can be associated with a wide range of side effects, most concerning respiratory depression and loss of airway protective reflexes. Other opioid side effects include dysphoria, euphoria, pruritus, nausea, vomiting, and constipation. (85-87). Nonetheless, opioids have an established role in the treatment of severe acute pain in the wilderness or austere environment (85, 88, 89). Choice of opioid, route of administration, and dosage are dependent upon many factors, including scope of practice, mechanism and extent of injuries, logistical issues of extraction and transport of the patient, comorbidities, and presence of hypovolemic shock.

Respiratory depression and loss of protective airway reflexes caused by opioids should always be considered when administering opioid analgesia, especially when using fast acting or

potent formulations. With careful titration and administration by experienced physicians and other providers, these problems can usually be avoided (85, 88). Whether opioids are part of the analgesic plan for an expedition, personal use, or wilderness EMS, consideration should be given to management of complications, and a risk/benefit analysis must be performed in advance of carrying and administering opioids. At a minimum, providers administering opioid analgesia should have basic training in airway management as this is the primary means of managing respiratory depression due to opioid use. Naloxone is a competitive opioid receptor antagonist that is useful in temporarily reversing respiratory depression and sedation due to opioids but also carries the risk of precipitating acute opioid withdrawal in chronic users. Carrying naloxone as a rescue medication should be considered when more potent forms of opioid medications are in use.

Recommendation: We suggest that opioid analgesia can be an effective component of acute pain management in austere environments; however, consideration of risk should be weighed against the benefit of these medications (Weak recommendation, moderate-quality evidence).

Recommendation: We recommend that providers who administer opioid analgesics be prepared to recognize and manage respiratory depression. Naloxone availability should be considered when non-oral opioids are in use (Strong recommendation, low-quality evidence).

Oral Opioids

Compared with parenteral preparations, PO formulations of opioids have the benefit of being easier to carry, store, and administer. These medications carry the least concern among opioid

formulations for acute respiratory depression and are generally considered safe for unmonitored use. However, caution should be used in opioid naïve patients, and short-acting opioids should be chosen over long-acting formulations (90). Durability and ease of administration make PO opioids an attractive form of analgesia for the austere environment.

PO opioids are often combined by manufacturers with other analgesics such as APAP to provide synergistic pain relief (91). When using these combined medications, care should be taken to monitor the total dose of both medications. It is preferable to separate the medications when possible and carry opioid and non-opioid medications independently to allow for titration of each medication. Non-opioid PO analgesics have been shown to be equally effective as or non-inferior to PO opioid analgesics and opioid combination medications and they generally have a better side effect profile (92-94). This knowledge should prompt a serious consideration of whether PO opioid analgesia provides any additional benefit over non-opioids in an austere environment. Psychological first aid, PRICE, and non-opioid medications should also be employed where appropriate to reduce the requirement for opioids.

Recommendation: We recommend that PO opioids or opioid/APAP combination drugs are effective for the treatment of acute severe pain in austere settings. However, non-opioid PO analgesic regimens may be preferred for their equivalent efficacy and superior safety profile (Strong recommendation, moderate-quality evidence).

Intranasal, Transmucosal, Sublingual, and Transdermal Opioids

Noninvasive methods of parenteral opioid administration include IN, transmucosal (TM), SL, and transdermal (TD) routes. Fentanyl is the most frequently utilized opioid medication for non-

IV parenteral use given its relatively predictable pharmacokinetics (95, 96). PO fentanyl is generally ineffective due to extensive first pass metabolism. However, IN, TM, and SL opioids are not subject to this metabolism and have the advantage of providing acute pain relief without the requirement for IV access. TM and SL formulations have traditionally been utilized for treatment of breakthrough cancer pain with great success (97, 98).

Intranasal opioid administration also provides a method for acute pain relief without the need for IV access. The IN route provides for rapid absorption of opioids. IN Fentanyl is most widely utilized in EDs, by EMS systems, as well as in wilderness setting such as on-hill ski patrol care (99-103). It has also been found to be safe for pediatric use in multiple settings (104-108). IN opioids are best delivered in a volume of 0.2-0.3 ml into each nostril (109). Volumes greater than 1 ml per nostril are not absorbed nasally but rather swallowed, undergoing first pass metabolism. The drug dose administered should be split between the two nostrils to maximize the absorptive area available and to keep volumes as low as possible. IN medications are most effectively administered through use of an atomizer device (110, 111). The standard 50 ug/ml concentration IV fentanyl is used for IN administration.

When larger doses of opioids are desired IN, the volume of fentanyl required must be given as multiple doses over time to limit volumes per nostril. To overcome this volume limitation and provide greater analgesia in small volume doses, the more potent fentanyl analog sufentanil has been utilized. Sufentanil is nearly 100% bioavailable when given IN and at 20 minutes achieves equivalency with similar IV doses (112). It has been successfully and safely utilized in clinic and ED settings (113, 114).

Oral transmucosal fentanyl citrate (OTFC) in lozenge form delivers fentanyl by TM absorption. OTFC was originally developed for treatment of cancer-related pain. OTFC has been

used extensively by the US military and is safe when utilized in appropriate doses in a healthy military population (115,116). Safe use in the military has also led to safe and effective OTFC use in prehospital environments and recommendations for use in mountain rescue (18, 117).

Sublingual fentanyl administration has been shown to have pharmacokinetics similar to IV fentanyl (118). Recently, sufentanil SL has been approved for treatment of acute pain in opiate naive patients in a monitored setting, developed with support from the United States Department of Defense to provide a non-IV method of battlefield pain control. The SL formulation is safe, effective, and has a decreased side-effect and euphoric profile in comparison to fentanyl (119). Studies of sufentanil SL use have been limited to ED- and hospital-based settings, but pooled studies of 436 and 806 patients suggest that it is safe and efficacious in monitored healthcare settings (120-122). There have not yet been any published studies on prehospital use.

Transdermal opioids are designed for long term maintenance of pain control in opioid tolerant individuals and have limited utility in the wilderness setting due to their slow onset of action. Due to a paucity of evidence for the treatment of acute pain, transdermal opioids are not recommended for use in austere environments.

Recommendation: We recommend OTFC for safe and effective treatment of pain in austere settings (Strong recommendation, moderate-quality evidence).

Recommendation: We recommend intranasal fentanyl for safe and effective pain control in austere environments (Strong recommendation, moderate-quality evidence).

Recommendation: We suggest that despite limited documented prehospital use, sublingual fentanyl and sufentanil are likely safe and effective for austere use provided that patient monitoring is available (Weak recommendation, moderate quality evidence).

Intravenous, Intramuscular, and Intraosseous Opioids

IV opioid administration can provide fast and effective analgesia (85, 89, 123-125). However, obtaining and maintaining IV access may be difficult in austere environments as it requires appropriate skills, suitable equipment, and ongoing maintenance. Environmental factors such as cold or freezing temperatures may also weigh on the decision to pursue IV access.

IV opioids carry an increased licensing and regulatory burden and fall outside the scope of practice for many austere medical providers. The availability of IV medications may be more reasonable within organized wilderness EMS systems or in expedition base camps where prolonged stays and delayed evacuations may occur. Consideration should be given to the potential complications of parenteral opioid use and all providers administering these medications should be proficient at basic airway management (87, 88, 124). Naloxone can be considered as rescue medication.

Intraosseous access (IO) is an alternative route that is more easily established in certain situations and should be considered when IV access is needed and cannot be readily obtained (126). IV and IO routes have several advantages over the PO route: they provide rapid pain relief and reliable drug delivery, are titratable, and may be a familiar mode of administration for healthcare professionals.

Fentanyl has been recommended as the drug of choice for prehospital IV pain control in the EMS setting (88, 125, 127). In austere or prolonged settings, a longer acting opioid might be

preferred. Although the invasive nature of the IV route makes it impractical for many austere environments, once IV/IO access is established, these medications are still considered the gold standard for control of acute, severe pain (88). The IM route avoids the need to establish IV or IO access, but due to variable absorption, onset and efficacy may be inferior. IM fentanyl and morphine have been compared prospectively and proven equivalent in terms of analgesia and side effects when given in equipotent doses (128).

Recommendation: We recommend that parenteral opioids be used to provide analgesia for acute severe pain in austere environments when benefits outweigh risks and when more conservative modalities do not provide sufficient analgesia (Strong recommendation, moderate-quality evidence).

Ketamine

Ketamine is a dissociative anesthetic agent which is an N-methyl-D-aspartate (NMDA) antagonist suspected to block afferent effects of pain perception in the medullary reticular formation, alter certain CNS transmitter systems, and suppress spinal cord activity (129). In common formulation, ketamine is a 50:50 racemic mixture of *s*- and *r*-ketamine. The *s*-ketamine enantiomer is a more potent analgesic agent and in sub-dissociative doses has fewer psychiatric side effects than the racemic mixture, which has led to its growing use as an analgesic in some countries (130). The more widely available racemic form is discussed here.

In lower doses, ketamine has a long history of pre-hospital use for acute pain. Ketamine can be administered via several routes including IM, IV/IO, IN, PO, SL, and PR (131). Most wilderness applications involve IM, IN, and IV/IO routes.

Ketamine is unique among pain control medications in that it has variable effects at different doses, and care must be taken to remain within the range of ‘pain dose’ or ‘low dose’ ketamine (0.1-0.5 mg/kg IV or 1-2 mg/kg IM) (131,132). At higher doses (1-2 mg/kg IV or 3-5 mg/kg IM), ketamine causes dissociation, a type of moderate sedation often employed in hospital use (133). Between these dosing ranges, ketamine can cause hallucinations, euphoria/dysphoria, delirium, and other undesirable effects.

For pain control, ketamine is administered within the low dose range as above by slow IV push, IV infusion, or IM injection. If repeat dosing is undertaken, care must be taken to avoid rapid redosing and induction of hallucinations or sedation. Ketamine can be safely used in combination with opioids to enhance pain control if multiple modalities are needed (134). Ketamine is well tolerated and has been shown to be safe when administered in dissociative doses by non-physician practitioners to facilitate procedures, which reinforces its safety when used in pain control doses (135).

Ketamine has been reported to be safe, useful, and to have few associated adverse events when used by physicians working in an air medical mountain rescue service (136). It has also been safely used at dissociative doses for procedures at high altitude (>3900m) by primary care physicians without anesthesia training. This application in 11 cases found no significant complications or hypoxia associated with ketamine use (137).

In appropriately titrated pain doses, patients can maintain their own airway as pharyngeal reflexes and spontaneous ventilation are usually preserved. However, salivation is often increased, and airway obstruction has been reported with dissociative doses (138). Unlike many anesthetic and analgesic agents, cardiovascular function is often well preserved in the setting of

ketamine administration. This makes ketamine a suitable choice for trauma patients if hemorrhagic shock is suspected (139).

Recommendation: We recommend ketamine for the control of acute severe pain in the austere setting using pain-control doses (0.1-0.5mg/kg IV, 1-2mg/kg IM). Providers should be experienced in using the medication and equipped to manage airway complications should they arise (Strong recommendation, moderate-quality evidence).

Local Anesthetics

Local anesthetic (LA) agents act by prolonging deactivation of voltage-gated sodium channels. When acting on peripheral neurons, they reduce or reversibly block neuronal transmission, leading to local analgesia or anesthesia. However, when absorbed systemically, these medications can cause severe adverse effects by acting on sodium channels in the myocardium or CNS, such as cardiac arrest and seizure. Attention must thus be given to weight-based maximum doses to avoid toxicity (e.g. lidocaine 4-5 mg/kg, maximum 300 mg/dose and 2400 mg/day), and care must be taken to avoid intravascular or intraneural injection.

Worldwide, the most used LAs are lidocaine (lignocaine) and bupivacaine, although there are many variations and preparations within the category. The onset, duration of action, and density of the local analgesia or anesthesia produced by LAs depend upon multiple medication factors including the lipid solubility, dissociation constant, concentration, and total dose. Typically, agents with more-rapid onset tend to have shorter duration of action, whereas longer-acting drugs carry higher risks of toxicity (140).

LAs can be applied topically to the skin or mucous membranes, injected directly into peripheral tissues, or adjacent to regional nerves or even the spinal cord to produce regional or neuraxial analgesia or anesthesia.

Topical Local Anesthesia

Topical anesthetic agents are directly applied on or around peripheral sites of nociception in gel, liquid, ointment, or cream form, blocking noxious stimuli at their source (141-144). Administration is simple, and the low systemic absorption reduces the risk of side effects (145).

There are many available formulations of anesthetics for topical use including combination products and viscous formulations to facilitate easy application (e.g. lidocaine/epinephrine/tetracaine (LET) and the eutectic mixture of lidocaine/prilocaine (EMLA). The efficacy and safety of these formulations has been demonstrated for treatment of wounds and stings and for the facilitation of laceration repairs (146, 147). The efficacy of these formulations outside of recommended storage temperatures of 0-30° C has not been studied and may limit their use in some austere environments.

Recommendation: We recommend locally instilled and topically applied local anesthetic agents for safe and effective treatment of pain related to burns, stings, and soft tissue injuries in austere environments (Strong recommendation, moderate-quality evidence).

Ophthalmic/Otic Solutions

Ophthalmic and otic solutions of topical anesthetics are useful in the treatment of conditions including corneal abrasions or ulcerations, ultraviolet keratitis, otitis media, and otitis externa

(148, 149). These preparations differ from plain aqueous solutions of LAs in that they often contain preservatives, buffers, viscosity agents, and other additives to make them more effective and tolerable at the target site (150). Lacking alternatives, an aqueous solution of lidocaine, as used for local infiltration, can be used in topical ocular application. However, the lack of pH buffering reduces penetration into the tear film and causes significant discomfort on instillation, reducing efficacy and tolerability (151). Ophthalmic and otic solutions are generally similar in composition, though ophthalmic solutions are sterile and free of particulate matter, so ophthalmic solutions may be instilled in the ear, but otic solutions should not be used in the eye. Otic solutions are generally less expensive (152).

Ophthalmic solutions can be of great use in the care of a patient who has become debilitated by a corneal abrasion or keratitis, as analgesia can facilitate eye opening and allow the patient to participate in their evacuation. Concerns over delayed corneal healing have led many providers to withhold repeated use in the setting of corneal injury (153). However, a meta-analysis pooling two randomized, prospective studies in the emergency department showed no clinically significant adverse effects in the repeated use of ophthalmic proparacaine for less than 72 hours (154). Accordingly, it is reasonable to use a short course of ophthalmic anesthetics to aid in the evacuation of otherwise debilitated patients. Due to reduced capacity for self-monitoring, care should be taken to protect an anesthetized eye from insults, including dryness, exposure to foreign bodies, or to UV radiation (155).

Recommendation: We suggest that ophthalmic anesthetic solutions are, when used appropriately, safe, and useful for the treatment of acute ocular and auricular pain in austere environments (Strong recommendation, low-quality evidence).

651

652 *Regional Analgesia*

653 Regional analgesia employs LA injection near and about nerves or nerve plexi to achieve broad
654 areas of analgesia and anesthesia. The use of regional analgesia/anesthesia in the austere
655 environment has the potential to meet many goals of austere pain management, such as avoiding
656 sedation, respiratory depression, cardiovascular instability and other unwanted side-effects
657 (156,157). Regional anesthesia also uses a limited amount of compact equipment to provide
658 rapid and predictable onset of excellent pain control. However, such approaches require
659 additional equipment, pharmacological agents, training, and experience. Selection of patient,
660 setting, and technique must thus be carefully considered (157). Neuraxial (intrathecal/spinal and
661 epidural) injections are unlikely to be suited to austere conditions outside of specific
662 circumstances, and thus fall beyond the scope of these guidelines.

663

664 *Field-expedient regional blocks*

665 In contrast to neuraxial techniques, regional anesthesia and analgesia can be achieved in many
666 settings with minimal equipment and limited training while still retaining an acceptable margin
667 of safety. Well-executed nerve blocks can provide excellent and long-lasting analgesia, or full
668 anesthesia of a limb or region to allow interventions and even mobilization and self-rescue.
669 However, these advantages must be balanced with safety; nerve blocks may also hamper self-
670 rescue or render a limb unusable for hours.

671 The concept of field-expedient regional blocks (FERBs) refers to nerve block techniques
672 which:

- 673 • Have consistent approaches using reliable anatomical landmarks;

- Can be performed reliably with a predictable distribution of effect;
- Avoid structures which carry high risk or consequences of injury (e.g. pleura or major vessels);
- Have a low risk of unwanted effects (e.g. phrenic nerve block and diaphragmatic paralysis which frequently accompanies interscalene brachial plexus block);
- Require drug volumes which are well within the maximum safe dose of the selected LA;
- Tend towards slower drug absorption, reducing the risk of systemic toxicity;
- Are easily learned and can be practiced by austere medicine practitioners working in a variety of settings to become proficient before using them in the field;
- Do not require highly specialized equipment (158).

While a full discussion of suitable FERBs is beyond the scope of these guidelines, examples of the concept include certain orbital, dental and occipital blocks in the head and neck, coracoid, wrist, and interdigital blocks in the upper limb, the fascia iliaca plane block, popliteal, peroneal, ankle, and intermetatarsal blocks in the lower limb, and ring blockade of any digit (159-164).

Ultrasound has become the standard of care in-hospital for many regional blocks and is mandatory for some techniques. It allows visualization of tissue planes, nerves, blood vessels and other structures with minimal discomfort, and has many other uses in austere medicine (156). However, ultrasound requires additional training and practice to be used effectively for regional blocks, which should be obtained in a controlled setting.

Practitioners should know the potential risks and complications of regional nerve blocks and seek patient consent wherever possible before performing a block. Knowledge of the signs, symptoms, and management of local anesthetic systemic toxicity (LAST) is essential. IV access

should be obtained before all but the most simple blocks to facilitate treatment of complications as needed, and equipment and drugs for resuscitation should always be available before commencing a regional block (165). These techniques are therefore limited to well-equipped and trained rescue teams and responders.

Recommendation: We suggest that when performed by experienced practitioners with necessary administration and monitoring equipment, field-expedient regional nerve blocks may be an effective tool to manage pain and facilitate procedures in an austere environment. (Weak recommendation, low-quality evidence).

Recommendation: We recommend that portable ultrasound be used for FERBs in austere environments when appropriate equipment and skills are available (Strong recommendation, moderate quality evidence).

Care of patients with nerve blocks in the field

Local and regional analgesia may facilitate (but should not delay) life- and limb-saving interventions. Well-executed regional analgesia may improve a patient's ability to communicate or partake in self-care and self-rescue activities, but it does not negate the responsibility to perform ongoing monitoring and assessment. Care should be given to assessing for symptoms and signs of LAST and to the care of the region under analgesia/anesthesia to prevent further injury. Patients with well-executed blocks may not be able to feel pressure points, control an anesthetized limb, or perceive inadequate perfusion from compartment syndrome, dressings, or immobilization measures which are too restrictive.

720

721 *Recommendations: We recommend immobilization with adequate padding, pressure care, and*
722 *frequent assessments for perfusion and absence of clinical signs of compartment syndrome as*
723 *necessary for patients who have received regional blocks in austere environments (Strong*
724 *recommendation, low-quality evidence).*

725

726 *Hematoma blocks*

727 In contrast to all other modern regional analgesia techniques, a hematoma block aims to
728 deliberately aspirate blood by advancing a 20G needle directly into the hematoma surrounding a
729 fracture using sterile technique. It is nearly exclusively used for reduction of distal radius/ulna
730 fractures, although its use has also been described for fractures of the tibia (166, 167). As
731 fracture hematomas can be considered contiguous with the bone marrow space, absorption of
732 LAs via hematoma could theoretically be as rapid as IO drug administration (168). For this
733 reason, only lidocaine is used for this block, at a maximum dose of 1.5-2 mg/kg. For most adult
734 patients, a volume of 5-10 ml of 1% lidocaine solution is safe and effective (169, 170).

735

736 *Recommendations: We suggest that hematoma blocks with plain lidocaine can be performed by*
737 *suitably experienced practitioners in austere environments (Weak recommendation, moderate*
738 *quality evidence).*

739

740 *Field infiltration of local anesthetics*

741 Although not a true regional block, infiltration of LAs around the margins of a wound or in an
742 injured region is an effective means of providing analgesia and can extend to procedural

locoregional anesthesia. In this setting, there is little advantage to using concentrated LA, and dilutions of 0.5 to 1% lidocaine and as little as 0.1% of long-acting agents can be effective. Use of a narrow-gauge needle and injection with smooth continuous withdrawal reduces the risk of intravascular injection. Caution should still be employed in calculating the maximum safe dose; particularly when providing repeated injections within a limited timeframe.

Recommendations: We recommend field infiltration of dilute LA for safe management of acute pain due to soft tissue wounds in austere environments (Strong recommendation, moderate-quality evidence).

Infiltration of diphenhydramine

Diphenhydramine (DPH) is an antihistamine that is an inverse agonist at the H1 receptor. It is usually employed in the treatment of allergy or anaphylaxis and may be carried as a component of austere medical kits for this reason. While PO formulations of DPH are limited to these traditional indications, a 1% aqueous DPH formulation diluted from the 5% typical for IV use is also useful as a LA, particularly in those patients with allergies to typical LAs such as lidocaine. Local injection of 1% DPH achieves adequate analgesia for wound care within 5 minutes in 80% of patients (171-173). In comparison to lidocaine, the injection of DPH is more painful, the analgesic effect slightly less than that of lidocaine but adequate for suturing and wound care, and DPH analgesia has shorter duration than that of lidocaine (174, 175). DHP infiltration can cause dose-dependent mild sedation, and this should be taken into account if the patient is expected to ambulate or participate in self-rescue (171). The potential adverse effects of intraarterial injection DPH are minimal, particularly in comparison to lidocaine. Local DPH infiltration can,

rarely, cause skin irritation (171, 175). To date, DPH has not been studied for use in nerve, regional, and hematoma blocks.

Recommendation: We recommend aqueous DPH solution as safe and effective for local infiltration and wound care as an alternative to lidocaine in the austere environment when lidocaine is not available, or allergy limits its use. Consideration should be given to its sedating effects (Strong recommendation, moderate-quality evidence).

Intravenous Lidocaine

IV lidocaine use as an infusion for the treatment of pain has been widely reported in hospital-based applications and has shown equivalent effect relative to opioids in some applications, as well as opioid sparing effects (176, 177). However, anesthesia guidelines describe the use of IV lidocaine as “high risk” even in the hospital setting due to its narrow therapeutic window (178). In hospital application, access to lipid emulsion therapy is considered necessary to treat possible overdose. In a systematic review of its use in hospital application, of 289 patients treated with IV lidocaine, 44 adverse events were reported of which 36 (12.4%) were ‘serious’, including altered mental status and cardiac dysrhythmias (177). Meta-analysis of its use in the emergency department yielded similar adverse event rates but showed equivocal performance in comparison to opioids (176)

A case report is available describing two patients treated with IV lidocaine in an austere setting. Both patients experienced improved pain, however, both patients were treated without the availability of lipid emulsion and both experienced prodromal symptoms of LAST (179). This highlights the hazard of employing this method without hospital-level resources. Because of

the degree of monitoring and resuscitation equipment required, and given the very poor therapeutic index of this method, IV use of LAs is not recommended.

Recommendation: We recommend that IV lidocaine should not be used in the austere environment (Strong recommendation, low-quality evidence).

Inhalational Analgesics

Inhalational analgesia is provided by breathing specific gases or volatile agents which induce analgesic effects after absorption in the lung. This should be distinguished from topical/TM administration of agents using atomization or nebulization.

Inhalational analgesia is characterized by rapid on- and offset, ease of administration, lack of immediate requirement of IV access, and comparable efficacy to other systemic agents. Currently, there are only two inhalational analgesics in clinical use: nitrous oxide (a gas), and methoxyflurane (a vapor). Both require specific equipment and methods to administer and are discussed further below.

Nitrous Oxide

Nitrous oxide (N₂O) is a non-flammable, colorless gas with a slightly sweet scent which has euphoric and dissociative anesthetic properties at high partial pressures, and significant analgesia at lower doses. N₂O is typically self-administered by breathing through a demand-valve and mouthpiece that is held between the teeth, or directly through an anesthetic face mask. Onset of action is rapid, within several breaths; clinically significant analgesia occurs within 5 minutes (180). While the precise mechanism of action is not fully understood, anesthetic effects are likely

812 through non-competitive central NMDA-receptor antagonism, while analgesic effects are
813 mediated by endogenous opioid release in the midbrain (181). It is commonly used as a
814 synergistic agent during induction and maintenance of general anesthesia but has also been
815 extensively administered in combination with oxygen (O₂) as an analgesic for painful conditions
816 and procedures such as labor and delivery or dental work. Outside of the operating theater, it is
817 most commonly available in an equal, preblended (50%) mixture with oxygen, known as
818 Entonox. N₂O can also be mixed with a separate supply of pure O₂ using a blender, though this is
819 more cumbersome for prehospital use. N₂O has been used with considerable success as an
820 analgesic in pre-hospital care for many years (180, 182).

821 Patient-administered N₂O analgesia has been shown to have a low rate of adverse events
822 and sedation, in both children and adults (183, 184). The most-common side effects, such as
823 nausea, dizziness, and mild sedation, are normally short lived. At analgesic doses, it exhibits
824 cardiovascular stability and no respiratory depression and can provide analgesia on par with
825 systemic opiate administration (181, 185-187). The rapid diffusion of N₂O can lead to
826 accumulation in gas-filled spaces within the body, and it should be used cautiously in head and
827 chest trauma victims so as not to worsen pneumothorax or pneumocephalus, and its use is
828 contraindicated in recently-ascended divers (188).

829 Three important limitations for the use of N₂O in the austere setting exist. First, it is
830 stored in bulky, heavy, pressurized cylinders (e.g. Entonox), which require the use of regulators
831 and a demand valve, making transport and storage challenging.

832 Second, the nature of the pressurized gas mixture reduces the range of ambient
833 temperatures in which it can be used. Direct sunlight and hot temperatures can risk cylinder
834 rupture due to overpressure, and subzero (below -5.5°C) temperatures will cause the N₂O to

liquify. Initially, this can result in the inhalation of oxygen with limited N₂O in unpredictable concentrations. However, as the cylinder empties, an increasing concentration of N₂O is delivered, and the mixture becomes dangerously hypoxic (213). Accordingly, N₂O/O₂ mixtures must be administered with care at below-freezing temperatures.

Third, N₂O is delivered as a gas, and increasing altitude will cause a corresponding decrease in the inhaled partial pressure of gases. This in turn decreases the analgesic effect of N₂O at altitude (189). Thus, while it is a useful agent at sea level, the value of N₂O is greatly decreased at high altitude (190-192). Additionally, N₂O can accumulate in small and poorly ventilated spaces, so adequate ventilation should be assured to prevent unintentional inhalation among bystanders.

These restrictions on the portability of the agent and use under adverse conditions has discouraged austere use. Although N₂O has been employed among ski patrols in Canada, Australia, and the United States, its use on the ski hill, including efficacy and safety, has not been reported (193).

Recommendation: We suggest N₂O as a safe and effective analgesic in austere environments for short painful procedures or for limited periods of time. Although evidence pertinent to austere use is limited, it can be safely used at low altitude, with appropriate monitoring, and where appropriate storage of cylinders can be assured (Weak recommendation, low-quality evidence).

Methoxyflurane

Methoxyflurane is a halogenated volatile anesthetic with analgesic properties (194). Its mechanism of action is not well understood, but it is a known GABA receptor agonist and

reduces gap junction function in the brain. Although it is no longer used for anesthesia, it has been widely used as an analgesic in Australia and New Zealand by paramedic, military and civilian first aid providers for over 30 years (195).

Methoxyflurane is self-administered by inhalation through a hand-held whistle-like device which functions as a simple, disposable draw-over vaporizer. Onset of analgesia is rapid (within 5 minutes) and continues while the patient breathes through the device (196). When used continuously, a single dose (3 ml) lasts for approximately 25-30 minutes (197). Its effects are rapidly reversed once inhalation stops.

Changes in cardiovascular, respiratory and neurological function during methoxyflurane administration are not clinically significant and therefore IV access is not essential (196, 198). The most common side effects are dizziness, headache and feeling somnolent (197, 199). Serious adverse effects are rare, but methoxyflurane is contraindicated in patients with renal or hepatic impairment as well as those who have a personal or family history of malignant hyperthermia (101, 199).

Concerns have been raised as to whether methoxyflurane will suffer the same decrease in efficacy at altitude which is experienced with N_2O (200). However, as saturated vapor pressure is not influenced by ambient pressure, altitude should not affect the partial pressure delivered by the device, a theory which has been borne out in initial laboratory work (189, 201). Questions which remain to be studied are the effect of low ambient temperatures on device performance and whether environmental factors influence clinical efficacy of methoxyflurane in the field. Efficacy for analgo-sedation at high altitude (4470 m) appears to be preserved in one case study (202).

Methoxyflurane has been shown to have more rapid onset and comparable or better initial analgesia compared to IV opiates or APAP (203-206). In addition, its portability, stability, and ease of administration make it highly suitable for use in austere settings (101, 207, 208). The rapid onset of action and limited side effect profile make methoxyflurane useful as a temporizing measure until other modalities can be established or as a definitive short-term analgesic (195, 209).

Recommendation: We recommend methoxyflurane as safe and effective to treat pain in sub-anesthetic doses in austere environments (Strong recommendation, moderate-quality evidence).

Adjuncts

Benzodiazepines

Benzodiazepines (BZDs), which have long been used in the treatment of anxiety, have previously been recommended as a possible analgesic adjunct (15). This was rational because, as previously discussed, there is a correlation between anxiety states and increased perception of acute pain. However, no quality evidence has emerged to support the administration of BZDs for analgesia.

BZDs act centrally on GABA-A receptors, resulting in CNS depression (210). Additionally, the combination of BZDs with other CNS depressants can potentiate their concerning adverse effects. The combination of BZDs with opioids has been well-studied and has the potential to induce significant respiratory depression (211-213).

Recommendation: We recommend that BZDs should not be used for the treatment of pain in austere settings (Strong recommendation, high-quality evidence).

Further Modalities and Adjuncts

A variety of additional medication classes and techniques have been studied as adjuncts for pain in both the hospital and outpatient environment. These include antihistamines, battlefield acupuncture, antipsychotics, non-BZD muscle relaxants, capsaicin, and antidepressant medications. While there is heterogeneous but generally supportive evidence for the use of these medications in the long-term treatment of chronic pain, there is inadequate evidence to support their use for the treatment of acute pain in the austere environment, and austere use cannot be recommended.

Conclusion

In all cases, the comfort of the patient must be balanced with the risk of any tools used to address pain. In comparison to hospital-based or EMS medicine, evidence for pain control modalities specific to the austere environment is generally limited. Accordingly, the experience of the responder, consideration for available resources, and clinical judgment are of paramount importance in the selection of pain control modalities appropriate to the patient and the setting. Some interventions may enable the patient to participate in their own rescue while others may generate unacceptable risk to the patient or team. Backcountry recreationalists, expedition leaders, and professional rescuers alike should consider the resources, hazards, and limitations specific to their unique settings and should both plan for appropriate modalities to manage acute

925 pain, and for the training, monitors, equipment, and antidotes needed to manage the risks of
926 those interventions. The paucity of austere-specific research also opens numerous opportunities
927 for further study.

928 **Acknowledgements:** The authors would like to acknowledge the contributions of Katarzyna A.
929 Grela, MBBS, FANZCA, Katie W. Russell, MD; Courtney L. Scaife, MD, and Rom A. Stevens,
930 MD to the unpublished 2019 revision of this clinical practice guideline, which informed the
931 current text.

932

933 **Author Contributions:**

934 Acquisition of data: (PF, AW, WS, GBZ, JL, SM, CVT, IW, JW, RH, DW)

935 Analysis of the data: (PF, AW, WS, GBZ, JL, SM, CVT, IW, JW, RH, DW)

936 Drafting of the manuscript: (PF, AW, WS, GBZ, JL, SM, CVT, IW, JW, RH, DW)

937 Critical revision of the manuscript: (PF, AW, WS, GBZ, JL, SM, CVT, IW, JW, RH, DW)

938 Approval of the final manuscript: All authors approved the final manuscript.

939

940 **Financial/Material Support Statement:** None

941

942 **Disclosures:** None of the authors have any conflicts of interest to disclose. The opinions of
943 the authors expressed in this manuscript are their own, and do not reflect official policy of the
944 United States Army or the Department of Defense

945

946

947 *Table 1: The ABCDE mnemonic for the psychosocial treatment of pain.*

A	Anchoring, Attention & Acknowledgement	Provide the patient with an attentive provider who identifies themselves as responsible for their comfort and who acknowledges their distress.
B	Breathing	Encourage controlled breathing depth and rate, limiting rapid, shallow breathing or hyperventilation.
C	Control and Cognitive Shift	Provide the patient with a role that allows an element of control over their situation, and which shifts their mindset away from catastrophizing.
D	Decrease nociception, Distract, Diffuse.	Use other pharmacological/nonpharmacological interventions to treat nociceptive signals. Distract the patient from the source of pain. Diffuse tension using empathy and humor.

E	Explanations and Expectations	Provide the patient with an understanding of what to expect in novel situations, such as technical rescue or vehicle/helicopter transport. Explain procedures to be performed, personnel involved, timelines to be expected, etc.
---	-------------------------------	---

948

949 **Figure Legends**

950 Figure 1: Conceptual schematic of escalating pain treatment stratified by risk posed to the patient
951 and the need for monitoring when used. With greater anticipated or reported pain severity, a
952 clinician may elect to use one or more modalities appropriate to that degree of severity. IV,
953 intravenous; IM, intramuscular; IN, intranasal; NSAID, non-steroidal anti-inflammatory drug.

954

955 Table 1: The ABCDE mnemonic for the psychosocial treatment of pain: Suggested mnemonic
956 for the field application of psychological first aid to reduce acute pain.

References

1. Diekemper R, Patel S, Mette S, Ornelas J, Ouellette D, Casey K. Making the GRADE: CHEST Updates Its Methodology. *CHEST*. 2018;153(3):756-9.
2. Group ITW. Pain Terms and Definitions. International Association for the Study of Pain. <https://www.iasp-pain.org/resources/terminology/#pain>. Published 2011. Accessed 07/22/2022, 2022.
3. Leemon D, Schimelpfenig T. Wilderness Injury, Illness, and Evacuation: National Outdoor Leadership School's Incident Profiles, 1999–2002. *Wilderness Environ Med*. 2003;14:174-82.
4. McIntosh SE, Leemon D, Visitacion J, Schimelpfenig T, Fosnocht D. Medical Incidents and Evacuations on Wilderness Expeditions. *Wilderness Environ Med*. 2007;18:298-304.
5. Wells FC, Warden CR. Medical Incidents and Evacuations on Wilderness Expeditions for the Northwest Outward Bound School. *Wilderness Environ Med*. 2018;29(4):479-87.
6. *Advanced Trauma Life Support® Student Course Manual*. 10th ed: American College of Surgeons; 2018.
7. McIntosh SE, Freer L, Grissom CK, Auerbach PS, Rodway GW, Cochran A, et al. Wilderness Medical Society Clinical Practice Guidelines for the Prevention and Treatment of Frostbite: 2019 Update. *Wilderness Environ Med*. 2019;30(4S):S19-S32.
8. King K, Brewster G, Herbert M, Hoffman J. Medical Myth: Analgesia should not be given to patients with an acute abdomen because it obscures the diagnosis. *Western Journal of Medicine*. 2000;172(3):209-10.
9. Silen W. *Cope's Early Diagnosis of the Acute Abdomen*. 22nd Ed. ed: Oxford University Press; 2010.

- 980 10. Bernard A, Oyler D, Anglen J, Barrett M, Bhalla P, Cocanour C, et al. ACS Trauma
981 Quality Programs Best Practices Guidelines for Acute Pain Management in Trauma
982 Patients. 2020.
- 983 11. Hsu JR, Mir H, Wally MK, Seymour RB, Orthopaedic Trauma Association
984 Musculoskeletal Pain Task F. Clinical Practice Guidelines for Pain Management in Acute
985 Musculoskeletal Injury. *J Orthop Trauma*. 2019;33(5):e158-e82.
- 986 12. Hjerstad MJ, Fayers PM, Haugen DF, Caraceni A, Hanks GW, Loge JH, et al. Studies
987 comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales
988 for assessment of pain intensity in adults: a systematic literature review. *J Pain Symptom*
989 *Manage*. 2011;41(6):1073-93.
- 990 13. Vila H, Jr., Smith RA, Augustyniak MJ, Nagi PA, Soto RG, Ross TW, et al. The efficacy
991 and safety of pain management before and after implementation of hospital-wide pain
992 management standards: is patient safety compromised by treatment based solely on
993 numerical pain ratings? *Anesth Analg*. 2005;101(2):474-80.
- 994 14. Mularski RA, White-Chu F, Overbay D, Miller L, Asch SM, Ganzini L. Measuring pain
995 as the 5th vital sign does not improve quality of pain management. *J Gen Intern Med*.
996 2006;21(6):607-12.
- 997 15. Russell KW, Scaife CL, Weber DC, Windsor JS, Wheeler AR, Smith WR, et al.
998 Wilderness Medical Society practice guidelines for the treatment of acute pain in remote
999 environments: 2014 update. *Wilderness Environ Med*. 2014;25(4 Suppl):S96-104.
- 1000 16. Surrett G, Franklin J, Wedmore I. Pain Control In Austere Settings. *Current Sports*
1001 *Medicine Reports*. 2015;14(2):117-22.

- 1002 17. Wedmore IS, Johnson T, Czarnik J, Hendrix S. Pain management in the wilderness and
1003 operational setting. *Emerg Med Clin North Am.* 2005;23(2):585-601, xi-xii.
- 1004 18. Ellerton J, Milani M, Blancher M, Zen-Ruffinen G, Skaiaa SC, Brink B, et al. Managing
1005 moderate and severe pain in mountain rescue. *High Alt Med Biol.* 2014;15(1):8-14.
- 1006 19. Schauer SG, Naylor JF, Brown DJ, Gibbons RV, Syndergaard I, Cushing T. A Survey of
1007 Wilderness Medicine Analgesia Practice Patterns. *Wilderness Environ Med.*
1008 2018;29(2):211-4.
- 1009 20. Hansen G, Streltzer J. The psychology of pain. *Emerg Med Clinics.* 2005;23(2):339-48.
- 1010 21. Linton S, WS S. Impact of psychological factors in the experience of pain. *Phys Ther.*
1011 2011;91:700-11.
- 1012 22. Sambo CF, Howard M, Kopelman M, Williams S, Fotopoulou A. Knowing you care:
1013 effects of perceived empathy and attachment style on pain perception. *Pain.*
1014 2010;151(3):687-93.
- 1015 23. Fauchon C, Faillenot I, Perrin AM, Borg C, Pichot V, Chouchou F, et al. Does an
1016 observer's empathy influence my pain? Effect of perceived empathetic or unempathetic
1017 support on a pain test. *Eur J Neurosci.* 2017;46(10):2629-37.
- 1018 24. von Mohr M, Krahe C, Beck B, Fotopoulou A. The social buffering of pain by affective
1019 touch: a laser-evoked potential study in romantic couples. *Soc Cogn Affect Neurosci.*
1020 2018;13(11):1121-30.
- 1021 25. James J, Hardardottir D. Influence of attention focus and trait anxiety on tolerance of
1022 acute pain. *British Journal of Health Psychology.* 2002;7:149-62.

- 1023 26. Kapoor S, White J, Thorn BE, Block P. Patients Presenting to the Emergency Department
1024 with Acute Pain: The Significant Role of Pain Catastrophizing and State Anxiety. *Pain*
1025 *Med.* 2016;17(6):1069-78.
- 1026 27. Woo A. Depression and Anxiety in Pain. *Reviews in Pain.* 2010;4(1).
- 1027 28. Nicholls JL, Azam MA, Burns LC, Englesakis M, Sutherland AM, Weinrib AZ, et al.
1028 Psychological treatments for the management of postsurgical pain: a systematic review of
1029 randomized controlled trials. *Patient Relat Outcome Meas.* 2018;9:49-64.
- 1030 29. Edwards R. Psychological Treatment in Acute Pain. In: Edwards R, ed. *Encyclopedia of*
1031 *Pain.* Vol 2007.2007:2030-2.
- 1032 30. Rischer KM, Gonzalez-Roldan AM, Montoya P, Gigl S, Anton F, van der Meulen M.
1033 Distraction from pain: The role of selective attention and pain catastrophizing. *Eur J*
1034 *Pain.* 2020;24(10):1880-91.
- 1035 31. Psychological First Aid Field Operations Guide. 2006.
1036 https://www.ptsd.va.gov/professional/treat/type/psych_firstaid_manual.asp. Published
1037 2nd. Accessed 8/3/2022.
- 1038 32. Mcgladrey L. Psychological First Aid and Stress Injuries. In: Hawkins S, ed. *Wilderness*
1039 *EMS.* Illustrated ed.: Wolters Kluwer Health; 2017:292-308.
- 1040 33. Schimelpfenig T. Stress and the Rescuer. In: *NOLS Wilderness Medicine.* 7th ed.:
1041 Stackpole Books; 2021:300-5.
- 1042 34. Mortimer A, Mortimer R. Psychological First Aid for Wilderness Trauma: Interventions
1043 for Expedition or Search and Rescue Team Members. *Wilderness Environ Med.*
1044 2023;34(3):346-53.

- 1045 35. Hofmeyr R. Wilderness Pain Management: State of the Evidence and State of the Art.
1046 Wilderness Medicine Society Summer Conference; August 3, 2022, 2022; Snowmass,
1047 CO, modified with permission from Hofmeyr R, Duys RA. Managing pain in an
1048 emergency. PainSA 2019, Drakensberg, South Africa, 2019.
- 1049 36. van den Bekerom MP, Struijs PA, Blankevoort L, Welling L, van Dijk CN, Kerkhoffs
1050 GM. What is the evidence for rest, ice, compression, and elevation therapy in the
1051 treatment of ankle sprains in adults? *J Athl Train*. 2012;47(4):435-43.
- 1052 37. Miranda JP, Silva WT, Silva HJ, Mascarenhas RO, Oliveira VC. Effectiveness of
1053 cryotherapy on pain intensity, swelling, range of motion, function and recurrence in acute
1054 ankle sprain: A systematic review of randomized controlled trials. *Phys Ther Sport*.
1055 2021;49:243-9.
- 1056 38. Bleakley CM, O'Connor S, Tully MA, Rocke LG, Macauley DC, McDonough SM. The
1057 PRICE study (Protection Rest Ice Compression Elevation): design of a randomised
1058 controlled trial comparing standard versus cryokinetic ice applications in the management
1059 of acute ankle sprain [ISRCTN13903946]. *BMC Musculoskelet Disord*. 2007;8:125.
- 1060 39. Malanga G, Yan N, Stark J. Mechanisms and efficacy of heat and cold therapies for
1061 musculoskeletal injury. *Postgrad Med*. 2015;127(1):57-65.
- 1062 40. Bleakley CM, O'Connor SR, Tully MA, Rocke LG, Macauley DC, Bradbury I, et al.
1063 Effect of accelerated rehabilitation on function after ankle sprain: randomised controlled
1064 trial. *BMJ*. 2010;340:c1964.
- 1065 41. Sprouse R, McLaughlin A, Harris G. Braces and Splints for Common Musculoskeletal
1066 Conditions. *Am Fam Physician*. 2018;98(10):570-6.

- 1067 42. Algaflly AA, George KP. The effect of cryotherapy on nerve conduction velocity, pain
1068 threshold and pain tolerance. *Br J Sports Med.* 2007;41(6):365-9; discussion 9.
- 1069 43. Kwiecien SY, McHugh MP. The cold truth: the role of cryotherapy in the treatment of
1070 injury and recovery from exercise. *Eur J Appl Physiol.* 2021;121(8):2125-42.
- 1071 44. Dow J, Giesbrecht GG, Danzl DF, Brugger H, Sagalyn EB, Walpoth B, et al. Wilderness
1072 Medical Society Clinical Practice Guidelines for the Out-of-Hospital Evaluation and
1073 Treatment of Accidental Hypothermia: 2019 Update. *Wilderness Environ Med.*
1074 2019;30(4S):S47-S69.
- 1075 45. Kyaw S, Moore I, Oo M. A Systematic Review on the Effectiveness of Different
1076 Functional Treatments for Acute Ankle Sprains. *P Sports Med Dopng Stud.*
1077 2019;9(1):213.
- 1078 46. Amar P, Schiff E. Acetaminophen safety and hepatotoxicity--where do we go from here?
1079 *Expert Opin Drug Saf.* 2007;6(4):341-55.
- 1080 47. Ghanem CI, Perez MJ, Manautou JE, Mottino AD. Acetaminophen from liver to brain:
1081 New insights into drug pharmacological action and toxicity. *Pharmacol Res.*
1082 2016;109:119-31.
- 1083 48. Smith H. Potential analgesic mechanisms of acetaminophen. *Pain Physician.*
1084 2009;12(1):269-89.
- 1085 49. Zimmerman H, Maddrey W. Acetaminophen (paracetamol) hepatotoxicity with regular
1086 intake of alcohol: analysis of instances of therapeutic misadventure. *Hepatology.*
1087 1995;22(3):767-73.
- 1088 50. Rumack B. Acetaminophen hepatotoxicity: the first 35 years. *J Toxicol Clin Toxicol.*
1089 2002;40(1):3-20.

- 1090 51. Chiew AL, Buckley NA. Acetaminophen Poisoning. *Crit Care Clin.* 2021;37(3):543-61.
- 1091 52. Chiew AL, Buckley NA. Acetaminophen Poisoning. *Crit Care Clin.* 2021;37(3):543-61.
- 1092 53. Zimmerman H, Maddrey W. Acetaminophen (paracetamol) hepatotoxicity with regular
1093 intake of alcohol: analysis of instances of therapeutic misadventure. *Hepatology.*
1094 1995;22(3):767-73.
- 1095 54. Patel A, Pai BHP, Diskina D, Reardon B, Lai YH. Comparison of clinical outcomes of
1096 acetaminophen IV vs PO in the peri-operative setting for laparoscopic inguinal hernia
1097 repair surgeries: A triple-blinded, randomized controlled trial. *J Clin Anesth.*
1098 2020;61:109628.
- 1099 55. Tompkins DM, DiPasquale A, Segovia M, Cohn SM. Review of Intravenous
1100 Acetaminophen for Analgesia in the Postoperative Setting. *Am Surg.* 2021;87(11):1809-
1101 22.
- 1102 56. Bannwarth B, Péhourcq F. Pharmacologic basis for using paracetamol: pharmacokinetic
1103 and pharmacodynamic issues. *Drugs.* 2003;63(Spec No 2):5-13.
- 1104 57. Scialli AR, Ang R, Breitmeyer J, Royal MA. A review of the literature on the effects of
1105 acetaminophen on pregnancy outcome. *Reprod Toxicol.* 2010;30(4):495-507.
- 1106 58. Phillips W, Bradford C. Analgesic Pharmacology: II. Specific Analgesics. *J Am Acad*
1107 *Orthop Surg.* 2004;12(4):221-33.
- 1108 59. Vane J. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like
1109 drugs. *Nat New Biol.* 1971;231(25):232-5.
- 1110 60. Chaiamnuay S, Allison JJ, Curtis JR. Risks versus benefits of cyclooxygenase-2-selective
1111 nonsteroidal antiinflammatory drugs. *Am J Health Syst Pharm.* 2006;63(19):1837-51.

- 1112 61. Wheatley R. Analgesic efficacy of ketorolac. *Acta anaesthesiologica belgica*.
1113 1996;47(3):135-42.
- 1114 62. Gillis J, Brogden R. Ketorolac. A reappraisal of its pharmacodynamic and
1115 pharmacokinetic properties and therapeutic use in pain management. *Drugs*.
1116 1997;53(1):139-88.
- 1117 63. Sostres C, Gargallo CJ, Arroyo MT, Lanás A. Adverse effects of non-steroidal anti-
1118 inflammatory drugs (NSAIDs, aspirin and coxibs) on upper gastrointestinal tract. *Best*
1119 *Pract Res Clin Gastroenterol*. 2010;24(2):121-32.
- 1120 64. Rampal P, Moore N, Van Ganse E, Le Parc J-M, Wall R, Schneid H, Verriere F.
1121 Gastrointestinal Tolerability of Ibuprofen Compared with Paracetamol and Aspirin at
1122 Over-the-counter Doses. *J Int Med Res*. 2002;30:301-8.
- 1123 65. Le Parc J, Ganse E, Moore N, Schneid H, Verriere F. Comparative tolerability of
1124 paracetamol, aspirin and ibuprofen for short-term analgesia in patients with
1125 musculoskeletal conditions: results in 4291 patients. *Clin Rheumatol*. 2002;21(1):28-31.
- 1126 66. Schafer A. Effects of nonsteroidal anti-inflammatory therapy on platelets. *Am J Med*.
1127 1999;106(5B):25S-36S.
- 1128 67. Lewis SR, Nicholson A, Cardwell ME, Siviter G, Smith AF. Nonsteroidal anti-
1129 inflammatory drugs and perioperative bleeding in paediatric tonsillectomy. *Cochrane*
1130 *Database Syst Rev*. 2013(7):CD003591.
- 1131 68. Forrest J, Heitlinger E, Revell S. Ketorolac for Postoperative Pain Management in
1132 Children. *Drug Safety*. 1997;16(5):309-29.
- 1133 69. Whelton A. Nephrotoxicity of nonsteroidal anti-inflammatory drugs: physiologic
1134 foundations and clinical implications. *Am J Med*. 1999;106:13S-24S.

- 1135 70. Balestracci A, Ezquer M, Elmo ME, Molini A, Thorel C, Torrents M, Toledo I.
1136 Ibuprofen-associated acute kidney injury in dehydrated children with acute
1137 gastroenteritis. *Pediatr Nephrol*. 2015;30(10):1873-8.
- 1138 71. Motov S, Masoudi A, Drapkin J, Sotomayor C, Kim S, Butt M, et al. Comparison of Oral
1139 Ibuprofen at Three Single-Dose Regimens for Treating Acute Pain in the Emergency
1140 Department: A Randomized Controlled Trial. *Ann Emerg Med*. 2019;74(4):530-7.
- 1141 72. Turner NJ, Long DA, Bongiorno JR, Katoski TP, Jin LM, Horsch JP, Ahern BJ.
1142 Comparing two doses of intramuscular ketorolac for treatment of acute musculoskeletal
1143 pain in a military emergency department. *Am J Emerg Med*. 2021;50:142-7.
- 1144 73. Jones P, Lamdin R, Dalziel SR. Oral non-steroidal anti-inflammatory drugs versus other
1145 oral analgesic agents for acute soft tissue injury. *Cochrane Database Syst Rev*.
1146 2020;8:CD007789.
- 1147 74. Ong CK, Lirk P, Tan CH, Seymour RA. An evidence-based update on nonsteroidal anti-
1148 inflammatory drugs. *Clin Med Res*. 2007;5(1):19-34.
- 1149 75. Dwarica DS, Pickett SD, Zhao YD, Nihira MA, Quiroz LH. Comparing Ketorolac With
1150 Ibuprofen for Postoperative Pain: A Randomized Clinical Trial. *Female Pelvic Med*
1151 *Reconstr Surg*. 2020;26(4):233-8.
- 1152 76. Arora S, Wagner J, Herbert M. Myth: parenteral ketorolac provides more effective
1153 analgesia than oral ibuprofen. *Canadian Journal of Emergency Medicine*. 2007;9(1):30-2.
- 1154 77. Motov S, Masoudi A, Drapkin J, Sotomayor C, Kim S, Butt M, et al. Comparison of Oral
1155 Ibuprofen at Three Single-Dose Regimens for Treating Acute Pain in the Emergency
1156 Department: A Randomized Controlled Trial. *Ann Emerg Med*. 2019;74(4):530-7.

- 1157 78. Baranowski DC, Buchanan B, Dwyer HC, Gabriele JP, Kelly S, Araujo JA. Penetration
1158 and efficacy of transdermal NSAIDs in a model of acute joint inflammation. *J Pain Res.*
1159 2018;11:2809-19.
- 1160 79. Snizek PJ, Brodland DG, Zitelli JA. A randomized controlled trial comparing
1161 acetaminophen, acetaminophen and ibuprofen, and acetaminophen and codeine for
1162 postoperative pain relief after Mohs surgery and cutaneous reconstruction. *Dermatol*
1163 *Surg.* 2011;37(7):1007-13.
- 1164 80. Mitchell A, van Zanten SV, Inglis K, Porter G. A randomized controlled trial comparing
1165 acetaminophen plus ibuprofen versus acetaminophen plus codeine plus caffeine after
1166 outpatient general surgery. *J Am Coll Surg.* 2008;206(3):472-9.
- 1167 81. Clark E, Plint AC, Correll R, Gaboury I, Passi B. A randomized, controlled trial of
1168 acetaminophen, ibuprofen, and codeine for acute pain relief in children with
1169 musculoskeletal trauma. *Pediatrics.* 2007;119(3):460-7.
- 1170 82. Ong CK, Seymour RA, Lirk P, Merry AF. Combining paracetamol (acetaminophen) with
1171 nonsteroidal antiinflammatory drugs: a qualitative systematic review of analgesic
1172 efficacy for acute postoperative pain. *Anesth Analg.* 2010;110(4):1170-9.
- 1173 83. Black IH, McManus J. Pain management in current combat operations. *Prehosp Emerg*
1174 *Care.* 2009;13(2):223-7.
- 1175 84. Aldington DJ, McQuay HJ, Moore RA. End-to-end military pain management. *Philos*
1176 *Trans R Soc Lond B Biol Sci.* 2011;366(1562):268-75.
- 1177 85. Dalton MK, Semco RS, Ordoobadi AJ, Goralnick E, Chovanes J, Salim A, Jarman MP.
1178 Opioid administration in the prehospital setting for patients sustaining traumatic injuries:
1179 An evaluation of national emergency medical services data. *Injury.* 2022;53(9):2923-9.

- 1180 86. Cepeda MS, Farrar JT, Baumgarten M, Boston R, Carr DB, Strom BL. Side effects of
1181 opioids during short-term administration: effect of age, gender, and race. *Clin Pharmacol*
1182 *Ther.* 2003;74(2):102-12.
- 1183 87. Dinges HC, Otto S, Stay DK, Baumlein S, Waldmann S, Kranke P, et al. Side Effect
1184 Rates of Opioids in Equianalgesic Doses via Intravenous Patient-Controlled Analgesia: A
1185 Systematic Review and Network Meta-analysis. *Anesth Analg.* 2019;129(4):1153-62.
- 1186 88. ACEP. Policy Statement: Out-of-Hospital Use of Analgesia and Sedation. *Ann Emerg*
1187 *Med.* 2016;67(2):305-6.
- 1188 89. Gausche-Hill M, Brown K, Oliver Z, Sasson C, Dayan P, Eschmann N. An Evidence-
1189 based Guideline for Prehospital Analgesia in Trauma. *Prehosp Emerg Care.* 2014;18:25-
1190 34.
- 1191 90. Miller M, Barber CW, Leatherman S, Fonda J, Hermos JA, Cho K, Gagnon DR.
1192 Prescription opioid duration of action and the risk of unintentional overdose among
1193 patients receiving opioid therapy. *JAMA Intern Med.* 2015;175(4):608-15.
- 1194 91. Raffa RB. Pharmacology of oral combination analgesics: rational therapy for pain. *J Clin*
1195 *Pharm Ther.* 2001;26(4):257-64.
- 1196 92. Bijur PE, Friedman BW, Irizarry E, Chang AK, Gallagher EJ. A Randomized Trial
1197 Comparing the Efficacy of Five Oral Analgesics for Treatment of Acute Musculoskeletal
1198 Extremity Pain in the Emergency Department. *Ann Emerg Med.* 2021;77(3):345-56.
- 1199 93. Graudins A, Meek R, Parkinson J, Egerton-Warburton D, Meyer A. A randomised
1200 controlled trial of paracetamol and ibuprofen with or without codeine or oxycodone as
1201 initial analgesia for adults with moderate pain from limb injury. *Emerg Med Australas.*
1202 2016;28(6):666-72.

- 1203 94. Chang AK, Bijur PE, Esses D, Barnaby DP, Baer J. Effect of a Single Dose of Oral
1204 Opioid and Nonopioid Analgesics on Acute Extremity Pain in the Emergency
1205 Department: A Randomized Clinical Trial. *JAMA*. 2017;318(17):1661-7.
- 1206 95. Schug SA, Ting S. Fentanyl Formulations in the Management of Pain: An Update.
1207 *Drugs*. 2017;77(7):747-63.
- 1208 96. Lotsch J, Walter C, Parnham MJ, Oertel BG, Geisslinger G. Pharmacokinetics of non-
1209 intravenous formulations of fentanyl. *Clin Pharmacokinet*. 2013;52(1):23-36.
- 1210 97. Jandhyala R, Fullarton JR, Bennett MI. Efficacy of rapid-onset oral fentanyl formulations
1211 vs. oral morphine for cancer-related breakthrough pain: a meta-analysis of comparative
1212 trials. *J Pain Symptom Manage*. 2013;46(4):573-80.
- 1213 98. Brząkała J, Leppert W. The Role of Rapid Onset Fentanyl Products in the Management
1214 of Breakthrough Pain in Cancer Patients. *Pharmacological Reports*. 2019;71(3):438-42.
- 1215 99. Karlsen AP, Pedersen DM, Trautner S, Dahl JB, Hansen MS. Safety of intranasal
1216 fentanyl in the out-of-hospital setting: a prospective observational study. *Ann Emerg*
1217 *Med*. 2014;63(6):699-703.
- 1218 100. Rickard C, O'Meara P, McGrail M, Garner D, McLean A, Le Lievre P. A randomized
1219 controlled trial of intranasal fentanyl vs intravenous morphine for analgesia in the
1220 prehospital setting. *Am J Emerg Med*. 2007;25(8):911-7.
- 1221 101. Middleton PM, Simpson PM, Sinclair G, Dobbins TA, Math B, Bendall JC. Effectiveness
1222 of morphine, fentanyl, and methoxyflurane in the prehospital setting. *Prehosp Emerg*
1223 *Care*. 2010;14(4):439-47.

- 1224 102. Tanguay A, Lebon J, Hebert D, Begin F. Intranasal Fentanyl versus Subcutaneous
1225 Fentanyl for Pain Management in Prehospital Patients with Acute Pain: A Retrospective
1226 Analysis. *Prehosp Emerg Care*. 2020;24(6):760-8.
- 1227 103. Lynch TV, Callas PW, Peterson TD, Schlein SM. Intranasal Fentanyl for On-the-Hill
1228 Analgesia by Ski Patrol. *Wilderness Environ Med*. 2022;33(3):296-303.
- 1229 104. Borland M, Jacobs I, King B, O'Brien D. A randomized controlled trial comparing
1230 intranasal fentanyl to intravenous morphine for managing acute pain in children in the
1231 emergency department. *Ann Emerg Med*. 2007;49(3):335-40.
- 1232 105. Crellin D, Ling RX, Babl FE. Does the standard intravenous solution of fentanyl (50
1233 microg/mL) administered intranasally have analgesic efficacy? *Emerg Med Australas*.
1234 2010;22(1):62-7.
- 1235 106. Manjushree R, Lahiri A, Ghosh B, Laha A, Handa K. Intranasal fentanyl provides
1236 adequate postopera- tive analgesia in pediatric patients. *Canadian Journal of*
1237 *Anaesthesia*. 2002;49(2):190-3.
- 1238 107. Cole J, Shepherd M, Young P. Intranasal fentanyl in 1-3-year-olds: a prospective study of
1239 the effectiveness of intranasal fentanyl as acute analgesia. *Emerg Med Australas*.
1240 2009;21(5):395-400.
- 1241 108. Younge P, Nicol M, Kendall J, Harrington A. A prospective randomized pilot
1242 comparison of intranasal fentanyl and intramuscular morphine for analgesia in children
1243 presenting to the emergency department with clinical fractures. *Emergency Medicine*.
1244 1999;11(2):90-4.
- 1245 109. Dale O. Intranasal administration of opioids/fentanyl – Physiological and
1246 pharmacological aspects. *European Journal of Pain Supplements*. 2010;4(3):187-90.

- 1247 110. Hardy J, Lee S, Wilson C. Intranasal drug delivery by spray and drops. *J Pharm*
1248 *Pharmacol.* 1985;37(5):294-7.
- 1249 111. Daley-Yates P, Baker R. Systemic bioavailability of fluticasone propionate administered
1250 as nasal drops and aqueous nasal spray formulations. *Br J Clin Pharmacol.*
1251 2001;51(1):103-5.
- 1252 112. Helmers J, Noorduyn H, Van Leeuwen L, Zuurmond W. Comparison of intravenous and
1253 intranasal sufentanil absorption and sedation. *Canadian Journal of Anaesthesia.*
1254 1989;36(5):494-7.
- 1255 113. Steenblik J, Goodman M, Davis V, Gee C, Hopkins CL, Stephen R, Madsen T. Intranasal
1256 sufentanil for the treatment of acute pain in a winter resort clinic. *Am J Emerg Med.*
1257 2012;30(9):1817-21.
- 1258 114. Stephen R, Lingenfelter E, Broadwater-Hollifield C, Madsen T. Intranasal sufentanil
1259 provides adequate analgesia for emergency department patients with extremity injuries. *J*
1260 *Opioid Manag.* 2012;8(4):237-41.
- 1261 115. Wedmore IS, Kotwal RS, McManus JG, Pennardt A, Talbot TS, Fowler M, McGhee L.
1262 Safety and efficacy of oral transmucosal fentanyl citrate for prehospital pain control on
1263 the battlefield. *J Trauma Acute Care Surg.* 2012;73(6 Suppl 5):S490-5.
- 1264 116. Lewis P, Wright C, Hooper C. Opioid analgesia on the battlefield: a retrospective review
1265 of data from Operation HERRICK. *J R Army Med Corps.* 2018;164(5):328-31.
- 1266 117. Ellerton J, Greene M, Paal P. The use of analgesia in mountain rescue casualties with
1267 moderate or severe pain. *Emerg Med J.* 2013;30(6):501-5.

- 1268 118. Rauck RL, Oh DA, Singla N, Koch C, Parikh N, Nalamachu S, et al. Pharmacokinetics of
1269 Fentanyl Sublingual Spray in Opioid-Naive Participants: Results of a Phase 1, Multiple
1270 Ascending Dose Study. *Clin Drug Investig*. 2018;38(8):715-26.
- 1271 119. Kim SY, Buckenmaier CC, Howe EG, Choi KH. The Newest Battlefield Opioid,
1272 Sublingual Sufentanil: A Proposal to Refine Opioid Usage in the U.S. Military. *Mil Med*.
1273 2022;187(3-4):77-83.
- 1274 120. Miner JR, Rafique Z, Minkowitz HS, DiDonato KP, Palmer PP. Sufentanil sublingual
1275 tablet 30mcg for moderate-to-severe acute pain in the ED. *Am J Emerg Med*.
1276 2018;36(6):954-61.
- 1277 121. Hutchins JL, Leiman D, Rafique Z, DiDonato KP, Palmer PP. Pooled Dosing and
1278 Efficacy Analysis of the Sufentanil Sublingual Tablet 30 mcg Across Demographic
1279 Subgroups for the Management of Moderate-to-Severe Acute Pain. *J Perianesth Nurs*.
1280 2020;35(1):22-8.
- 1281 122. Miner JR, Melson T, Leiman D, Minkowitz HS, Chiang Y, DiDonato KP, Palmer PP.
1282 Pooled Phase III safety analysis of sufentanil sublingual tablets for short-term treatment
1283 of moderate-to-severe acute pain. *Pain Management*. 2019;9(3):259-71.
- 1284 123. Schauer SG, Mora A, Maddry J, Bebart V. Multicenter, prospective study of prehospital
1285 administration of analgesia in the U.S. combat theater of Afghanistan. *Prehosp Emerg*
1286 *Care*. 2017;21(6):744-9.
- 1287 124. Kuusniemi KS, Olkkola KT. Parenteral Opioids in emergency medicine - Are we treating
1288 pain adequately? *Scand J Pain*. 2011;2(4):185-6.

- 1289 125. Kiavialaitis GE, Muller S, Braun J, Rossler J, Spahn DR, Stein P, Kaserer A. Clinical
1290 practice of pre-hospital analgesia: An observational study of 20,978 missions in
1291 Switzerland. *Am J Emerg Med.* 2020;38(11):2318-23.
- 1292 126. Buck ML, Wiggins BS, Sesler JM. Intraosseous drug administration in children and
1293 adults during cardiopulmonary resuscitation. *Ann Pharmacother.* 2007;41(10):1679-86.
- 1294 127. Thomas SH, Rago O, Harrison T, Biddinger PD, Wedel SK. Fentanyl trauma analgesia
1295 use in air medical scene transports. *J Emerg Med.* 2005;29(2):179-87.
- 1296 128. Galinski M, Dolveck F, Borron SW, Tual L, Van Laer V, Lardeur JY, et al. A
1297 randomized, double-blind study comparing morphine with fentanyl in prehospital
1298 analgesia. *Am J Emerg Med.* 2005;23(2):114-9.
- 1299 129. Mion G, Villevieille T. Ketamine pharmacology: an update (pharmacodynamics and
1300 molecular aspects, recent findings). *CNS Neurosci Ther.* 2013;19(6):370-80.
- 1301 130. Trimmel H, Helbok R, Staudinger T, Jaksch W, Messerer B, Schochl H, Likar R. S(+)-
1302 ketamine : Current trends in emergency and intensive care medicine. *Wien Klin*
1303 *Wochenschr.* 2018;130(9-10):356-66.
- 1304 131. Balzer N, McLeod S, Walsh C, Grewal K. Low-dose Ketamine For Acute Pain Control in
1305 the Emergency Department: A Systematic Review and Meta-analysis. *Acad Emerg Med.*
1306 2021;28(4):444-54.
- 1307 132. Mo H, Campbell. M, Fertel B, Lam S, Wells E, Casserly E, Meldon S. Ketamine Safety
1308 and Use in the Emergency Department for Pain and Agitation/Delirium: A Health System
1309 Experience. *West J Emerg Med.* 2019;21(2):272-81.
- 1310 133. Sin B, Ternas T, Motov S. The use of subdissociative-dose ketamine for acute pain in the
1311 emergency department. *Acad Emerg Med.* 2015;22(3):251-7.

- 1312 134. Sandberg M, Hyldmo PK, Kongstad P, Dahl Friesgaard K, Raatiniemi L, Larsen R, et al.
1313 Ketamine for the treatment of prehospital acute pain: a systematic review of benefit and
1314 harm. *BMJ Open*. 2020;10(11):e038134.
- 1315 135. Bisanzo M, Nichols K, Hammerstedt H, Dreifuss B, Nelson SW, Chamberlain S, et al.
1316 Nurse-administered ketamine sedation in an emergency department in rural Uganda. *Ann*
1317 *Emerg Med*. 2012;59(4):268-75.
- 1318 136. Vanolli K, Hugli O, Eidenbenz D, Suter MR, Pasquier M. Prehospital Use of Ketamine in
1319 Mountain Rescue: A Survey of Emergency Physicians of a Single-Center Alpine
1320 Helicopter-Based Emergency Service. *Wilderness Environ Med*. 2020;31(4):385-93.
- 1321 137. Bishop R, Litch J, Stanton J. Ketamine anaesthesia at high altitude. *High Alt Med Biol*.
1322 2004;1(2):111-4.
- 1323 138. Green SM, Roback MG, Krauss B, Brown L, McGlone RG, Agrawal D, et al. Predictors
1324 of airway and respiratory adverse events with ketamine sedation in the emergency
1325 department: an individual-patient data meta-analysis of 8,282 children. *Ann Emerg Med*.
1326 2009;54(2):158-68 e1-4.
- 1327 139. Morgan MM, Perina DG, Acquisto NM, Fallat ME, Gallagher JM, Brown KM, et al.
1328 Ketamine Use in Prehospital and Hospital Treatment of the Acute Trauma Patient: A
1329 Joint Position Statement. *Prehosp Emerg Care*. 2021;25(4):588-92.
- 1330 140. Harmatz A. Local anesthetics: uses and toxicities. *Surg Clin North Am*. 2009;89(3):587-
1331 98.
- 1332 141. Birsa L, Verity P, Lee R. Evaluation of the effects of various chemicals on discharge of
1333 and pain caused by jellyfish nematocysts. *Comp Biochem Physiol C Toxicol Pharmacol*.
1334 2010;151(4):426-30.

- 1335 142. Ward NT, Darracq MA, Tomaszewski C, Clark RF. Evidence-based treatment of jellyfish
1336 stings in North America and Hawaii. *Ann Emerg Med.* 2012;60(4):399-414.
- 1337 143. Aksel G, Guler S, Dogan NO, Corbacioglu SK. A randomized trial comparing
1338 intravenous paracetamol, topical lidocaine, and ice application for treatment of pain
1339 associated with scorpion stings. *Hum Exp Toxicol.* 2015;34(6):662-7.
- 1340 144. Turgut K, Yavuz E, Gulacti U, Aydin I, Sonmez C, Aktas N, Arslan E. Comparison of
1341 Intravenous Paracetamol, Dexketoprofen Trometamol, or Topical Lidocaine Use for Pain
1342 Relief in Scorpion Stings: A Placebo-Controlled, Randomized Study. *Wilderness Environ*
1343 *Med.* 2022;33(4):379-85.
- 1344 145. Tran A, Koo J. Risk of systemic toxicity with topical lidocaine/prilocaine: a review. *J*
1345 *Drugs Dermatol.* 2014;13(9):1118-22.
- 1346 146. Königs I, Wenskus J, Boettcher J, Reinschagen K, Boettcher M. Lidocaine-Epinephrine-
1347 Tetracaine Gel Is More Efficient than Eutectic Mixture of Local Anesthetics and
1348 Mepivacaine Injection for Pain Control during Skin Repair in Children: A Prospective,
1349 Propensity Score Matched Two-Center Study. *Eur J Pediatr Surg.* 2020;30(6):512-6.
- 1350 147. Zaki H, Elarref M, Iftikhar H, Al-Marri N, Masood M, Fayed M, et al. Efficacy of Emla
1351 (Eutectic Mixture of Local Anaesthetics) and Let (Lidocaine, Epinephrine, Tetracaine)
1352 for Topical Use in Wound Management for Children: A Systematic Review and Meta-
1353 Analysis. *Cureus.* 2022;14(11):e31447.
- 1354 148. Bolt P, Barnett P, Babl FE, Sharwood LN. Topical lignocaine for pain relief in acute
1355 otitis media: results of a double-blind placebo-controlled randomised trial. *Arch Dis*
1356 *Child.* 2008;93(1):40-4.

- 1357 149. Willman G. Ultraviolet Keratitis: From the Pathophysiological Basis to Prevention and
1358 Clinical Management. *High Alt Med Biol.* 2015;16(4):277-82.
- 1359 150. Wilson C. Specialised Pharmaceutical Formulation: The Science and Technology of
1360 Dosage Forms. In: Tovey G, ed.: RSC Publishing; 2022:1-44.
- 1361 151. Page M, Fraunfelder F. Safety, efficacy, and patient acceptability of lidocaine
1362 hydrochloride ophthalmic gel as a topical ocular anesthetic for use in ophthalmic
1363 procedures. *Clinical Ophthalmology.* 2009;3:601-9.
- 1364 152. Valentini C, Lin J, Golub JS, Lustig L. Price Differences Between Otic and Ophthalmic
1365 Drops. *Otol Neurotol.* 2021;42(2):274-7.
- 1366 153. Swaminathan A, Otterness K, Milne K, Rezaie S. The Safety of Topical Anesthetics in
1367 the Treatment of Corneal Abrasions: A Review. *J Emerg Med.* 2015;49(5):810-5.
- 1368 154. Puls HA, Cabrera D, Murad MH, Erwin PJ, Bellolio MF. Safety and Effectiveness of
1369 Topical Anesthetics in Corneal Abrasions: Systematic Review and Meta-Analysis. *J*
1370 *Emerg Med.* 2015;49(5):816-24.
- 1371 155. Akorn. Proparacaine Hydrochloride Ophthalmic Solution. Package Insert. 2000.
- 1372 156. Gadsden J. Prehospital regional anesthesia. In: *Regional Anesthesia in Trauma: A Case-*
1373 *Based Approach.* Cambridge University Press; 2012:11-8.
- 1374 157. Fleming I, Egeler C. Regional anaesthesia for trauma: an update. *Continuing Education*
1375 *in Anaesthesia Critical Care & Pain.* 2014;14(3):136-41.
- 1376 158. Hofmeyr R. Field Expedient Regional Blocks (FERBs). Wilderness Medicine Society
1377 Summer Conference; 2022; Snowmass.
- 1378 159. Whiffler K. Coracoid block--a safe and easy technique. *Br J Anaesth.* 1981;53(8):845-8.

- 1379 160. Bocquet J, N'Takpe N, Draganescu C, Ridarch A, Jullien Y. [The coracoid block:
1380 demonstration of a simple approach using the pectoralis minor as landmark]. *Can J*
1381 *Anaesth*. 2005;52(10):1040-6.
- 1382 161. Simpson PM, McCabe B, Bendall JC, Cone DC, Middleton PM. Paramedic-performed
1383 digital nerve block to facilitate field reduction of a dislocated finger. *Prehosp Emerg*
1384 *Care*. 2012;16(3):415-7.
- 1385 162. Lefort H, Mendibil A, Romanat P, Tourtier J. Anesthésie locorégionale préhospitalière :
1386 le bloc iliofascial. *Ann Fr Med Urgence*. 2013;3:363-6.
- 1387 163. Dochez E, van Geffen G, Bruh J, Hoogerwerf N, de Pas H, Scheffer G. Prehospital
1388 administered fascia iliaca compartment block by emergency medical service nurses, a
1389 feasibility study. *Scand J trauma Resusc Emerg*. 2014;22(1):38.
- 1390 164. Hards M, Brewer A, Bessant G, Lahiri S. Efficacy of Prehospital Analgesia with Fascia
1391 Iliaca Compartment Block for Femoral Bone Fractures: A Systematic Review. *Prehosp*
1392 *Disaster Med*. 2018;33(3):299-307.
- 1393 165. Hofmeyr R. Anesthesia outside a hospital setting. SASA Practice Guidelines. *S Afr J*
1394 *Anaesth Analg*. 2022.
- 1395 166. Case R. Haematoma block—a safe method of reducing Colles' fractures. *Injury*.
1396 1985;16(7):469-70.
- 1397 167. Ross A, Catanzariti AR, Mendicino RW. The hematoma block: a simple, effective
1398 technique for closed reduction of ankle fracture dislocations. *J Foot Ankle Surg*.
1399 2011;50(4):507-9.

- 1400 168. Dezfuli B, Edwards C, DeSilva G. Distal radius fracture hematoma block with combined
1401 lidocaine and bupivacaine can induce seizures while within therapeutic window. *J Orthop*
1402 *Case Rep.* 2012;2(4):10-3.
- 1403 169. Myderrizi N, Mema B. The hematoma block an effective alternative for fracture
1404 reduction in distal radius fractures. *Med Arh.* 2011;65(4):239-42.
- 1405 170. Maleitzke T, Plachel F, Fleckenstein FN, Wichlas F, Tsitsilonis S. Haematoma block: a
1406 safe method for pre-surgical reduction of distal radius fractures. *J Orthop Surg Res.*
1407 2020;15(1):351.
- 1408 171. Pavlidakey P, Brodell E, Helms S. Diphenhydramine as an alternative local anesthetic
1409 agent. *J Clin Aesthetic Dermatol.* 2009;2(10):37-40.
- 1410 172. Green SM. What is the role of diphenhydramine in local anesthesia? *Acad Emerg Med.*
1411 1996;3(3):198-200.
- 1412 173. Bartfield J, Jandreau S, Raccio-Robak N. Randomized trial of diphenhydramine versus
1413 benzyl alcohol with epinephrine as an alternative to lidocaine local anesthesia. *Ann*
1414 *Emerg Med.* 1998;32(6):650-4.
- 1415 174. Ernst A, Anand P, Nick T, Wassmuth S. Lidocaine versus diphenhydramine for
1416 anesthesia in the repair of minor lacerations. *J Trauma.* 1993;34(3):354-7.
- 1417 175. Dire D, Hogan D. Double-blinded comparison of diphenhydramine versus lidocaine as a
1418 local anesthetic. *Ann Emerg Med.* 1993;22(9):1419-22.
- 1419 176. Silva E, Scherber K, Cabrera D, Motov S, Erwin PJ, West CP, et al. Safety and Efficacy
1420 of Intravenous Lidocaine for Pain Management in the Emergency Department: A
1421 Systematic Review. *Ann Emerg Med.* 2018;72(2):135-44 e3.

- 1422 177. Masic D, Liang E, Long C, Sterk EJ, Barbas B, Rech MA. Intravenous Lidocaine for
1423 Acute Pain: A Systematic Review. *Pharmacotherapy*. 2018;38(12):1250-9.
- 1424 178. Foo I, Macfarlane AJR, Srivastava D, Bhaskar A, Barker H, Knaggs R, et al. The use of
1425 intravenous lidocaine for postoperative pain and recovery: international consensus
1426 statement on efficacy and safety. *Anaesthesia*. 2021;76(2):238-50.
- 1427 179. Dryden B, Kerr WB, Higgins S, Tou K, Dhanjal ST. The Use of Intravenous Lidocaine as
1428 an Analgesic Modality in the Austere Environment: Two Cases. *Wilderness Environ*
1429 *Med*. 2022;33(3):348-50.
- 1430 180. Ducasse JL, Siksik G, Durand-Bechu M, Couarraze S, Valle B, Lecoules N, et al. Nitrous
1431 oxide for early analgesia in the emergency setting: a randomized, double-blind
1432 multicenter prehospital trial. *Acad Emerg Med*. 2013;20(2):178-84.
- 1433 181. Knuf K, Maani C. Nitrous Oxide. In: *StatPearls [Internet]*. StatPearls Publishing; 2022.
- 1434 182. Faddy SC, Garlick SR. A systematic review of the safety of analgesia with 50% nitrous
1435 oxide: can lay responders use analgesic gases in the prehospital setting? *Emerg Med J*.
1436 2005;22(12):901-8.
- 1437 183. Kuivalainen AM, Ebeling F, Poikonen E, Rosenberg PH. Nitrous oxide analgesia for
1438 bone marrow aspiration and biopsy - A randomized, controlled and patient blinded study.
1439 *Scand J Pain*. 2015;7(1):28-34.
- 1440 184. Heinrich M, Menzel C, Hoffman F, Berger M, von Schweinitz D. Self-administered
1441 procedural analgesia using nitrous oxide/oxygen (50:50) in the pediatric surgery
1442 emergency room: effectiveness and limitations. *Eur J Pediatr Surg*. 2015;25(3):250-6.

- 1443 185. Kariman H, Majidi A, Amini A, Dolatabadi AA, Derakhshanfar H, Hatamabadi H, et al.
1444 Nitrous oxide/oxygen compared with fentanyl in reducing pain among adults with
1445 isolated extremity trauma: a randomized trial. *Emerg Med Australas*. 2011;23(6):761-8.
- 1446 186. Masoumi K, Forouzan A, Saghari S, Feli M, Sattari AR, Asgari Darian A. Sedative and
1447 Analgesic Effects of Entonox Gas Compared with Midazolam and Fentanyl in
1448 Synchronized Cardioversion. *Crit Care Res Pract*. 2015;2015:798478.
- 1449 187. Xing Y, Zhou L, Yu J, Wang Z, Ding Z, Xie C, et al. Analgesic efficacy of nitrous oxide
1450 in adults in the emergency department: A meta-analysis of randomized controlled trials.
1451 *Am J Emerg Med*. 2022;56:92-9.
- 1452 188. O’Sullivan I, Bengner J. Nitrous Oxide in Emergency Medicine. *Emerg Med J*.
1453 2003;20(3):214-7.
- 1454 189. James MF, Hofmeyr R, Grocott MP. Losing concentration: time for a new MAPP? *Br J*
1455 *Anaesth*. 2015;115(6):824-6.
- 1456 190. James MF, Manson ED, Dennett JE. Nitrous oxide analgesia and altitude. *Anaesthesia*.
1457 1982;37(3):285-8.
- 1458 191. James MF, White J. Anesthetic Considerations at Moderate Altitude. *Anesth Analg*.
1459 1984;63(12):1097-105.
- 1460 192. Wood C, Arbet J, Amura C, Nodine P, Collins M, Orlando B, et al. Multicenter Study
1461 Evaluating Nitrous Oxide Use for Labor Analgesia at High- and Low-Altitude
1462 Institutions. *Anesth & Analg*. 2022;134(2):294-302.
- 1463 193. Federation Internationale des Patrouilles de Ski 2014 Congress Part B - Detailed Activity
1464 Report. Paper presented at: FIPS World Congress2014; Big White, Canada.

1465 194. Porter KM, Dayan AD, Dickerson S, Middleton PM. The role of inhaled methoxyflurane
1466 in acute pain management. *Open Access Emerg Med.* 2018;10:149-64.

1467 195. Jephcott C, Grummet J, Nguyen N, Spruyt O. A review of the safety and efficacy of
1468 inhaled methoxyflurane as an analgesic for outpatient procedures. *Br J Anaesth.*
1469 2018;120(5):1040-8.

1470 196. Coffey F, Wright J, Hartshorn S, Hunt P, Locker T, Mirza K, Dissman P. STOP!: a
1471 randomised, double-blind, placebo-controlled study of the efficacy and safety of
1472 methoxyflurane for the treatment of acute pain. *Emerg Med J.* 2014;31:613-8.

1473 197. Coffey F, Dissmann P, Mirza K, Lomax M. Methoxyflurane Analgesia in Adult Patients
1474 in the Emergency Department: A Subgroup Analysis of a Randomized, Double-blind,
1475 Placebo-controlled Study (STOP!). *Adv Ther.* 2016;33(11):2012-31.

1476 198. Oxer H. Effects of Pentrox® (methoxyflurane) as an analgesic on cardiovascular and
1477 respiratory functions in the pre-hospital setting. *J Mil Veteran's Health.* 24(2):14-20.

1478 199. Hartshorn S, Dissmann P, Coffey F, Lomax M. Low-dose methoxyflurane analgesia in
1479 adolescent patients with moderate-to-severe trauma pain: a subgroup analysis of the
1480 STOP! study. *J Pain Res.* 2019;12:689-700.

1481 200. Windsor J, van der Kaaij J, Ellerton J, Oxer H, Hillebrandt D, Rodway G.
1482 Methoxyflurane as an analgesic for prehospital use at high altitude. *High Alt Med Biol.*
1483 2009;10(2):201-2.

1484 201. Hofmeyr R, Moon R, Dong T. Influence of altitude on performance of the
1485 methoxyflurane inhalational analgesic device: a hypobaric laboratory assessment. *J Pain*
1486 *S Africa.* 2019;14(2):11.

- 1487 202. Wilkes M, Heath EC, Mason NP. Methoxyflurane for Procedural Analgesia at 4470 m
1488 Altitude. *Wilderness Environ Med.* 2018;29(3):388-91.
- 1489 203. Mercadante S, Voza A, Serra S, Ruggiano G, Carpinteri G, Gangitano G, et al. Analgesic
1490 Efficacy, Practicality and Safety of Inhaled Methoxyflurane Versus Standard Analgesic
1491 Treatment for Acute Trauma Pain in the Emergency Setting: A Randomised, Open-Label,
1492 Active-Controlled, Multicentre Trial in Italy (MEDITA). *Adv Ther.* 2019;36(11):3030-
1493 46.
- 1494 204. Borobia AM, Collado SG, Cardona CC, Pueyo RC, Alonso CF, Torres IP, et al. Inhaled
1495 Methoxyflurane Provides Greater Analgesia and Faster Onset of Action Versus Standard
1496 Analgesia in Patients With Trauma Pain: InMEDIATE: A Randomized Controlled Trial
1497 in Emergency Departments. *Ann Emerg Med.* 2020;75(3):315-28.
- 1498 205. Voza A, Ruggiano G, Serra S, Carpinteri G, Gangitano G, Intelligente F, et al. Inhaled
1499 Methoxyflurane versus Intravenous Morphine for Severe Trauma Pain in the Emergency
1500 Setting: Subgroup Analysis of MEDITA, a Multicenter, Randomized, Controlled, Open-
1501 Label Trial. *J Pain Res.* 2020;13:491-502.
- 1502 206. Smith MD, Rowan E, Spaight R, Siriwardena AN. Evaluation of the effectiveness and
1503 costs of inhaled methoxyflurane versus usual analgesia for prehospital injury and trauma:
1504 non-randomised clinical study. *BMC Emerg Med.* 2022;22(1):122.
- 1505 207. Hofmeyr R. Methoxyflurane: inspiration of the old to breathe life into the new. *S Afr J*
1506 *Anaesth Analg.* 2020;26(5):218-22.
- 1507 208. Bussienne F, Reynaud T. Methoxyflurane May Be a Suitable Analgesic for Extreme
1508 Prehospital Conditions. *Wilderness Environ Med.* 2021;32(2):261-4.

- 1509 209. Trimmel H, Egger A, Doppler R, Pimiskern M, Voelckel WG. Usability and
1510 effectiveness of inhaled methoxyflurane for prehospital analgesia - a prospective,
1511 observational study. *BMC Emerg Med.* 2022;22(1):8.
- 1512 210. Guina J, Merrill B. Benzodiazepines I: Upping the Care on Downers: The Evidence of
1513 Risks, Benefits and Alternatives. *J Clin Med.* 2018;7(2).
- 1514 211. Bailey P, Pace N, Ashburn M, Moll J, East K, Stanley T. Frequent hypoxemia and apnea
1515 after sedation with midazolam and fentanyl. *Anesthesiology.* 1990;73(5):826-30.
- 1516 212. Horsfall JT, Sprague JE. The Pharmacology and Toxicology of the 'Holy Trinity'. *Basic*
1517 *Clin Pharmacol Toxicol.* 2017;120(2):115-9.
- 1518 213. Boon M, van Dorp E, Broens S, Overdyk F. Combining opioids and benzodiazepines:
1519 effects on mortality and severe adverse respiratory events. *Ann Palliat Med.*
1520 2020;9(2):542-57.
- 1521
1522