

Central Lancashire Online Knowledge (CLOK)

Title	A recursive formula for the product of element orders of finite abelian groups
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/51785/
DOI	doi:10.46298/cm.10996
Date	2023
Citation	Saha, Subhrajyoti (2023) A recursive formula for the product of element orders of finite abelian groups. Communications in Mathematics, 32 (1). ISSN 2336-1298
Creators	Saha, Subhrajyoti

It is advisable to refer to the publisher's version if you intend to cite from the work.
doi:10.46298/cm.10996

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

A recursive formula for the product of element orders of finite abelian groups

Subhrajyoti Saha

Abstract. Let G be a finite group and let $\psi(G)$ denote the sum of element orders of G ; later this concept has been used to define $R(G)$ which is the product of the element orders of G . Motivated by the recursive formula for $\psi(G)$, we consider a finite abelian group G and obtain a similar formula for $R(G)$.

1 Introduction

Let G be a finite group. For any non-empty subset S of G , let $\psi(S)$ denote the sum of element orders of S . This has been introduced in [2] and later in [4], the notion $R(G)$ was introduced which stands for the product of element orders of G . In the same paper, a formula for computing $R(G)$ when G is a finite abelian group was obtained. In [3], [5], an explicit recursive formula for computing $\psi(G)$ were obtained in case G is abelian. Motivated by these results, in this paper, we obtain a similar recursive formula for computing $R(G)$ when G is a finite abelian group.

Throughout this paper, we let $\varphi(n)$ denote the Euler totient function of the positive integer n and let p denote a prime number. A cyclic group of order n will be denoted by C_n whereas $C_p^{(r)}$ will denote the elementary abelian p -group of rank r . We always assume G to be finite. For a group G and element $x \in G$, the notation $o(x)$ denotes the order of x . For any group G , we take

$$R(G) = \prod_{x \in G} o(x).$$

For a group G , the notation $\exp(G)$ denotes the exponent of G which is the smallest positive integer z such that $g^z = 1_G$ for all $g \in G$ where 1_G is the identity element of G ; without any ambiguity we will denote this identity element as 1.

MSC 2020: 20D60, 20K01

Keywords: Finite group; Abelian group; Element order

Affiliation:

Monash University, Australia and University of Liverpool, UK

E-mail: subhrajyotimath@gmail.com

2 Explicit formulas for finite abelian groups

In this section, we obtain explicit recursive formula for $R(G)$ where G is a finite abelian group. We present this in different cases starting from a finite cyclic group. This format is inspired by [3, Section 2]. We will then consider the direct product of a finite cyclic p -group and a (not necessarily abelian) p -group. Finally, we will consider a the most general case of finite abelian groups. The proofs of our results in this section are motivated by the methods used in [3]. We begin with the following important preliminary results.

Theorem 2.1 ([4, Proposition 1.1]). *Let $G_1, G_2, \dots, G_k$ be finite groups having co-prime orders and $G \cong G_1 \times G_2 \times \dots \times G_k$. Then*

$$R(G) = \prod_{i=1}^k R(G_i)^{n_i}, \quad \text{where } n_i = \prod_{j=1, j \neq i}^k |G_j|, i = 1, 2, \dots, k.$$

Lemma 2.2 ([3, Lemma 2.6]). *Let $H \cong C_{p^{r_1}} \times C_{p^{r_2}} \times \dots \times C_{p^{r_n}}$ where $1 \leq r_1 \leq r_j$ for all j with $2 \leq j \leq n$. Then for any $i \in \{1, \dots, r_1\}$, there are $(p^i)^n - (p^{i-1})^n$ elements of H of order p^i*

The following lemma, motivated by [3, Lemma 1.1], follows easily from the fact that C_n has exactly $\varphi(d)$ elements of order d for each divisor d of n .

Lemma 2.3. *Let n be any positive integer. Then*

$$\psi(C_n) = \sum_{d|n} d\varphi(d) \quad \text{and} \quad R(C_n) = \prod_{d|n} d^{\varphi(d)}$$

2.1 Finite Cyclic Groups

Let G be a cyclic group of order n . Then we know that $G \cong C_{m_1} \times \dots \times C_{m_k}$, where $m_1, \dots, m_k$ are co-prime to each other and $n = m_1 \dots m_k$.

Lemma 2.4. *Let G be a cyclic group of order p^n where p is a prime number and n is a positive integer, then $R(G) = p^{\left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)}$.*

Proof. Using Lemma 2.3, we get $R(G) = \prod_{r=1}^n p^{\varphi(p^r)r} = p^{\sum_{r=1}^n \varphi(p^r)r}$. So we have $R(G) = p^z$ where $z = \sum_{r=1}^n r(p^r - p^{r-1}) = (1 - \frac{1}{p}) \sum_{r=1}^n rp^r$. Now the sum $\sum_{r=1}^n rp^r$ is an arithmetic-geometric series with common difference 1 and common ration p . Thus

$$\sum_{r=1}^n rp^r = \frac{p + np^{n+2} - (n+1)p^{n+1}}{(p-1)^2}.$$

This shows that $z = \frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}$. So we get $R(G) = p^{\left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)}$. □

Lemma 2.5. *Let G be a cyclic group of order $s = p_1^{r_1} \dots p_k^{r_k}$ where the p_i are distinct primes with $r_i \geq 1$ for $i = 1, \dots, k$. Then*

$$R(G) = \prod_{i=1}^k p_i^{\left(\frac{p_i + r_i p_i^{r_i+2} - (r_i+1)p_i^{r_i+1}}{p_i(p_i-1)} \right) n_i}, \quad n_i = p_i^{-r_i} \prod_{j=1}^k p_j^{r_j}.$$

Proof. We know that $G \cong C_{p_1^{r_1}} \times \dots \times C_{p_k^{r_k}}$ and p_i are all distinct primes. Hence we can apply Theorem 2.1. Thus by Lemma 2.4, we arrive at the required result. $\square$

2.2 Direct product of a cyclic p -Group and a p -Group

In this section, we obtain a recursive formula for $R(G)$ where G is a direct product of a finite cyclic p -group and any p -group. We begin with the following lemma from [3, Lemma 2.3].

Lemma 2.6. *If H and K are p -groups, then $o((x_1, x_2)) = \max \{o(x_1), o(x_2)\}$ for all for any $x_1 \in H$ and $x_2 \in K$.*

We now prove the following using techniques motivated by [3].

Proposition 2.7. *Let $G = C_{p^r} \times H$ where $r \geq 1$, and H is a p -group with $\exp(H) \geq p^r$. Let N_j be the number of elements in H that have order p^j . Then*

$$R(G) = \begin{cases} p^{p-1} R(H)^{p^r} \prod_{i=2}^r \left(\prod_{j=1}^{i-1} (p^{i-j})^{N_j} \right)^{p^i - p^{i-1}}, & \text{if } r > 1 \\ p^{p-1} R(H)^p, & \text{if } r = 1 \end{cases}$$

Proof. Note that G is a finite group whose elements are of the form (x, y) where $x \in C_{p^r}$ and $y \in H$. We now partition G based on the order of the elements in the first component. In particular we have, $G = \bigcup_{k=0}^r F_k$ where $F_k = \{(x_1, x_2) \in G \mid o(x_1) = p^k\}$. Since the $F_i \cap F_j = \emptyset$ for $i \neq j$, we have $R(G) = \prod_{k=0}^r R(F_k)$. Now let $x_1 \in C_{p^r}$ with $o(x_1) = p^i$ for some i with $0 \leq i \leq r$. For each such x_1 , define $F_{i,x_1} = \{(x_1, x_2) \mid x_2 \in H\}$. Then we have $F_i = \bigcup_{x_1 \in C_{p^r}, o(x_1)=p^i} F_{i,x_1}$ and $R(F_i) = \prod_{x_1 \in C_{p^r}, o(x_1)=p^i} R(F_{i,x_1})$ for $i = 0, 1, \dots, r$. There are $(p^i - p^{i-1})$ elements of order p^i in C_{p^r} , see Lemma 2.2. As a result,

$$R(F_i) = R(F_{i,x_1})^{(p^i - p^{i-1})}.$$

Taking $i = 0$, we have $F_0 = F_{0,1}$ and thus, $R(F_0) = R(H)$. For $i = 1$, each element (x_1, x_2) in F_{1,x_1} has order same as $o(x_2)$ except for $(x_1, 1)$ which has order p . Thus $R(F_{1,x_1}) = pR(H)$. So $R(F_1) = (pR(H))^{p-1}$.

If $r = 1$ then $R(G) = R(F_0) R(F_1) = p^{p-1} R(H)^p$.

Now consider $r > 1$ and let $i \in \{2, \dots, r\}$. For $(x_1, x_2) \in F_{i,x_1}$ with $o(x_2) = p^j$, we have

$o((x_1, x_2)) = p^i$ if $j < i$ and $o((x_1, x_2)) = p^j$ if $j \geq i$. If $r' = \exp(H)$ then

$$\begin{aligned} R(F_{i,x_1}) &= \prod_{j=0}^{i-1} (p^i)^{N_j} \prod_{j=i}^{r'} (p^j)^{N_j} \\ &= p^i \prod_{j=1}^{i-1} (p^i)^{N_j} \prod_{j=i}^{r'} (p^j)^{N_j} \\ &= p^i \prod_{j=1}^{r'} (p^j)^{N_j} \prod_{j=1}^{i-1} (p^{i-j})^{N_j} \\ &= p^i R(H) \prod_{j=1}^{i-1} (p^{i-j})^{N_j} \end{aligned}$$

Thus $\prod_{i=2}^r R(F_i) = R(H)^{(p^r-p)} \prod_{i=2}^r \left(\prod_{j=1}^{i-1} (p^{i-j})^{N_j} \right)^{p^i-p^{i-1}}$ and finally the result follows from a direct calculation $R(G) = R(F_0) R_l(F_1) \prod_{i=2}^r R_l(F_i)$. $\square$

2.3 Finite abelian groups

We can now state how to compute $R(G)$ for any finite abelian group G . In view of Theorem 2.1, the following is a direct application of Proposition 2.7 and Lemma 2.2.

Theorem 2.8. *Let G be a finite abelian group with $G \cong H_1 \times \cdots \times H_k$ where each H_i is an abelian p_i -group and p_i are distinct primes for $i = 1, \dots, k$. Then*

$$R(G) = R(H_1) \dots R(H_k)$$

where $R(H_i)$ for $i = 1, \dots, k$ are computed as follows:

a) If $H_i \cong C_{p_i^n}$ then

$$R(H_i) = p_i^{\left(\frac{p_i + n p_i^{n+2} - (n+1) p_i^{n+1}}{p_i(p_i-1)} \right)}.$$

b) If $H_i \cong C_{p_i^{r_1}} \times C_{p_i^{r_2}} \times \cdots \times C_{p_i^{r_n}}$, where $1 \leq r_1 \leq r_2 \leq \cdots \leq r_n$, and $r_1 + \dots + r_n = r$ then $R(H_i)$ can be determined recursively as follows

i) If $r_1 > 1$ then

$$R(H_i) = p_i^{p_i-1} R(C_{p_i^{r_2}} \times \cdots \times C_{p_i^{r_n}})^{p^{r_1}} \prod_{z=2}^{r_1} \left(\prod_{j=1}^{z-1} (p_i^{z-j})^{N_j} \right)^{p_i^z - p_i^{z-1}},$$

$$\text{where } N_j = \left((p_i^j)^{n-1} - (p_i^{j-1})^{n-1} \right)$$

ii) If $r_1 = 1$ then

$$R(H_i) = p_i^{p_i-1} R(C_{p_i^{r_2}} \times \cdots \times C_{p_i^{r_n}})^{p_i}.$$

3 Some Examples

In this final section we compute some examples using Theorem 2.8.

Example 3.1. We compute $R(C_p^{(r)} \times C_{p^n})$ where n and r are positive integers. By part a) of Theorem 2.8 we have

$$R(C_{p^n}) = p^{\left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)}.$$

Then by part b) of Theorem 2.8 we have

$$R(C_p \times C_{p^n}) = p^{p-1} p^{\left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)} = p^{\left(\frac{(p-1)^2+(p+np^{n+2}-(n+1)p^{n+1})}{(p-1)}\right)}.$$

Similarly

$$R(C_p \times C_p \times C_{p^n}) = p^{\left(\frac{(p-1)^2(1+p)+p(p+np^{n+2}-(n+1)p^{n+1})}{(p-1)}\right)}.$$

Thus inductively it is easy to show that

$$R(C_p^{(r)} \times C_{p^n}) = p^{\left(\frac{(p-1)^2(1+p+\dots+p^{r-1})+p^{r-1}(p+np^{n+2}-(n+1)p^{n+1})}{(p-1)}\right)}.$$

Thus we have

$$R(C_p^{(r)} \times C_{p^n}) = p^{\left(\frac{(p-1)(p^r-1)+p^{r-1}(p+np^{n+2}-(n+1)p^{n+1})}{(p-1)}\right)}.$$

The next example is an application of Theorem 2.8 in the case where $r_1 > 1$.

Example 3.2. In this example we compute $R(C_{p^2} \times C_{p^n})$ where n is a positive integers. By part a) of Theorem 2.8 we have

$$R(C_{p^n}) = p^{\left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)}.$$

Then by part b) of Theorem 2.8 we have

$$R(C_{p^2} \times C_{p^n}) = p^{p-1} p^{p^2 \left(\frac{p+np^{n+2}-(n+1)p^{n+1}}{p(p-1)}\right)} p^{p(p-1)(p^{n-1}-1)}.$$

This gives

$$R(C_{p^2} \times C_{p^n}) = p^{\left(\frac{(p-1)^2(p^n+1-p)+p(p+np^{n+2}-(n+1)p^{n+1})}{p-1}\right)}.$$

In the following example we compute $R(C_p^{(r)} \times C_{p^2} \times C_{p^n})$ where n and r are positive integers with $n \geq 2$.

Example 3.3. In example 3.2 we have computed $R(C_{p^2} \times C_{p^n})$. Then by part b) of Theorem 2.8 we have

$$R(C_p \times C_{p^2} \times C_{p^n}) = p^{p-1} p^{\left(\frac{(p-1)^2(p^n+1-p)+p(p+np^{n+2}-(n+1)p^{n+1})}{p-1}\right)}.$$

This gives

$$R(C_p \times C_{p^2} \times C_{p^n}) = p^{\left(\frac{(p-1)^2(1+p-p^2+p^{n+1})+p^2(p+np^{n+2}-(n+1)p^{n+1})}{p-1}\right)}.$$

Then inductively one can show that

$$R(C_p^{(r)} \times C_{p^2} \times C_{p^n}) = p^{\left(\frac{(p-1)^2(1+p+\dots+p^r-p^{r+1}+p^{n+r})+p^{r+1}(p+np^{n+2}-(n+1)p^{n+1})}{p-1}\right)}$$

where n and r are integers with $n \geq 2$.

Note that formulas obtained in Examples 3.1, 3.2 and 3.3 are straightforward to compute even when n and r are large. For any finite abelian p -group, an explicit formula for computing $R(G)$ was obtained in [4, Theorem 1.1]. By expanding and simplifying the formula obtained in [4, Theorem 1.1] one can compare the computations for $R(C_p^{(r)} \times C_{p^2} \times C_{p^n})$; the recursive formula used in Example 3.3 provides a more efficient method for obtaining $R(C_p^{(r)} \times C_{p^2} \times C_{p^n})$.

References

- [1] Amiri H. and Jafarian Amiri S. M. : Sum of element orders on finite groups of the same order. *Journal of Algebra and its Applications* 10 (2) (2011) 187–190.
- [2] Amiri H., Jafarian Amiri S. M. and Isaacs I. M.: Sums of element orders in finite groups. *Communications in Algebra* 37 (9) (2009) 2978–2980.
- [3] Chew C. Y., Chin A. Y. M. and Lim C. S.: A recursive formula for the sum of element orders of finite abelian groups. *Results in Mathematics* 72 (4) (2017) 1897–1905.
- [4] Tarnauceanu M.: A note on the product of element orders of finite abelian groups. *Bulletin of the Malaysian Mathematical Sciences Society. Second Series* 36 (4) (2013) 1123–1126.
- [5] Saha S.: Sum of the powers of the orders of elements in finite abelian groups. *Advances in Group Theory and Applications* 13 (2022) 1–11.

Received: April 12, 2021

Accepted for publication: June 25, 2022

Communicated by: Pasha Zusmanovich