

Central Lancashire Online Knowledge (CLoK)

Title	The use of accelerometers to improve clinical outcomes in patients with implantable cardioverter defibrillators or at high risk of sudden cardiac death	
Туре	Article	
URL	https://clok.uclan.ac.uk/id/eprint/53884/	
DOI	10.12968/bjca.2024.0068	
Date	2025	
Citation	Cheng, Hannah Yee Yan and Hill, James Edward (2025) The use of accelerometers to improve clinical outcomes in patients with implantable cardioverter defibrillators or at high risk of sudden cardiac death. British Journal of Cardiac Nursing. ISSN 1749-6403	
Creators	Cheng, Hannah Yee Yan and Hill, James Edward	

It is advisable to refer to the publisher's version if you intend to cite from the work. 10.12968/bjca.2024.0068

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the http://clok.uclan.ac.uk/policies/

<u>Title</u>: A commentary on the use of accelerometers to improve clinical outcomes in patients with implantable cardioverter-defibrillators

<u>Commentary on:</u> Kolk MZH, Frodi DM, Andersen TO, Langford J, Diederichsen SZ, Svendsen JH, Tan HL, Knops RE, Tjong FVY. Accelerometer-assessed physical behavior and the association with clinical outcomes in implantable cardioverter-defibrillator recipients: A systematic review. Cardiovasc Digit Health J. 2021 Nov 24;3(1):46-55. doi: 10.1016/j.cvdhj.2021.11.006. PMID: 35265934; PMCID: PMC8890329.

Abstract

The Implantable Cardioverter-Defibrillator (ICD) is pivotal in preventing sudden cardiac death, monitoring cardiac activity, and restoring normal sinus rhythm with electrical shocks. With sudden cardiac death accounting for half of cardiac-related fatalities, ICD significantly reduces mortality in high-risk patients. Integrated with accelerometers, ICDs track physical activity (PA), which helps predict cardiac support needs and hospitalisation. Enhanced PA post-ICD implantation is also associated with improved outcomes. Wearable accelerometers hold promise in assessing individual physical behaviour (PB), potentially serving as prognostic predictors of future events. This commentary critically evaluates a recent systematic review by Kolk et al. (2022), which examines the association between PB and key clinical outcomes in patients with ICD or at risk of sudden cardiac death.

Key Points

- Declining physical activity for individuals with implantable cardioverter-defibrillator may be associated with increased risk of mortality and hospitalisation.
- Individuals with implantable cardioverter-defibrillator and low physical activity may be at increased risk of mortality and hospitalisation.
- Future primary research should focus on standardisation of time of outcome reporting and outcomes assessed.

Introduction

Implantable cardioverter-defibrillator (ICD) is a battery-powered device implanted under the skin for continuous cardiac monitoring, detecting any life-threatening arrhythmias, and prompt restoration of normal sinus rhythm via electrical shocks when necessary (Mirowski 1985). Sudden cardiac death accounts for approximately half of all cardiac-related deaths (Wong et al. 2019) therefore, ICD represents a significant preventative measure that reduces cardiac mortality among survivors of sudden cardiac death and patients at high risk for ventricular arrhythmias (Friedman et al. 2022).

Currently, ICDs are designed to include a device-embedded accelerometer, which facilitates in detecting physical activity (PA) changes (Kolk et al. 2022). It has been suggested that increased PA, often achieved through cardiac rehabilitation, is associated with a reduced risk of further cardiac complications and improved outcomes in patients with heart failure (Atwater et al. 2021). Conversely, low PA following implantation of ICD has been shown to increase the risk of mortality, ICD shock, and hospitalisation (Zhao et al. 2017).

Therefore, it is possibly valuable to explore the potential of accelerometer-based methods, which could provide detailed insights into individual physical behaviour (PB) and serve as a more precise indicator of clinical deterioration in cardiac health (Kolk et al. 2022). PB encompasses various dimensions, including PA, sleep patterns, daily activities, postures and sedentary behaviours (Bussmann and van den Berg-Emons 2013). A previous systematic review by Kolk et al. (2022) assessed the association of PB with clinical deterioration leading to implantable cardioverter-defibrillator therapy, heart failure hospitalization, and mortality (Kolk et al. 2022).

Aim of commentary

This commentary aims to critically appraise the methods used within the systematic review by Kolk et al. (2022) and expand upon the findings in the context of clinical practice.

Critical appraisal and methods of the review by Kolk et al. (2022)

Using the Joanna Briggs Institute (2017) critical appraisal tool for systematic reviews, three out of nine criteria were judged to be satisfactory for this review (Aromataris et al. 2015). See Table 1 for full critical appraisal. Firstly, from a methodological standpoint there was a non-specific question proposed. When designing a systematic review, it is helpful to structure your question around one of the standardised systematic review types (Munn et al. 2018). Based upon the reviews question and the results presented in the review it appears that this review aims to assess both prognostic and diagnostic test accuracy simultaneously (Campbell et al. 2015; O'KeeffeGreene and Kearney 2014). Due to this lack of clarity, it makes it difficult to appraise certain methodological processes within the systematic review as most methods are critically appraised in context to the aims of the review. Due to this mixture of methodological designs, it brings into question the use of the Quality in Prognosis Studies tool to assess the risk of bias as this tool is not designed to additionally assess test accuracy studies (Hayden et al. 2013). Consequently, the accuracy outcomes of device-embedded accelerometers, in comparison to validated wearable accelerometers, have not been adequately assessed in the context of potential bias.

The third area of concern was related to the search strategy and the number of databases searches. Within the search strategy there was no clear rationale why the review only searched from 2000 onwards. Where this may be justified based upon certain technologies only being introduced at this time point this was not stipulated within the paper. Due to potential indexing bias in the resources used for searching, it is possible that important papers may have been missed. Previous recommendations for healthcare-related subjects suggest that, at a minimum, Embase, MEDLINE, Web of Science, and Google Scholar should be included as essential databases for comprehensive searches (Bramer et al. 2017).

The fourth area of concern was regarding data extraction, the review did indicate that two reviewers did undertake the data extraction process however there was lack of clarity regarding this process being undertaken independently. The current gold standard for this process is that two reviewers undertake data extraction independently (Higgins et al. 2021). Thus, it is possible that errors may have occurred within these processes.

The final areas of concern pertained to the methods of synthesis and the assessment of publication bias. Concerning the synthesis methods, the effect estimates for directional association were not always presented as ratios. This omission made it challenging to ascertain whether the directional association was representative of all the studies assessed, rather than only those studies that demonstrated that specific direction of association. Additionally, there appeared to be consistency regarding the reporting of hazard ratios for a range of outcomes, but no attempt of meta-analysis was undertaken. With no explanation of any breaches of assumptions needed to carry out this procedure.

Table 1 Critical appraisal using the Joanna Briggs Institute (JBI) critical appraisal checklist for systematic reviews and research syntheses (delete the tool which is not relevant)

JBI critical appraisal checklist items	Responses	
1. Is the review question clearly and explicitly	No - The review presents a broad clinical question	
stated?	that does not specifically focus on diagnostic test	
	accuracy or prognosis studies. With the review stating	
	that the question of interest is "what is the clinical	
	value of PB for identification of clinical deterioration	
2. Were the inclusion criteria appropriate for the	Yes - The inclusion criteria are appropriate. Patients	
review question?	with an Implantable cardioverter-defibrillator,	
	regardless of the use of cardiac resynchronisation	
	therapy or wearable cardioverter-defibrillator, were all	
	included. Patients who are at high risk of developing	
	sudden cardiac death without Implantable	
	cardioverter-defibrillator were included. Eligible	
	accelerometer-based methods included wearable or	
	device-embedded accelerometry, with outcomes of	
	interest being ICD therapy for ventricular	
	arrhythmias, heart failure hospitalization, mortality,	
	functional status (e.g., NYHA class), and quality of	
	life. The included population allows the review to	
	have a detailed comparison between Implantable	
	cardioverter-defibrillator and wearable accelerometer.	
3. Was the search strategy appropriate?	Yes- The review used 2 electronic data bases	
	MEDLINE and EMBASE to identify studies	
	published between Jan 2000 and Aug 2020. Relevant	
	keywords and terms were used to search papers,	
	together with the use of reference lists of relevant	
	papers to identify potentially missed papers.	
4. Were the sources and resources used to search for No - This review only covers 2 electronic data bas		
studies adequate?	This review should ideally cover multiple electronic	
	databases, as well as trial registries to minimise	
	publication bias.	

5. Were the criteria for appraising studies	No - QUIPS (Quality in Prognosis Studies) tool was
appropriate?	used in the review to appraise the included studies.
	However as indicated in the results three studies
	explored validation of wearable and device-embedded
	accelerometry based upon the studies the validation
	studies should be assessed accordingly.
6. Was critical appraisal conducted by two or more	Yes - The review stated that 2 independent reviewers
reviewers independently?	conducted the appraisal. If there were any
	disagreement, a third reviewer will be consulted.
7. Were there methods to minimize errors in data	No - Data extraction was completed by two reviewers;
extraction?	however, it was not specifically stated whether the
	screening process was undertaken independently.
8. Were the methods used to combine studies	No - Within the method section there is no specific
appropriate?	indication of exactly how the descriptive analysis was
	undertaken. It appears that a vote counting method has
	been used however study direction was not presented
	as a ratio making it difficult to assess the degree of
	association for all outcomes. Similarly, within the
	results there appears to be consistency in regard to
	hazard ratios presented within the studies although
	there was no explanation why a meta-analysis was
	performed.
9. Was the likelihood of publication bias assessed?	No - There was no specific reference used to assess
	publication bias.
· · · · · · · · · · · · · · · · · · ·	·

Results of the review by Kolk, et al. (2022)

After duplicate removal 4222 studies were identified of which after screening 52 studies were included in the systematic review. Of these studies, 22 were identified as having a low risk of bias, 19 as having a moderate risk, and 8 as having a high risk of bias.

Among the studies which measured device-embedded accelerometery there was an association between low PA following device implantation and increased risk of mortality (5 studies, n = 101,617,12-31 months), hospitalization (3 studies, n = 1,715,15-36 months), combined hospitalization or mortality (3 studies, n = 1,715,15-36 months) atrial arrhythmias (1 study, n = 770,25 months) and ICD shock (1 study, n = 4,057,1 months). A similar association was observed between low PA, measured using wearable accelerometery, and an increased risk of hospitalization and mortality (3 studies, n = 286,12-36 months).

Similarly declining PA was associated with increased risk of mortality (4 studies, n = 126,234,26-28 months), hospitalization (5 studies, n = 3,522,11.7-17 months), combined hospitalization or mortality (3 studies, n = 1,715,15-36 months) atrial arrhythmias (1 study, n = 770,25 months) and ICD shock (1 study, n = 4,057,1 months).

In the studies that utilized device-embedded accelerometery, several factors were identified as reducing PA. These included seasonal variation (1 study, n = 102, 12 months) and pandemic lockdown (1 study, n = 24, 80 days), both of which were associated with decreased PA. Additionally, implantable cardioverter-defibrillator placement (2 studies, n = 2,944, 12–22 months) and onset of atrial fibrillation (1 study, n = 266, 51.6 months) were associated with a decreased risk of PA reduction. In the studies that utilized wearable accelerometer, several factors were identified as associated with PA. Ischaemic heart failure patients had a lower mean activity level compared to healthy adults (1 study, n = 102, n =

Unfortunately, other outcomes related to wearable devices were reported, but there were significant inconsistencies and a lack of data concerning important variables. For instance, although it was mentioned that ischemic heart failure patients spent more time engaging in vigorous activities, there was no clarification regarding what this was compared to or the specific period during which this observation was made. Similarly, it was noted that sleep behaviour and physical activity were linked to patient-reported physical function, quality of life, and cognitive function. However, details such as the number of studies and participants for each specific association, the observation period, and the type of sleep behaviour observed remained unclear.

Commentary

Overall, it was deemed that this systematic review is likely to provide an accurate and comprehensive summary of the results of the available studies that address the question of interest. However, more research needs to be done to provide a more detailed and holistic review of this subject.

This review highlights the promising potential of using accelerometer-based methods, (implantable) in identifying the potential risks of patients with an ICD. Based on the current evidence in the review, there is not enough evidence to indicate that accelerometers should be used as a stand-alone detection method for increased risk of negative clinical outcomes (e.g., mortality and heart failure). However, it can possibly act as a moderating factor in a multifactorial assessment in identifying clinical deterioration. It is challenging to determine the precise impact that low or declining PAwould have within a predictive model based on this systematic review. As no single estimate of association was generated within the review for any outcome. Therefore, to generate an accurate weighted model, a weighted assessment of association (meta-analysis) for both low and declining PA across the outcomes assessed in this review is required. The individual estimates could then be incorporated into multifactorial predictive models, alongside other key prognostic factors identified in previous systematic reviews. Previous studies in this area have identified that other such factors such as anxiety and depression (Lindekilde et al. 2022), dialysis, chronic renal disease, cancer, advanced age (Alhakak et al. 2022) and ICD shocks (Qian et al. 2016) are also associated with increased risk of mortality. Accelerometer data on PA, if available, could enhance clinical risk stratification by serving as an additional indicator of potential clinical deterioration. This data can be combined with other key risk factors, such as anxiety, depression, and comorbidities, to provide a more comprehensive assessment of a patient's overall risk profile. Due to the significant inconsistency across studies and insufficient reporting within the review, it is challenging to draw any clear recommendations or conclusions regarding the findings on wearable accelerometry based upon this review.

Primary research in this area should aim to standardize both the exposed and unexposed groups, as well as establish consistent outcomes measured at multiple time points. Given the perceived but untested heterogeneity identified in this review, further investigation into potential moderating factors is recommended. Future systematic reviews in this area should aim to focus on a single type of study—either aetiology or test accuracy. While combining these two assessments in one review is possible, it requires two separate methodologies for critical appraisal and evidence synthesis. Separating them may also enhance the search strategy by allowing for more specific term sets tailored to the review being conducted. Future systematic reviews in this area should aim for a more comprehensive search strategy, as this review's focus on only two databases may have led to the omission of relevant studies. To

improve data synthesis, future studies should attempt a meta-analysis when there is sufficient data commonality. If the data proves too heterogeneous for meta-analysis, alternative methods such as vote counting based on the direction of association, reporting medians and quartile ranges, or using visualization techniques like Harvest Plots, Albatross Plots, or Strip Charts should be employed. Given the methodological limitations of these approaches, findings should be interpreted with caution unless more robust synthesis methods are feasible. Additionally, future research must emphasize the complete and consistent reporting of all key variables to allow for more accurate interpretation of the results.

CPD reflective questions

- 1. What are the key limitations of the systematic review presented in this commentary?
- 2. What advice can be given on lifestyle modifications and adherence to cardiac rehabilitation post-surgery to improve prognosis?
- 3. What are the other moderating factors of sudden cardiac death that should be considered in future research?

Funding statement (*must be included in the published article)

This research was partly-funded by the National Institute for Health and Care Research Applied Research Collaboration North West Coast (NIHR ARC NWC). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

References

- Alhakak A, Østergaard L, Butt JH, Vinther M, Philbert BT, Jacobsen PK, Yafasova A, Torp-Pedersen C, Køber L, Fosbøl EL et al. 2022. Cause-specific death and risk factors of 1-year mortality after implantable cardioverter-defibrillator implantation: A nationwide study. Eur Heart J Qual Care Clin Outcomes. 8(1):39-49.
- Aromataris E, Fernandez R, Godfrey C, Holly C, Kahlil H, Tungpunkom P. 2015. Summarizing systematic reviews: Methodological development, conduct and reporting of an umbrella review approach. Int J Evid Based Healthc. 13(3):132-140.
- Atwater BD, Li Z, Pritchard J, Greiner MA, Nabutovsky Y, Hammill BG. 2021. Early increased physical activity, cardiac rehabilitation, and survival after implantable cardioverter-defibrillator implantation. Circ Cardiovasc Qual Outcomes. 14(8):e007580.
- Bramer WM, Rethlefsen ML, Kleijnen J, Franco OH. 2017. Optimal database combinations for literature searches in systematic reviews: A prospective exploratory study. Systematic Reviews. 6(1):245.
- Bussmann JB, van den Berg-Emons RJ. 2013. To total amount of activity..... And beyond: Perspectives on measuring physical behavior. Front Psychol. 4:463.
- Campbell JM, Klugar M, Ding S, Carmody DP, Hakonsen SJ, Jadotte YT, White S, Munn Z. 2015. Diagnostic test accuracy: Methods for systematic review and meta-analysis. Int J Evid Based Healthc. 13(3):154-162.
- Friedman P, Murgatroyd F, Boersma LVA, Manlucu J, O'Donnell D, Knight BP, Clémenty N, Leclercq C, Amin A, Merkely BP et al. 2022. Efficacy and safety of an extravascular implantable cardioverter-defibrillator. N Engl J Med. 387(14):1292-1302.
- Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. 2013. Assessing bias in studies of prognostic factors. Ann Intern Med. 158(4):280-286.
- Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA. 2021. Cochrane handbook for systematic reviews of interventions. Hoboken, NJ: Wiley-Blackwell.
- Kolk MZH, Frodi DM, Andersen TO, Langford J, Diederichsen SZ, Svendsen JH, Tan HL, Knops RE, Tjong FVY. 2022. Accelerometer-assessed physical behavior and the association with clinical outcomes in implantable cardioverter-defibrillator recipients: A systematic review. Cardiovasc Digit Health J. 3(1):46-55.
- Lindekilde N, Skov O, Skovbakke SJ, Johansen JB, Nielsen JC, Pedersen SS. 2022. Anxiety and depression as risk factors for icd shocks and mortality in patients with an implantable cardioverter defibrillator a systematic review. Gen Hosp Psychiatry. 78:96-107.
- Mirowski M. 1985. The automatic implantable cardioverter-defibrillator: An overview. Journal of the American College of Cardiology. 6(2):461-466.

- Munn Z, Stern C, Aromataris E, Lockwood C, Jordan Z. 2018. What kind of systematic review should i conduct? A proposed typology and guidance for systematic reviewers in the medical and health sciences. BMC Medical Research Methodology. 18(1):5.
- O'Keeffe LM, Greene RA, Kearney PM. 2014. The effect of moderate gestational alcohol consumption during pregnancy on speech and language outcomes in children: A systematic review. Systematic Reviews. 3(1):1.
- Qian Z, Zhang Z, Guo J, Wang Y, Hou X, Feng G, Zou J. 2016. Association of implantable cardioverter defibrillator therapy with all-cause mortality-a systematic review and meta-analysis. Pacing Clin Electrophysiol. 39(1):81-88.
- Wong CX, Brown A, Lau DH, Chugh SS, Albert CM, Kalman JM, Sanders P. 2019. Epidemiology of sudden cardiac death: Global and regional perspectives. Heart Lung Circ. 28(1):6-14.
- Zhao S, Chen K, Su Y, Hua W, Chen S, Liang Z, Xu W, Dai Y, Liu Z, Fan X et al. 2017. Association between patient activity and long-term cardiac death in patients with implantable cardioverter-defibrillators and cardiac resynchronization therapy defibrillators. Eur J Prev Cardiol. 24(7):760-767.