

## Research paper

# Impact of tillage practices and soil texture on soil health and earthworms in the Pannonian region: A comparative study from Austria and Hungary

Barbara Simon<sup>a</sup>, Igor Dekemati<sup>b</sup>, Hanaa T.M. Ibrahim<sup>a</sup>, Maxwell M. Modiba<sup>a,b</sup>, Márta Birkás<sup>b</sup>, János Grósz<sup>c</sup>, Martin Kulhanek<sup>d</sup>, Reinhard W. Neugschwandtner<sup>e</sup>, Anna Hofer<sup>e</sup>, Viola Wagner<sup>e</sup>, Marion Windisch<sup>e</sup>, Karin Hage-Ahmed<sup>f</sup>, Kevin R. Butt<sup>g</sup>, Pia Euteneuer<sup>h,\*</sup>

<sup>a</sup> Hungarian University of Agriculture and Life Sciences, Department of Soil Science, Institute of Environmental Sciences, Gödöllő, Hungary

<sup>b</sup> Hungarian University of Agriculture and Life Sciences, Department of Agronomy, Institute of Crop Production Sciences, Gödöllő, Hungary

<sup>c</sup> Hungarian University of Agriculture and Life Sciences, Department of Water Management and Climate Adaptation, Institute of Environmental Sciences, Gödöllő, Hungary

<sup>d</sup> Czech University of Life Sciences in Prague, Department of Agroenvironmental Chemistry and Plant Nutrition, Faculty of Agrobiolgy, Food and Natural Resources, Czech Republic

<sup>e</sup> University of Natural Resources and Life Sciences, Vienna, Department of Crop Sciences, Institute of Agronomy, Tulln, Austria

<sup>f</sup> University of Natural Resources and Life Sciences, Vienna, Department of Crop Sciences, Institute of Plant Protection, Tulln, Austria

<sup>g</sup> University of Central Lancashire, Ecological Engineering, Preston, United Kingdom

<sup>h</sup> University of Natural Resources and Life Sciences, Vienna, Department of Crop Sciences, Experimental farm, Gross-Enzersdorf, Austria

## ARTICLE INFO

## Keywords:

Conventional tillage

Conservation tillage

Clay content

Soil pH value

Arbuscular mycorrhizal fungi

## ABSTRACT

Soil tillage has major impacts on physical, chemical and biological soil parameters. Two long-term soil tillage experiments in North-East Austria (AT) and Central-North Hungary (HU), both located in the Pannonian region, were studied in 2020 and in 2021. The three soil tillage systems comprised mouldboard ploughing (P), shallow cultivation (C), and no-till (NT) in a completely randomised block design with four blocks. Major differences between AT and HU concerned clay content and pH value (clay: 20 %; 36 % and pH: 7.7; 4.5, respectively). This affected most parameters such as dissolved organic carbon, soil aggregate stability, bulk density, earthworm abundance, and biomass and Shannon index. These parameters decreased with soil tillage intensification at both sites. In addition, for all mentioned parameters, C and P were similar in AT, while in HU this was the case for NT and C. Additionally, epi-aneic *Lumbricus terrestris* was only found in AT, while endogeic *Aporrectodea georgii* was only present in HU. Arbuscular mycorrhizal root colonization was not responsive to different tillage practices when sampled in August in AT. In conclusion, reduced soil tillage such as C and NT can show similar affects towards soil health, but site-specific properties such as soil texture need to be considered for a final evaluation.

## 1. Introduction

Soil tillage is an important management practice in agriculture with a high impact on the soil physical (Dekemati et al., 2019; Euteneuer et al., 2024), biological (Dekemati et al., 2020, 2021; Euteneuer and Butt, 2025), and chemical properties (Neugschwandtner et al., 2020; Weidhuner et al., 2021). Soil tillage is practised for the incorporation of plant residues, seedbed preparation, weed and pest control (Gajri et al., 2002; Mairhofer et al., 2019), but there are several disadvantages of soil

tillage depending on the management practices. Ploughing (P) can cause intensive disturbances by inverting the top 30 cm of soil (Obour et al., 2017), and the resulting bare soil surface (Jug et al., 2019) is exposed to wind erosion, decreased soil aggregate stability (SAS) (Klik and Rosner, 2020), and loss of soil biodiversity (Briones and Schmidt, 2017). As opposed to ploughing, conservation agriculture is based on complete omission (no-till) or reduced tillage practices (Busari et al., 2015). No-till (NT) soils are more resilient to abiotic impacts and usually have higher water retention (Liebhard et al., 2022), soil carbon storage

**Abbreviations:** AMF, arbuscular mycorrhizal fungi; AT, Austria; BD, bulk density; DOC, dissolved organic carbon; EEG, easily extractable glomalin; HU, Hungary; NMDS, Non-metric multidimensional scaling; NT, no-till; P, ploughing; PCA, Principal component analysis; SAS, soil aggregate stability; C, shallow cultivation; SOC, soil organic carbon; SPR, soil penetration resistance.

\* Corresponding author.

E-mail address: [pia.euteneuer@boku.ac.at](mailto:pia.euteneuer@boku.ac.at) (P. Euteneuer).

<https://doi.org/10.1016/j.apsoil.2025.105863>

Received 12 July 2024; Received in revised form 29 December 2024; Accepted 1 January 2025

Available online 5 January 2025

0929-1393/© 2025 Published by Elsevier B.V.

(Rosinger et al., 2022; Sae-Tun et al., 2022), more stable soil aggregates (Sae-Tun et al., 2022) and earthworm abundance (Dekemati et al., 2019; Euteneuer et al., 2024). Moreover, shallow cultivation (C) represents a form of conservation agriculture and is considered to have an intermediate impact on soil physics and biodiversity compared to NT and P (Klik and Rosner, 2020; Liebhard et al., 2022; Sae-Tun et al., 2022).

Soil organisms, such as earthworms and arbuscular mycorrhiza fungi (AMF) are also affected by soil tillage (Jansa et al., 2002; Säle et al., 2015; Briones and Schmidt, 2017; Rosner et al., 2018). Earthworms and AMF are important for agricultural soil functionality, especially in the development of soil structure (Jongmans et al., 2003), soil porosity (Pères et al., 2010), and changes in soil organic carbon (SOC) (Sae-Tun et al., 2022). AMF contribute to soil carbon sequestration and soil aggregation via the release of a soil glycoprotein initially termed 'glomalin' during AMF hyphae turnover (Wright and Upadhyaya, 1998; Rillig et al., 2001; Driver et al., 2005; Thomopoulos et al., 2023). Glomalin or glomalin-related soil protein (Irving et al., 2021) contains >85 % sugars and thus provides a strong connection between mineral and organic soil particles (Gunina and Kuzyakov, 2015). Furthermore, it enhances carbon sequestration due to its recalcitrant property (Singh et al., 2022). In addition, dissolved organic carbon (DOC) comprises photosynthetically-derived carbon such as mucilage and exudates of organic compounds (Kalbitz et al., 2000) and can be metabolized by soil bacteria and fungi into more complex carbon compounds and lead to higher soil aggregate stabilization (SAS) (van Groenigen et al., 2010; Sokol et al., 2019; Sae-Tun et al., 2022). In addition, earthworms also have an important role in SAS through casting and burrowing (Lehmann et al., 2017; Euteneuer et al., 2024). Hence, SAS governs ecosystem services such as SOC sequestration (Six et al., 2000, 2004; Klik and Rosner, 2020), prevents soil erosion (Klik and Rosner, 2020), allows gas and water fluxes (Amézqueta, 1999; Finn et al., 2017) and thus supports soil fertility (Blanchart et al., 2009; Arai et al., 2018). High-intensity soil tillage affects SAS negatively, while soil biological activity has a positive effect (Lehmann et al., 2017; Sae-Tun et al., 2022).

The current study aimed to investigate the effects of tillage (NT; C; P) on parameters under similar climatic conditions with calcareous or endocalcic Chernozem soil with respect to i) physical (soil penetration resistance; bulk density; SAS); ii) chemical (pH, DOC, glomalin), and iii) biological properties (earthworm abundance, biomass, community composition; AMF root colonization rates) in two long-term tillage experiments in Austria and Hungary. It was hypothesized that the effects of soil tillage intensity on soil properties and biological communities would vary between Chernozems from Austria and Hungary mostly due to significant differences in soil pH and texture. Specifically, we hypothesized that regional differences in soil properties would interact with tillage practices to produce distinctive impacts on soil health, with these effects being more pronounced in clay-rich, acidic soils in Hungary compared to the calcareous soils in Austria. The interaction between tillage practices and site-specific conditions (e.g., soil pH and texture) is expected to influence soil physical properties, nutrient availability, and biological parameters differently across the Pannonian region.

## 2. Materials and methods

### 2.1. Study sites and years

The experiment in Austria (AT) was instigated in 1996 at the Experimental farm of the University of Natural Resources and Life Sciences, Vienna, in Raasdorf (48°14'N, 16°33'E; 153 m a.s.l.). The experimental field site is located to the east of Vienna (Lower Austria) on the edge of the Marchfeld plain in the north-western part of the Pannonian Basin. The silt loam soil is classified as a calcareous Chernozem of alluvial origin (WRB, 2014). The experiment is conducted as a complete random block design with four replications of plot size 24 × 40 m and different tillage systems, P (30 cm depth), C (10 cm depth) and NT (0 cm). The cropping sequence in AT from 2018 to 2021 was sorghum (*Sorghum*

*bicolor* Mönch), winter wheat (*Triticum aestivum* L.), maize (*Zea mays* L.) (study year 2020), winter wheat (study year 2021).

The experiment in Hungary (HU) is located at the Józsefmajor Experimental and Training farm of the Hungarian University of Agricultural and Life Sciences (47°41'N, 19°36'E; 110 m a.s.l.), in Pest County, Central Hungary. The long-term tillage experiment was set up in 2002 with three tillage systems, P (30 cm depth), C (18 cm depth) and NT (0 cm). The clay loam soil is classified as an endocalcic Chernozem (WRB, 2014). The experiment is arranged in a complete random block design with four replicates and plot size of 13 × 180 m. The cropping sequence between 2018 and 2021 in HU was soybean (*Glycine max* Merr.), winter wheat, winter oat (*Avena sativa* L.) (study year 2020), sunflower (*Helianthus annuus* L.) (study year 2021).

Both sites had a similar continental climate Dfb (1961–1990) in the Köppen-Geiger climate classification (Kottek et al., 2006) with a shift to Csa (1991–2020) for AT and Dfa for HU and an upcoming transition for HU to Csa (2041–2071; Beck et al., 2023). The long-term mean annual temperature for AT is 11.2 °C and 10.3 °C for HU, and the long-term annual precipitation is 560 mm for both countries. For details of soil properties, weather data and time of sampling, see Table 1.

### 2.2. Soil chemical analyses

Soil samples for SOC measurements in HU were taken from depths of 0–10 and 25–30 cm, from four random locations within each plot in autumn 2020. The SOC content of the HU samples was determined by wet oxidation using the Walkley (1947) method. SOC data and soil texture of AT were taken from Liebhard et al. (2022) derived from 2020 in the same plots as the current study. SOC was sampled at 0–10 and 0–30 cm and analysed according to Austrian Standards with dry combustion (ÖNORM L 1080, 1999) and hence, results are not directly comparable with SOC from HU. Therefore, parameters were not statistically analysed but are provided to complete site soil parameters.

For  $\text{pH}_{\text{CaCl}_2}$  measurement, the 0.01 mol L<sup>-1</sup> CaCl<sub>2</sub> extracts were performed in the ratio 1:10 w v<sup>-1</sup> (4 g of fresh soil (≤ 5 mm) 40 ml<sup>-1</sup> of extract) (Houba et al., 2000). After 2 h of shaking, the pH value was measured. The suspension was subsequently centrifuged at 5000g for 5 min. The content of DOC was measured using SKALAR SAN<sup>PLUS</sup> SYSTEM (Netherlands) and was recalculated on soil dry mass obtained from mass difference after 105 °C for 24 h. Glomalin-related soil protein or easily extractable glomalin (EEG) was extracted after Wright and Upadhyaya

**Table 1**

Site parameters of soil tillage trials in Austria and Hungary with treatments no-till (NT), cultivator (C) and plough (P). Current precipitation and average temperature during growth period in 2020 and 2021, soil organic carbon (SOC) and sampling times of physical soil parameters such as soil penetration resistance (SPR), bulk density (BD), soil aggregate stability (SAS); soil chemical parameters such as pH, dissolved organic carbon (DOC), easily extractable glomalin (EEG) and soil biological parameters such as earthworm sampling and plant roots colonised by arbuscular mycorrhiza fungi (AMF). Note that SOC in AT is dry combustion and in HU it is based on wet oxidation.

Parameter	Austria		Hungary	
	2020	2021	2020	2021
Sand/clay/silt (%)	27/20/53		37/36/27	
SOC (10 cm) NT/C/P	2.7/2.5/2.3		2.3/2.1/1.8	
SOC (30 cm) NT/C/P	2.5/2.4/2.3		1.7/1.9/1.7	
Current precipitation (mm)	496	268	359	302
SPR	May	May	Mar	Mar
BD	May	May	Mar	Mar
SAS	Nov	May	Mar	May
$\text{pH}_{\text{CaCl}_2}$	Nov	Jul	Sep	Jul
EEG	Nov	—	—	Sep
DOC	Nov	—	—	Sep
SOC	Jun	—	Sep	—
Earthworm	May & Oct	Mar	Mar & Sep	Mar & Nov
AMF	Aug	—	—	—

(1998) with 20 mmol L<sup>-1</sup> sodium citrate at pH<sub>CaCl2</sub> 7.0 for 30 min at 121 °C. The air-dried soil ( $\leq 2$  mm) solution<sup>-1</sup> ratio was 1:8. After extraction, samples were centrifuged at 5000g for 5 min. The EEG content was measured colorimetrically using an extract of dye brilliant blue reagent and Bovine serum albumin as a standard (Protein assay kit II, Bio-Rad, USA). For the analysis, 10  $\mu$ l of the extract was dispensed onto a microplate, followed by the addition of 200  $\mu$ l of dye reagent (diluted at a 1:4 ratio). Subsequently, the microplate was horizontally shaken at 593 rpm for 30 s and then measured after a 5-minute stabilization period at a wavelength of 590 nm (Spark, Tecan Ltd., Zurich, Switzerland).

### 2.3. Soil physical analyses

SAS was measured with the smartphone application MOULDER Version 2.0 (formerly: SLAKES) (Fajardo et al., 2016; Fajardo and McBratney, 2019) and according to the analytical protocol established by Flynn et al. (2020) and utilising an iPhone 8 (Apple Inc. Cupertino, California, USA) in AT and Huawei P30 lite MAR-LX 1A (Huawei Technologies Co., Ltd., Bantian, Longgang, Shenzhen, China) in HU. In summary, air-dried soil samples were sieved to a particle size range of 2–5 mm, with three soil aggregates per sample and five subsamples per plot analysed (Euteneuer et al., 2024). The initial step in the protocol involves capturing a reference photo of the dry soil aggregates positioned in a dry Petri dish against a high-contrast background. Subsequently, the aggregates are transferred to a Petri dish containing deionized water, ensuring that they are placed in the same orientation as in the reference image. The application then measures the expansion of the dispersed soil area, starting from the reference image and determining the final area after a 10-minute interval. Upon completion of the ten-minute period, the  $\alpha$ -coefficient is displayed on the screen. The  $\alpha$ -coefficient represents the maximum predicted dispersion of a soil aggregate and is derived from the Slaking index, which is fitted to the Gompertz function (Gompertz, 1833). For comprehensive details regarding the  $\alpha$ -coefficient, the Slaking index, and the methodology, refer to Fajardo et al. (2016). A lower  $\alpha$ -coefficient indicates more stable soil, with zero being the most stable value. For comparison of MOULDER to more established methods, such as Cornell Rainfall Simulator (Moebius-Clune et al., 2016), wet sieve procedure (Kemper and Roseau, 1986; Nimmo and Perkins, 2002) or water stable aggregate mean weight diameter (Franzuebbers et al., 2000) related to earthworm processed soil or soil health, refer to Euteneuer et al. (2024) and Rieke et al. (2022).

Soil bulk density (BD) samples were taken in the topsoil (0–5 cm) with an undisturbed soil sampler, using 100 cm<sup>3</sup> rings in four subplots. Soil penetration resistance (SPR) was measured at both sites by an electronic penetrometer (Penetrologger, Royal Eijkelpark, Giesbeck, Netherlands). The measurements were carried out at 20 random points plot<sup>-1</sup>. SPR values were measured by 1 cm, 1 N accuracy, between 0 and 40 cm, with a penetration speed of 2 cm s<sup>-1</sup>, with a 1 cm<sup>2</sup> cone.

### 2.4. Soil biological analyses

Earthworms were sampled by hand-sorting 25 × 25 × 30 cm soil blocks, in four randomly chosen places of each treatment at both sites (ISO 23611-1, 2018). Parameters measured were earthworm abundance (individuals m<sup>-2</sup>), earthworm biomass (g m<sup>-2</sup>), juvenile:adult ratio, juvenile:adult biomass ratio, average biomass of endogeic adult earthworm (g individual<sup>-1</sup>) and species composition according to Christian and Zicsi (1999) and Csuzdi and Zicsi (2003) for AT and HU, respectively. In addition to hand-sorting, a common practice is the use of a vermifuge, such as a mustard suspension (5 g L<sup>-1</sup> mustard powder), allyl isothiocyanate or formaldehyde (formalin) to expel earthworms from soil, particularly deep burrowing earthworms such as *Lumbricus terrestris* (Linnaeus, 1758) (Butt and Grigoropoulou, 2010). The disadvantage of a vermifuge for silty soil such as in AT is the increased infiltration rate

(1–2 h; Euteneuer unpublished). Silty Chernozem soil is highly sensitive to soil tillage, which can block infiltration of any liquid (Weninger et al., 2019) and was therefore not applicable. As an alternative for an expellant, we counted middens of *L. terrestris*. To determine the middens, wheat straw (178 g m<sup>-2</sup>) was added onto an area of 7 m<sup>2</sup> in May and June 2020 to allow *L. terrestris* to build middens (Euteneuer et al., 2024). These middens were then counted in November 2020. Normally, there is one *L. terrestris* burrow under each midden, but sometimes a burrow can have two openings to the soil surface, which can appear as two individual middens and could lead to an overestimation of *L. terrestris* abundance (Butt and Grigoropoulou, 2010; Grigoropoulou and Butt, 2010). To verify our approach, we checked ten randomly selected middens plot<sup>-1</sup> for occupancy using mustard suspension (Stroud et al., 2016). Earthworms were taken to the laboratory and biomass were recorded (g individual<sup>-1</sup>), but not added to the earthworms from hand-sorting. The proportion of *L. terrestris* (total; adult; juvenile) expelled by mustard suspension from the middens (occupancy rate), was calculated by number of middens occupied divided by ten middens plot<sup>-1</sup>. Subsequently, recorded middens m<sup>-2</sup> were corrected by total occupancy rate of middens.

For AMF assessment, 5 randomly chosen plants plot<sup>-1</sup> were sampled in late August 2020, but only at the AT site due to COVID-19 travelling restrictions. Roots were stained according to the method of Vierheilig et al. (1998). Briefly, maize roots were cleared in 10 % potassium hydroxide (KOH) for 6 min at 90 °C and then roots were stained in 5 % ink-vinegar solution for 6 min at 90 °C. The stained roots were then stored in 30 % ethanol at 4 °C. For each sample, 30 root pieces were mounted on a microscope slide and AMF colonization was determined according to the modified method of Trouvelot et al. (1986) using the INOQ calculator Advanced (Mercy, 2017). Parameters measured were mycorrhizal frequency (F %), mycorrhizal intensity (M %), arbuscule abundance (A %), vesicle abundance (V %) and hyphal abundance (H %). Details of sampling time at both sites are provided in Table 1.

### 2.5. Statistical analyses

Two-way linear mixed model (2-way LMM) with the fixed effect site (2 levels; AT, HU) and tillage (3 levels; P, C, NT) were used to analyse parameters BD, SAS, SPR, pH, DOC, EEG, earthworm abundance, earthworm biomass, juvenile:adult ratio, juvenile:adult biomass ratio, average biomass of endogeic adult earthworms and Shannon index. Shannon index was obtained from package ‘vegan’ (Oksanen et al., 2020) function ‘diversity’. Middens were only found in AT and parameters occupancy rate, numbers of middens, proportion of adult and immature *L. terrestris* and their biomasses were analysed with fixed effect tillage and random effect replicate using a one-way LMM (1-way LMM).

For 2-way LMM replicates, years (2 levels; 2020, 2021), sites and sampling dates (according to Table 1) were set random. All LMMs applied function ‘lmer’ (‘lme4’ package; Bates, 2015) in RStudio 6.1.524 (Posit team, 2023) using R 4.3.1 (R Core Team, 2023) with compound symmetry as a variance-covariance structure for repeated measurements and fitted for residual maximum likelihood (REML) method was used. Compound symmetry was selected as both long-term trials have existed for >20 years and variance in data are considered as momentary effects due to weather conditions as soil tillage systems are believed to have reached some form of stabilization after two decades. Function ‘Anova’ was applied for the analyses of variance with Wald-type *F*-tests and the Satterthwaite’s method for denominator degrees of freedom and type III hypotheses. Tukey post-hoc test with function ‘emmeans’ (package ‘emmeans’; Lenth, 2022) was applied in multiple mean comparisons (*P* < 0.05) for factor combinations. All data provided are mean values and standard deviation (mean  $\pm$  SD). Residual distributions were checked visually by frequency of residuals and homogeneity of the variance by residuals against fitted values per model. Parameters that did not meet these assumptions, such as EEG, average biomass of earthworm,

occupancy rate, number of middens, proportion and biomass of adult and immature *L. terrestris*, were square root transformed. Total earthworm biomass, juvenile:adult ratio, juvenile:adult biomass ratio were log transformed.

Non-metric multidimensional scaling (NMDS) was applied for ordination of rank orders for soil parameters (Table 1) and mean earthworm species abundance for both sites (Paliy and Shankar, 2016). For NMDS, package 'vegan' and function 'metaMDS' with Bray-Curtis distances was used and was solved with  $k = 2$  and a stress score of 0.141 after an interaction of 20 tries (Kenkel and Orloci, 1986; Clarke, 1993). Plotting of NMDS was done with package 'ggplot2' (Wickham, 2016) and 'score' function to extract the results of vector fitting by function 'envfit' (package 'vegan') with scaling 'species' for earthworm species and 'site' for soil and site parameters.

A Principal Component Analysis (PCA) with AT data was performed on the soil parameters, earthworm data from October 2020 and AMF data from August 2020 to depict the interplay and the contribution of the selected variables. For PCA analysis and visualization the packages 'FactoMineR' (Le et al., 2008) and 'factoextra' (Kassambara and Mundt, 2020) were used.

### 3. Results

#### 3.1. Soil chemistry and physics

The results of the 2-way LMM showed interaction of site  $\times$  soil tillage for pH (Table 2), with a higher pH in AT than HU, and within HU highest for P, whereas pH did not differ between tillage treatments in AT (Fig. 1A). The EEG was only affected by site and had a 25 % lower concentration in AT than HU (Fig. 1B). In addition, DOC differed between sites and was 2.7 times higher in AT than HU and decreased with soil tillage intensity at both sites (AT: NT > C = P > HU: NT = C > P) (Fig. 1C). Soil physical parameters SAS, BD and SPR showed an interaction of site  $\times$  soil tillage (Table 2). Aggregate stability was 2.25 times greater for NT in AT than C and P, whereas in HU C  $\geq$  NT were 1.7 times more stable than P (Fig. 1D). Bulk density followed a similar pattern and decreased with soil tillage intensity and was 1.2 times higher in HU than AT (Fig. 1E). Two-way LMM of SPR showed that resistance across depths was lowest in NT and P in HU, followed by P  $\leq$  C in AT and  $\leq$  C in HU and  $\leq$  NT in AT (Fig. 1F).

**Table 2**

ANOVA results of chemical, physical and biological soil parameters (2-way LMM) with fixed factors site (S; Austria, Hungary) and tillage (T; no-till, cultivation, plough). Easily extractable glomalin (EEG), dissolved organic carbon (DOC), soil aggregate stability (SAS), bulk density (BD), soil penetration resistance (SPR) and earthworm parameters. Degrees of freedom: S = 1, T = 2, S  $\times$  T = 2, N = 4.

Parameter	F-value			P-value		
	S	T	S $\times$ T	S	T	S $\times$ T
pH <sub>CaCl2</sub>	3544	1.61	13.06	<0.001	0.218	<0.001
EEG	59.9	3.24	0.423	<0.001	0.054	0.66
DOC	267	31.2	5.08	<0.001	<0.001	0.013
SAS	0.053	17.9	5.12	0.819	<0.001	0.013
BD	159	18.7	9.89	<0.001	<0.001	<0.001
SPR	0.694	5.79	15.0	0.406	0.004	<0.001
Earthworm abundance	4.22	16.7	8.51	0.046	<0.001	<0.001
Earthworm biomass	14.6	7.64	30.5	0.001	<0.001	<0.001
Adult biomass	2.66	0.22	9.38	0.108	0.801	<0.001
Juvenile:adult ratio	3.57	9.05	6.04	0.065	<0.001	0.004
Juvenile:adult biomass ratio	17.3	6.25	3.53	<0.001	0.004	0.036
Shannon diversity	4.38	3.48	5.05	0.043	0.045	0.013


#### 3.2. Soil biology and biodiversity

All earthworm parameters were affected by site  $\times$  soil tillage, except for adult biomass which was only affected by site (Table 2). In detail, earthworm abundance and biomass were 1.6–1.9 or 4–6 times higher for NT in AT or HU, respectively, and decreased with soil tillage intensity and was lowest in P in HU, while in AT, P was similar to C (Fig. 2A, B). Overall, NT in AT showed the greatest juvenile:adult ratio compared to the remaining treatments (Fig. 2C), but the juvenile:adult biomass ratio for C, P and NT in HU showed similar results to NT in AT (Fig. 2D). In addition, average biomass of an adult earthworm was similar within sites but higher for C in AT than P in HU (Fig. 2F). The Shannon diversity index was highest in NT and C in HU and NT in AT, followed by C and P in AT and lowest in P in HU (Fig. 2E). The earthworm community at the two sites consisted mainly of endogeic earthworms dominated by *Aporrectodea caliginosa* (Savigny, 1826), *A. rosea* (Savigny, 1826), plus *Allolobophora chlorotica* (Savigny, 1826). Differences were found, as *Aporrectodea georgii* (Michaelsen, 1890) was present in HU, but absent in AT, while *L. terrestris* was found in AT, but not in HU (Fig. 3). In addition, the presence of *L. terrestris* at both sites were also assessed by counting middens and corrected by the occupancy rate of middens. No middens were found in HU and the number of middens in AT were 10–12 times higher in NT than C and P (Table 3). Similar was seen with the occupancy rate, which was 30–40 % higher in NT than C and P (Table 3). In addition, the proportion of adults did not differ between the tillage system, but smaller and more immature *L. terrestris* were expelled in NT than C, while biomass of adult was similar between the tillage systems (Table 3). The presence and absence of the earthworm species mainly affected the outcome of the NMDS (Fig. 4A). Thus, a clear separation of soil tillage systems P and NT and sites was observed by vectors EEG ( $R^2 = 0.467$ ), SAS ( $R^2 = 0.294$ ), BD ( $R^2 = 0.539$ ), clay and sand content ( $R^2 = 0.459$ ) at the HU site (Fig. 4B).

The AMF parameters F % ( $F_{1,2} = 0.375$ ,  $P = 0.690$ ), M % ( $F_{1,2} = 1.066$ ,  $P = 0.354$ ) and A % ( $F_{1,2} = 2.312$ ,  $P = 0.113$ ) were not affected by soil management. AMF frequency reached between 98 and 99 % (Table 4). AMF intensities ranged between 29 and 31 %. Arbuscule abundance was in the range of 26 to 29 %. Vesicle abundance ( $F_{1,2} = 5.378$ ,  $P < 0.01$ ) and hyphal abundance ( $F_{1,2} = 7.036$ ,  $P < 0.01$ ) were significantly affected by soil management. Vesicle abundance was lowest in NT (5 %) and highest in the C treatment (9 %). For hyphal abundance, the lowest values (5 %) were seen in the C treatment and highest (9 %) in the NT treatment.

Results of the PCA that depict the interplay between AMF root colonization parameters, earthworm parameters and soil properties are provided in Fig. 5. Parameters identified in a first PCA (data not shown) as less important, such as DOC, soil penetration resistance from 0 to 10 cm, mean soil penetration resistance from 0 to 40 cm and soil pH were excluded from the data analysis. The PCA identified 3 principal components explaining 68.9 % of the total variance. Juvenile earthworm mass (0.920), juvenile earthworm abundance (0.825) followed by the soil parameters bulk density (0.734) and soil organic carbon 0–10 cm (0.637) correlated to a great extent to PC1 explaining 30.2 % of the variance (Fig. 5). Furthermore, soil aggregate stability was negatively correlated ( $-0.639$ ) with PC1. The second PC explained 22.8 % of the variance. The most highly correlated variables were mycorrhizal intensity (M %) (0.874) and arbuscule abundance (A %) (0.822). Additional variables correlated with PC2 were SOC 0–30 cm (0.634) and vesicle abundance (V %) (0.592). The third PC explained 16.0 % of the variance and was excluded from the PCA plot to increase readability. Adult earthworm biomass (0.766) and adult earthworm abundance (0.739) correlated positively with PC3 and hyphal abundance ( $-0.583$ ), and glomalin ( $-0.613$ ) correlated negatively with PC3.

Additionally, individual data points of the respective treatments were included in the PCA biplot (Fig. 5). C and P samples were completely separated from the NT samples. This separation mainly depended on parameters such as BD, SAS and SOC and earthworms and



**Fig. 1.** Soil chemical parameters  $\text{pH}_{\text{CaCl}_2}$  (A), Easily extractable glomalin (B), dissolved organic carbon (C), and soil physical parameters soil aggregate stability ( $\alpha$ -coefficient; D), bulk density (E) and mean soil penetrations resistance of 0–40 cm (F) in Austria or Hungary of three tillage treatments in 2020 and 2021. Tillage treatments having no letter in common are significantly different by pairwise comparison (2-way LMM, Tukey;  $P < 0.05$ ). Mean + SD,  $N = 4$ . Note: More stable soil aggregates have a lower  $\alpha$ -coefficient.

was associated with NT. PCA (Fig. 5) showed that PC1 accounted for 35.9 % of variance and was associated with earthworm parameters, EEG, SOC and soil physical parameters such as BD and SAS. Parameters of AMF were linked to PC2 and explained 25.4 % of variance. Soil tillage system ellipses overlapped slightly in P and C and all AMF parameters, except hyphal abundance, were related to C. Hyphal abundance was not linked to any soil tillage system and negatively correlated to remaining AMF parameters. By contrast, all earthworm parameters were clearly associated with NT, while SAS decreased with soil tillage intensity and was negatively correlated to earthworm parameters. AMF and earthworm data appeared to be independent at the AT site.

## 4. Discussion

### 4.1. Soil chemistry and physics

Overall, results of soil chemistry and physics were highly affected by soil tillage systems and differences for soil pH and clay or silt content. The calcareous loess parent material in AT was relevant for the high pH value (Bolan et al., 2023) and was not affected by soil tillage treatments as seen in Neugschwandtner et al. (2022) and mainly affected EEG and DOC. EEG was higher in HU than AT and clearly benefited from a lower pH, as seen by Singh et al. (2016), due to a reduced nutrient availability, which can affect AMF growth negatively (Rillig et al., 2001; Wright and

Upadhyaya, 1996). The greater DOC values in AT than HU were also pH-related and decreased with soil tillage intensity at both sites. Sae-Tun et al. (2022) showed that DOC decreased with soil tillage intensification in the order of NT > C > P in a long-term soil tillage trial in the same area of AT, with a similar silty loam Chernozem and soil pH as in the current study. Soil pH can determine the release of DOC, as the availability of dissolved organic substances in soils increases with increasing pH (Kalbitz et al., 2000; Jones and Willett, 2006) and explains the higher concentrations of DOC in AT (pH 7.7) than HU (pH 4.5). In addition, the clay loam texture in HU (36 %; compared with 20 % clay content in AT) may have caused a stronger adsorption of DOC to clay minerals, as suggested by Ussiri and Johnson (2004) and Saïdy et al. (2015).

For soil physical parameters, SAS decreased with soil tillage intensification at both sites. While Flynn et al. (2020) found a clear separation of SAS between soil tillage systems in clayey soils, aggregates in HU showed similar results for NT and C. But when Bagnall and Morgan (2021) compared tillage systems for soils with 27–38 % clay content, they found no differences for SAS between NT and chisel-ploughing (20–25 cm depth), which can be considered as less destructive than P, but with higher intensity than C. Moreover, Bagnall and Morgan (2021) stated that SAS is independent of clay content, but that clayey soils need a higher content of organic carbon to show similar SAS than soils with lower clay content. In addition, Schrader and Zhang (1997) found that soil texture is a predictor for SAS and that soils which are more sensitive




Fig. 2. Earthworm parameters total abundance (A), biomass (B), juvenile:adult ratio (C), juvenile:adult biomass ratio (D), Shannon diversity index (E), and average biomass of endogeic adult earthworm (F) in Austria or Hungary of three tillage treatments in 2020 and 2021. Tillage treatments having no letter in common are significantly different by pairwise comparison, except for adult biomass (F) which was only affected by site (2-way LMM, Tukey;  $P < 0.05$ ). Mean  $\pm$  SD,  $N = 4$ .





Fig. 3. Earthworm community composition of *Aporrectodea caliginosa* (Acal), *A. rosea* (Aros), *A. georgii* (Ageo), *Allolobophora chlorotica* (Achl) and *Lumbricus terrestris* (Lter) in Austria or Hungary of three tillage systems in 2020 and 2021 assessed by hand-sorting. Mean,  $N = 4$ .

Table 3

ANOVA results of rate of occupied burrows of *Lumbricus terrestris* (occupancy rate), number of middens corrected by the occupancy rate (middens), rate of adult and immature *L. terrestris* burrow occupancy and biomass with fixed factor tillage in Austria (2020). Tillage treatments having no letter in common are significantly different by pairwise comparison (1-way LMM, Tukey;  $P < 0.05$ ). Mean  $\pm$  SD,  $N = 4$ , degrees of freedom = 2.

Parameter	No-till	Cultivator	Plough	F-value	P-value
Occupancy rate	0.825 $\pm$ 0.126 <sup>b</sup>	0.525 $\pm$ 0.171 <sup>a</sup>	0.425 $\pm$ 0.096 <sup>a</sup>	14.3	0.005
Middens	28.7 $\pm$ 7.23 <sup>b</sup>	2.85 $\pm$ 2.76 <sup>a</sup>	2.28 $\pm$ 1.42 <sup>a</sup>	56.5	>0.001
Adult rate	0.325 $\pm$ 0.222 <sup>a</sup>	0.4 $\pm$ 0.125 <sup>a</sup>	0.275 $\pm$ 0.096 <sup>a</sup>	1.28	0.344
Immature rate	0.5 $\pm$ 0.141 <sup>b</sup>	0.125 $\pm$ 0.126 <sup>a</sup>	0.15 $\pm$ 0.1 <sup>ab</sup>	7.63	0.011
Biomass adult	3.33 $\pm$ 0.463 <sup>a</sup>	4.22 $\pm$ 0.708 <sup>a</sup>	3.43 $\pm$ 0.468 <sup>a</sup>	2.94	0.104
Biomass immature	1.18 $\pm$ 0.167 <sup>a</sup>	2.74 $\pm$ 0.366 <sup>b</sup>	2.6 $\pm$ 0.358 <sup>b</sup>	215	>0.001

to physical disturbance, such as soils with high silt content (Weninger et al., 2019), show the highest effect on SAS through earthworm casting, while clayey soils have an inverse effect. These results are in line with current findings, when data from AT showed a gradient in SAS from NT



**Fig. 4.** Non-metric multidimensional scaling (NMDS) of earthworm species (A) in Austria and Hungary under three soil tillage treatments from 2020 to 2021. Earthworm species found were *Lumbricus terrestris* (Lter), *Aporrectodea caliginosa* (Acal), *A. rosea* (Aros), *A. georgii* (Ageo) and *Allolobophora chlorotica* (Achl). Site parameters of NMDS (B) include soil aggregate stability (alpha), bulk density (BD), soil organic matter from 0 to 10 cm and 0–30 cm (SOC\_10, SOC\_30 cm; respectively), dissolved organic carbon (DOC), soil texture (Sand, Clay, Silt), pH, easily extractable glomalin (EEG), long-term temperature and precipitation (LT\_temp, LT\_pre; respectively) and annual precipitation (ST\_pre).

**Table 4**


Arbuscular mycorrhiza colonization parameters for maize roots under three tillage treatments at the Austrian site (2020). Tillage treatments having no letter in common are significantly different by pairwise comparison (2-way LMM, Tukey;  $P < 0.05$ ). Mean,  $\pm$ SD,  $N = 4$ . F (% mycorrhizal frequency), M (% mycorrhizal intensity), A (% arbuscule abundance), V (% vesicle abundance), H (% hyphal abundance) in the root system.

Tillage	F (%)	M (%)	A (%)	V (%)	H (%)
No-till	98 $\pm$ 5 a	31 $\pm$ 3 a	27 $\pm$ 5 a	5 $\pm$ 3 <sup>a</sup>	9 $\pm$ 4 <sup>b</sup>
Cultivator	99 $\pm$ 4 a	31 $\pm$ 4 a	29 $\pm$ 5 a	9 $\pm$ 7 <sup>b</sup>	5 $\pm$ 4 <sup>a</sup>
Plough	99 $\pm$ 2 a	29 $\pm$ 4 a	26 $\pm$ 6 a	6 $\pm$ 4 <sup>ab</sup>	7 $\pm$ 5 <sup>ab</sup>

$> C = P$  and with similar results of Euteneuer et al. (2024) from the same area in AT. In addition, BD was increased by  $P \geq C > NT$ , while BD was higher in HU than AT. Klik and Rosner (2020) showed similar with BD increased with reduced soil tillage independently of soil texture within 0–20 cm soil depth in three long-term soil tillage trials in AT. However, SPR differed between soil tillage systems, where NT in AT was higher than P, but NT in HU was similar to P. These results in AT concur with previous publications (e.g., Dekemati et al., 2019).

**4.2. Earthworm community and AMF**

Earthworm communities at both sites were affected by soil tillage systems, showed highest abundance and biomass in NT and decreased with intensity, similar to previous studies of Dekemati et al. (2019) at the HU site. This finding is supported by a meta-analysis of Briones and Schmidt (2017), they reported that anecic and larger-sized species such as *L. terrestris* benefit more from reduced soil tillage than endogeic and smaller-sized earthworms such as *A. chlorotica* and *A. caliginosa*. In the current study, *A. caliginosa* was less sensitive to soil tillage compared to *L. terrestris*, but also increased with reduced soil tillage intensity, but not as much as *L. terrestris* in AT, when 12-times more middens were counted in NT than P (Briones and Schmidt, 2017; Capowiec et al., 2009; Euteneuer et al., 2024). Overall, endogeic earthworms were found in all soil tillage systems except for *A. chlorotica* for P in HU, therefore the Shannon index was higher in NT at both sites and lower for P in HU. Interestingly, *A. georgii* and *L. terrestris* were found mostly in reduced tillage



**Fig. 5.** Principal Component Analysis (PCA) biplot depicting the interplay between arbuscular mycorrhizal root colonization parameters, earthworm parameters, and soil properties at the Austrian site (2020). Ellipses are confidence ellipses based on the 0.95 level. F (mycorrhizal frequency), M (mycorrhizal intensity), A (arbuscule abundance), V (vesicle abundance), H (hyphal abundance) in the root system, BD (bulk density), alpha (soil aggregate stability), EEG (easily extractable glomalin), SOC\_10 (soil organic carbon from 0 to 10 cm), SOC\_30 (soil organic carbon from 0 to 30 cm), ad (adult earthworm abundance (individuals  $m^{-2}$ ), ad\_mass (adult earthworm biomass ( $g\ m^{-2}$ ), juv (juvenile earthworm abundance (individuals  $m^{-2}$ ), juv\_mass (juvenile earthworm biomass ( $g\ m^{-2}$ ).

and seem to have the same demands related to undisturbed soils, but *A. georgii* was absence in AT, while *L. terrestris* was missing in HU. Similar to Briones and Schmidt (2017), we believe that ecological groups or body size are not sufficient to explain the presence or absence in certain soil tillage systems or site. Currently, ecological groups and functional groups are undergoing some revisions and clarifications

(Bottinelli et al., 2020; Bottinelli and Capowicz, 2021; Capowicz et al., 2024). For example, anecic for *L. terrestris* is used by Briones and Schmidt (2017) and Butt et al. (2022) or epi-anecic by Bouché (1972), Bottinelli and Capowicz (2021), and Capowicz et al. (2024). In addition, Capowicz et al. (2024) categorised *L. terrestris* to functional group burrower in relation to burrowing, feeding and casting activities close to the surface or at the surface. *A. caliginosa* is also determined as epi-anecic with functional group burrower by Capowicz et al. (2024) and for ecological group as endogeic by Bouché (1972), Bottinelli and Capowicz (2021) and Capowicz et al. (2024). Perhaps neither ecological groups nor functional groups are sufficient to explain the sensitivity of certain earthworm species to soil tillage systems, but additional information is needed, such as preferred burrowing mode (ingestion or cavity expansion; Arrázola-Vásquez et al., 2022), habitat requirements, survival strategies, soil properties and site history (Briones and Schmidt, 2017) to evaluate the sensitivity, presence or absence of certain earthworm species. As the climatic data at both sites are similar, the NMDS result suggested that differences in earthworm community composition are mainly related to soil properties pH and soil texture, but leaving out the other mentioned factors such as burrowing mode, survival and site history. The pH and clay data dominated the NMDS analysis, but according to Misirlioglu et al. (2016), *L. terrestris* can be found in soils with low pH values (3.5–3.7) and Edwards and Lofty (1975) emphasized that the abundance of the same earthworm species enhanced as the acidic pH values (pH 3.7) increased to neutral (pH 7.0), suggesting that earthworms favour neutral pH ranges to acidic soils. The high clay content in HU might explain why *A. georgii* was only detected at HU site, which is in line with Csuzdi and Zicsi (2003), who reported that *A. georgii* can be found in moist clayey soils in southern Europe. In addition, Zicsi (1994) found *A. georgii* only 8 km from the AT site, in a pasture in the Lobau (an alluvial forest in Donau-Auen National Park) with mostly loamy sand Fluvisol (WRB, 2014; BFW, 2023). The moisture content of the soil might therefore be more important than the clay content, but cannot be addressed in this study as unknown factors may have caused the absence/presence of *A. georgii*.

Apart from earthworms, AMF are an important indicator for soil health, therefore the parameters of AMF colonization and EEG were selected to monitor AMF development. AMF colonization rates did not differ between the treatments at the sampling date in August in AT, at the corn filling stage. As obligate biotrophs, AMF development is strongly dependent on an available host plant and fluctuates over the growing season (Abbott and Robson, 1985; Kabir et al., 1997). In a field study with maize, hyphal density and density of metabolic active hyphae were the lowest in spring before maize was sown and increased thereafter at the 12–14 leaf stage, peaked at the silking stage and decreased again following root senescence (Kabir et al., 1997). In addition, differences between soil management practices also depended on the sampling time. Treatments with P showed diminished AMF parameters compared to C and NT only at the silking stage. In the current study, root samples were taken at the corn filling stage where differences were probably no longer evident. Furthermore, EEG can be used as a proxy for arbuscules and hyphal lengths of AMF, as it correlates closely with phospholipid fatty acids 16:1 $\omega$ 5 (Thomopoulos et al., 2023). EEG did not follow the expectation that soil tillage should influence EEG values. However, by tendency, EEG was lower in the P treatment than in the C and NT treatments as seen by Thomopoulos et al. (2023). The same tendency was seen in HU when the magnitude of the EEG values was in general higher than in AT. In another study investigating the link between EEG and AMF hyphal biomass in different soil aggregate classes, Helgason et al. (2010) could find only a correlation between EEG and AMF biomass in the P treatment, but not in the NT treatment, indicating a potential threshold from where AMF ceased to affect aggregate stability. Although EEG is proposed as a useful proxy for assessing AMF biomass and its potential impact on SAS, its association with soil tillage practices may be complex and dependent on site-specific factors. Holátko et al. (2021) recently reported that any correlation between

AMF and EEG may be indirect or coincidental, due to unknown interplays of the soil food web, turnover rates of glomalin or differences in glomalin productions rates of AMF species.

To date, studies linking AMF and earthworm data are scarce. In a combined principal components analysis, AMF and earthworm parameters at the AT site were not correlated. Earthworm parameters were clearly linked to NT, while AMF parameters were not affected by tillage. As discussed above, results for earthworm parameters are supported by many previous studies, but AMF parameters were also expected to increase in NT and decrease with soil tillage intensity (Kabir et al., 1997; Rosner et al., 2018; Thomopoulos et al., 2023). Non-responsiveness of AMF data might be linked to the sampling date, or a site-specific physiological maximum for root colonization.

Earthworm activity, in addition to AMF hyphal activity/degradation, contributes to SAS (Six et al., 2004; Helgason et al., 2010; Euteneuer et al., 2024). In a recent review, it was highlighted that earthworms ingest AMF and that AMF propagules stay active in the digestive system resulting in the dispersal of AMF propagules and enhancement of root colonization (Pelosi et al., 2024). However, grazing and burrowing activity can simultaneously result in a decrease in hyphal length and disruption of hyphal networks (Meng et al., 2022; Pelosi et al., 2024). As the contributions of AMF and earthworms to SAS follow very distinct pathways, it remains to be determined if their contributions can be linked and may provide synergistic effects. Investigating these potential links may require consideration of the behaviour and niche preferences of different earthworm species, specific soil and site factors, and the overall history of soil management.

#### 4.3. Future perspectives

As a causal relationship between AMF and earthworm activity remains elusive, further studies on different soils and at varying scales could provide more insight. For example, investigating aggregate size classes, i.e., moving from bulk soil to a finer scale, could offer a clearer understanding of potential synergistic effects (Helgason et al., 2010). In addition, microcosm experiments using a range of field soils may serve as the preferred test system in such cases. To enhance clarity, future research should aim to define the nature of the AMF-earthworm relationship, address specific knowledge gaps and guide investigations at different scales and soil types.

Soil tillage showed different impacts on soil and earthworm parameters related to soil properties rather than soil tillage treatments. The higher silt content in AT made the soil particularly more sensitive to soil tillage than in HU. Even reduced tillage showed a similar impact on DOC, SAS, BD, SPR and earthworms compared with conventional tillage, so our hypothesis cannot be fully accepted or rejected. While NT improved soil health parameters such as DOC, SAS and earthworms, reduced soil management depended on soil texture. Further research and reviews on various topics might usefully consider soil texture, when comparing soil tillage systems.

## 5. Conclusion

The current study showed that within the same soil type, site specific soil properties such as pH and soil texture have a high impact on soil chemistry and the earthworm community. In addition, other parameters such as DOC, earthworm abundance and biomass, and soil physical properties were clearly affected by soil tillage. Nevertheless, the extent of the influence of the soil tillage system depended on soil properties. For most parameters, it was seen that C showed some intermediate position between P and NT in HU, while in AT, C was rather similar to P. Hence, soil tillage, and especially reduced tillage such as C, needs site-specific evaluation with respect to soil health.

## CRedit authorship contribution statement

**Barbara Simon:** Writing – original draft, Methodology, Investigation, Conceptualization. **Igor Dekemati:** Writing – original draft, Methodology, Data curation. **Hanaa T.M. Ibrahim:** Writing – original draft, Investigation. **Maxwell M. Modiba:** Writing – original draft, Investigation. **Márta Birkás:** Validation, Resources, Methodology, Conceptualization. **János Grósz:** Writing – original draft, Investigation. **Martin Kulhanek:** Writing – original draft, Methodology, Investigation. **Reinhard W. Neugschwandtner:** Writing – original draft, Resources, Project administration, Conceptualization. **Anna Hofer:** Investigation. **Viola Wagner:** Investigation. **Marion Windisch:** Investigation. **Karin Hage-Ahmed:** Writing – original draft, Visualization, Validation, Methodology, Investigation, Formal analysis, Data curation. **Kevin R. Butt:** Writing – original draft, Methodology, Investigation. **Pia Euteneuer:** Visualization, Supervision, Methodology, Investigation, Formal analysis, Data curation, Conceptualization.

## Acknowledgements and funding

We like to thank: The University of Natural Resources and Life Sciences, Vienna (Austria) for the possibility to publish this paper open access, all bachelor students and technicians who assisted with data collection in the trials. Special thanks to Sabine Daxböck-Horvath for the excellent technical assistance with the AMF. This research received funding from the Austrian-Hungarian Scientific and Technological Cooperation from the Austrian Agency for Education and Internationalisation (OeAD) and the Hungarian National Research Development and Innovation Office (NKFIH).

## Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Pia Euteneuer reports travel was provided by Austrian-Hungarian Scientific and Technological Cooperation from the Austrian Agency for Education and Internationalisation (OeAD). Barbara Simon reports travel was provided by Austrian-Hungarian Scientific and Technological Cooperation from National Research, Development and Innovation Office of Hungary (NKFIH). Hanaa T. M. Ibrahim reports travel was provided by Austrian-Hungarian Scientific and Technological Cooperation from National Research, Development and Innovation Office of Hungary (NKFIH). Reinhard Neugschwandtner reports travel was provided by Austrian-Hungarian Scientific and Technological Cooperation from the Austrian Agency for Education and Internationalisation (OeAD). Karin Hage-Ahmed reports travel was provided by Austrian-Hungarian Scientific and Technological Cooperation from the Austrian Agency for Education and Internationalisation (OeAD). If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Data availability

Data will be made available on request.

## References

- Abbott, L., Robson, A., 1985. Formation of external hyphae in soil by four species of vesicular arbuscular. *New Phytol.* 245–255. <https://doi.org/10.1111/j.1469-8137.1985.tb03653.x>.
- Amézketa, E., 1999. Soil aggregate stability: a review. *J. Sustain. Agric.* 14, 83–151. [https://doi.org/10.1300/J064v14n02\\_08](https://doi.org/10.1300/J064v14n02_08).
- Arai, M., Miura, T., Tsuzura, H., Minamiya, Y., Kaneko, N., 2018. Two-year responses of earthworm abundance, soil aggregates, and soil carbon to no-tillage and fertilization. *Geoderma* 332, 135–141. <https://doi.org/10.1016/j.geoderma.2017.10.021>.
- Arrázola-Vásquez, E., Larsbo, M., Capowicz, Y., Taylor, A., Sandin, M., Iseskog, D., Keller, T., 2022. Earthworm burrowing modes and rates depend on earthworm

- species and soil mechanical resistance. *Appl. Soil Ecol.* 178, 104568. <https://doi.org/10.1016/j.apsoil.2022.104568>.
- Bagnall, D.K., Morgan, C.L.S., 2021. SLAKES and 3D scans characterize management effects on soil structure in farm fields. *Soil Tillage Res.* 208, 104893. <https://doi.org/10.1016/j.still.2020.104893>.
- Bates, D., 2015. Using lme4: Mixed-Effects Modeling in R 158.
- Beck, H.E., McVicar, T.R., Vergopolan, N., Berg, A., Lutsko, N.J., Dufour, A., Zeng, Z., Jiang, X., Van Dijk, A.I.J.M., Miralles, D.G., 2023. High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Sci. Data* 10, 724. <https://doi.org/10.1038/s41597-023-02549-6>.
- BFW (Bundesforschungs- und Ausbildungszentrum für Wald, Naturgefahren und Landschaft), 2023. “eBOD” Digitale Bodenkarte Österreichs. <https://bodenkarte.at> (accessed 20 October 2024).
- Blanchart, E., Marilleau, N., Chotte, J.-L., Drogo ul, A., Perrier, E., Cambier, Ch., 2009. SWORM: an agent-based model to simulate the effect of earthworms on soil structure. *Eur. J. Soil Sci.* 60, 13–21. <https://doi.org/10.1111/j.1365-2389.2008.01091.x>.
- Bolan, N., Srivastava, P., Rao, C.S., Satyanaraya, P.V., Anderson, G.C., Bolan, S., Nortjé, G.P., Kronenberg, R., Bardhan, S., Abbott, L.K., Zhao, H., Mehra, P., Satyanarayana, S.V., Khan, N., Wang, H., Rinklebe, J., Siddique, K.H.M., Kirkham, M.B., 2023. Distribution, characteristics and management of calcareous soils. In: Sparks, Donald L. (Ed.), *Advances in Agronomy*. Academic Press, pp. 81–130. <https://doi.org/10.1016/bs.agron.2023.06.002>.
- Bottinelli, N., Capowicz, Y., 2021. Earthworm ecological categories are not functional groups. *Biol. Fertil. Soils* 57, 329–331. <https://doi.org/10.1007/s00374-020-01517-1>.
- Bottinelli, N., Hedde, M., Jouquet, P., Capowicz, Y., 2020. An explicit definition of earthworm ecological categories – Marcel Bouché’s triangle revisited. *Geoderma* 372, 114361. <https://doi.org/10.1016/j.geoderma.2020.114361>.
- Bouché, M.B., 1972. *Lombriciens de France - Ecologie et systématique (Paris)*.
- Briones, M.J.I., Schmidt, O., 2017. Conventional tillage decreases the abundance and biomass of earthworms and alters their community structure in a global meta-analysis. *Glob. Change Biol.* 23, 4396–4419. <https://doi.org/10.1111/gcb.13744>.
- Busari, M.A., Kukal, S.S., Kaur, A., Bhatt, R., Dulazi, A.A., 2015. Conservation tillage impacts on soil, crop and the environment. *Int. Soil Water Conserv. Res.* 3, 119–129. <https://doi.org/10.1016/j.iswcr.2015.05.002>.
- Butt, K.R., Grigoropoulou, N., 2010. Basic research tools for earthworm ecology. *Appl. Environ. Soil Sci.* 2010, 1–12. <https://doi.org/10.1155/2010/562816>.
- Butt, K.R., Gilbert, J.A., Kostecka, J., Lowe, C.N., Quigg, S.M., Euteneuer, P., 2022. Two decades of monitoring earthworms in translocated grasslands at Manchester Airport. *Eur. J. Soil Biol.* 113, 103443. <https://doi.org/10.1016/j.ejsobi.2022.103443>.
- Capowicz, Y., Cadoux, S., Bouchant, P., Ruy, S., Roger-Estrade, J., Richard, G., Boizard, H., 2009. The effect of tillage type and cropping system on earthworm communities, macroporosity and water infiltration. *Soil Tillage Res.* 105, 209–216. <https://doi.org/10.1016/j.still.2009.09.002>.
- Capowicz, Y., Marchán, D., Decaëns, T., Hedde, M., Bottinelli, N., 2024. Let earthworms be functional - definition of new functional groups based on their bioturbation behavior. *Soil Biol. Biochem.* 188, 109209. <https://doi.org/10.1016/j.soilbio.2023.109209>.
- Christian, E., Zicsi, A., 1999. Ein synoptischer Bestimmungsschlüssel der Regenwürmer Österreichs (Oligochaeta: Lumbricidae). *Bodenkult* 121–131.
- Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Austral Ecol.* 18, 117–143. <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>.
- Csuzdi, C., Zicsi, A., 2003. *Earthworms of Hungary: Annelida: Oligochaeta, Lumbricidae. Pedozoologica Hungarica, Hungarian Natural History Museum, Budapest*.
- Dekemati, I., Simon, B., Vinogradov, S., Birkás, M., 2019. The effects of various tillage treatments on soil physical properties, earthworm abundance and crop yield in Hungary. *Soil Tillage Res.* 194, 104334. <https://doi.org/10.1016/j.still.2019.104334>.
- Dekemati, I., Simon, B., Bogunovic, I., Kiscic, I., Kassai, K., Kende, Z., Birkás, M., 2020. Long term effects of ploughing and conservation tillage methods on earthworm abundance and crumb ratio. *Agronomy* 10, 1552. <https://doi.org/10.3390/agronomy10101552>.
- Dekemati, I., Simon, B., Bogunovic, I., Vinogradov, S., Modiba, M.M., Gyuricza, C., Birkás, M., 2021. Three-year investigation of tillage management on the soil physical environment, earthworm populations and crop yields in Croatia. *Agronomy* 11, 825. <https://doi.org/10.3390/agronomy11050825>.
- Driver, J.D., Holben, W.E., Rillig, M.C., 2005. Characterization of glomalin as a hyphal wall component of arbuscular mycorrhizal fungi. *Soil Biol. Biochem.* 37, 101–106. <https://doi.org/10.1016/j.soilbio.2004.06.011>.
- Edwards, C., Lofty, J.R., 1975. The invertebrate fauna of the park grass plots I. *Soil fauna. Rothamsted Exp. Stn. Rep.* 1974 (Part 2), 133–154. <https://doi.org/10.23637/ERADOC-1-9>.
- Euteneuer, P., Butt, K.R., 2025. Potential protocol for earthworm monitoring under adverse soil conditions or in small plots. *Appl. Soil Ecol.* 206, 105828. <https://doi.org/10.1016/j.apsoil.2024.105828>.
- Euteneuer, P., Butt, K.R., Wagentristl, H., Mayerová, M., Fér, M., 2024. What is the best way to measure earthworm-processed soil? A comparison of common water stable aggregates, the smartphone app MOULDER, and a novel SlakeLight method. *Appl. Soil Ecol.* 201, 105517. <https://doi.org/10.1016/j.apsoil.2024.105517>.
- Fajardo, M., McBratney, A.B., 2019. *Slakes: A Soil Aggregate Stability Smart-Phone App [Software]*.
- Fajardo, M., McBratney, Alex.B., Field, D.J., Minasny, B., 2016. Soil slaking assessment using image recognition. *Soil Tillage Res.* 163, 119–129. <https://doi.org/10.1016/j.still.2016.05.018>.

- Finn, D., Kopittke, P.M., Dennis, P.G., Dalal, R.C., 2017. Microbial energy and matter transformation in agricultural soils. *Soil Biol. Biochem.* 111, 176–192. <https://doi.org/10.1016/j.soilbio.2017.04.010>.
- Flynn, K.D., Bagnall, D.K., Morgan, C.L.S., 2020. Evaluation of SLAKES, a smartphone application for quantifying aggregate stability, in high-clay soils. *Soil Sci. Soc. Am. J.* 84, 345–353. <https://doi.org/10.1002/saj2.20012>.
- Franzluebbers, A.J., Wright, S.F., Stuedemann, J.A., 2000. Soil aggregation and glomalin under pastures in the southern Piedmont USA. *Soil Sci. Soc. Am. J.* 64, 1018–1026. <https://doi.org/10.2136/sssaj2000.6431018x>.
- Gajri, P., Arora, V., Prihar, S., 2002. *Tillage for Sustainable Cropping*. The Haworth Press, New York.
- Gompertz, B., 1833. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. In a letter to Francis Baily, Esq. F. R. S. &c. By Benjamin Gompertz. *Abstr. Pap. Print. Philos. Trans. R. Soc. Lond.* 2, 252–253. doi:<https://doi.org/10.1098/rsp1.1815.0271>.
- Grigoropoulou, N., Butt, K.R., 2010. Field investigations of *Lumbricus terrestris* spatial distribution and dispersal through monitoring of manipulated, enclosed plots. *Soil Biol. Biochem.* 42, 40–47. <https://doi.org/10.1016/j.soilbio.2009.09.021>.
- Gunina, A., Kuz'yakov, Y., 2015. Sugars in soil and sweets for microorganisms: review of origin, content, composition and fate. *Soil Biol. Biochem.* 90, 87–100. <https://doi.org/10.1016/j.soilbio.2015.07.021>.
- Helgason, B.L., Walley, F.L., Germda, J.J., 2010. No-till soil management increases microbial biomass and alters community profiles in soil aggregates. *Appl. Soil Ecol.* 46, 390–397. <https://doi.org/10.1016/j.apsoil.2010.10.002>.
- Holátko, J., Brtnický, M., Kučerík, J., Kotianová, M., Elbl, J., Kintl, A., Kynický, J., Benada, O., Datta, R., Jansa, J., 2021. Glomalin – truths, myths, and the future of this elusive soil glycoprotein. *Soil Biol. Biochem.* 153, 108116. <https://doi.org/10.1016/j.soilbio.2020.108116>.
- Houba, V.J.G., Temminghoff, E.J.M., Gaikhorst, G.A., Van Vark, W., 2000. Soil analysis procedures using 0.01 M calcium chloride as extraction reagent. *Commun. Soil Sci. Plant Anal.* 31, 1299–1396. <https://doi.org/10.1080/00103620009370514>.
- Irving, T.B., Alptekin, B., Kleven, B., Ané, J., 2021. A critical review of 25 years of glomalin research: a better mechanical understanding and robust quantification techniques are required. *New Phytol.* 232, 1572–1581. <https://doi.org/10.1111/nph.17713>.
- ISO 23611-1, 2018. DIN EN ISO 23611-1:2018-10, Bodenbeschaffenheit - Probenahme von Wirbellosen im Boden - Teil 1: Handauslese und Extraktion von Regenwürmern (ISO 23611-1:2018); Deutsche Fassung EN ISO 23611-1:2018. Beuth Verlag GmbH. <https://doi.org/10.31030/2823611>.
- Jansa, J., Mozafar, A., Anken, T., Ruh, R., Sanders, I., Frossard, E., 2002. Diversity and structure of AMF communities as affected by tillage in a temperate soil. *Mycorrhiza* 12, 225–234. <https://doi.org/10.1007/s00572-002-0163-z>.
- Jones, D., Willett, V., 2006. Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol. Biochem.* 38, 991–999. <https://doi.org/10.1016/j.soilbio.2005.08.012>.
- Jongmans, A.G., Pulleman, M.M., Balabane, M., Van Oort, F., Marinissen, J.C.Y., 2003. Soil structure and characteristics of organic matter in two orchards differing in earthworm activity. *Appl. Soil Ecol.* 24, 219–232. [https://doi.org/10.1016/S0929-1393\(03\)00072-6](https://doi.org/10.1016/S0929-1393(03)00072-6).
- Jug, D., Đurđević, B., Birkás, M., Brozović, B., Lipiec, J., Vukadinović, V., Jug, I., 2019. Effect of conservation tillage on crop productivity and nitrogen use efficiency. *Soil Tillage Res.* 194, 104327. <https://doi.org/10.1016/j.still.2019.104327>.
- Kabir, Z., O'Halloran, I.P., Fyles, J.W., Hamel, C., 1997. Seasonal changes of arbuscular mycorrhizal fungi as affected by tillage practices and fertilization: hyphal density and mycorrhizal root colonization. *Plant and Soil* 285–293.
- Kalbitz, K., Solinger, S., Park, J.-H., Michalzik, B., Matzner, E., 2000. Controls on the dynamics of dissolved organic matter in soils: a review. *Soil Sci.* 165, 277–304. <https://doi.org/10.1097/00010694-200004000-00001>.
- Kassambara, A., Mundt, F., 2020. *factoextra: Extract and Visualize the Results of Multivariate Data Analyses*.
- Kemper, W., Rosenau, R., 1986. Aggregate stability and size distribution. In: Sparks, D. (Ed.), *Methods of Soil Analysis. Part 1 – Physical and Mineralogical Methods*. Soil Science Society of America, Madison, pp. 425–442.
- Kenkel, N.C., Orloci, L., 1986. Applying metric and nonmetric multidimensional scaling to ecological studies: some new results. *Ecology* 67, 919–928. <https://doi.org/10.2307/1939814>.
- Klik, A., Rosner, J., 2020. Long-term experience with conservation tillage practices in Austria: impacts on soil erosion processes. *Soil Tillage Res.* 203, 104669. <https://doi.org/10.1016/j.still.2020.104669>.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World Map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* 15, 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>.
- Le, S., Josse, J., Husson, F., 2008. FactoMineR: an R package for multivariate analysis. *J. Stat. Softw.* 25, 1–18. <https://doi.org/10.18637/jss.v025.i01>.
- Lehmann, A., Zheng, W., Rillig, M.C., 2017. Soil biota contributions to soil aggregation. *Nat. Ecol. Evol.* 1, 1828–1835. <https://doi.org/10.1038/s41559-017-0344-y>.
- Lenth, R., 2022. *Emmeans: estimated marginal means, aka least-squares means*. R package version 1 (7), 5.
- Liebhart, G., Klik, A., Neugschwandner, Nolz, R., 2022. Effects of tillage systems on soil water distribution, crop development, and evaporation and transpiration rates of soybean. *Agric. Water Manag.* 12.
- Mairhofer, F., Barta, N., Euteneuer, P., Schuster, J., Bauer, A., Gronauer, A., 2019. Influence of tillage depth of a cultivator on the incorporation of crop residues of winter barley in a chernozem soil. *Bodenkult. J. Land Manag. Food Environ.* 70, 69–79. <https://doi.org/10.2478/boku-2019-0006>.
- Meng, L.-L., Srivastava, A.K., Kuća, K., Wu, Q.-S., 2022. Earthworm (*Pheretima guillelmi*)-mycorrhizal fungi (*Funneliformis mosseae*) association mediates rhizosphere responses in white clover. *Appl. Soil Ecol.* 172, 104371. <https://doi.org/10.1016/j.apsoil.2021.104371>.
- Mercy, L., 2017. INOQ Calculator Advanced. Evaluate the mycorrhizal rate according to a modified Trouvelot method. <https://doi.org/10.13140/RG.2.2.13641.03684>.
- Misirlioglu, I.M., Tsekova, R., Stojanović, M., 2016. On the presence of *Lumbricus terrestris* Linnaeus 1758 (Oligochaeta, Lumbricidae) on the Balkan Peninsula: some aspects of ecology and distribution. *Turk. J. Zool.* 40, 438–444. <https://doi.org/10.3906/zoo-1509-12>.
- Moebius-Clune, B., Moebius-Clune, D.J., Gugino, B., Idowu, O., Schindelbeck, R., Ristow, A., van Es, H., Thies, J., Shayler, H.A., McBride, M.B., Wolfe, D., Abawi, G., 2016. *Comprehensive Assessment of Soil Health- The Cornell Framework Manual*. Cornell University, Geneva, New York.
- Neugschwandner, R., Száková, J., Pachtrog, V., Tlustoš, P., Černý, J., Kulhánek, M., Kaul, H.-P., Euteneuer, P., Moitzi, G., Wagenstristl, H., 2020. Basic soil chemical properties after 15 years in a long-term tillage and crop rotation experiment. *Int. Agrophysics* 1, 133–140. <https://doi.org/10.31545/intagr/114880>.
- Neugschwandner, R.W., Száková, J., Pachtrog, V., Tlustoš, P., Kulhánek, M., Černý, J., Kaul, H.-P., Wagenstristl, H., Moitzi, G., Euteneuer, P., 2022. Exchangeable and plant-available macronutrients in a long-term tillage and crop rotation experiment after 15 years. *Plants* 11, 565. <https://doi.org/10.3390/plants11040565>.
- Nimmo, J., Perkins, K., 2002. Aggregate stability and size distribution. In: Dane, Jacob H., Topp, G. Clarke (Eds.), *Physical Methods, Methods of Soil Analysis*. SSSA, Madison, pp. 317–328. <https://doi.org/10.2136/sssabookser5.4.c14>.
- Obour, P.B., Lamané, M., Edwards, G., Sørensen, C.G., Munkholm, L.J., 2017. Predicting soil workability and fragmentation in tillage: a review. *Soil Use Manage.* 33, 288–298. <https://doi.org/10.1111/sum.12340>.
- Oksanen, J.F., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., O'Hara, R. B., Simpson, G., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2020. *vegan: community ecology package*. R package version 2, 5–7.
- ÖNORM L 1080, 1999. *Chemische Bodenuntersuchungen – Bestimmung des organischen Kohlenstoffs durch trockene Verbrennung*.
- Paliy, O., Shankar, V., 2016. Application of multivariate statistical techniques in microbial ecology. *Mol. Ecol.* 25, 1032–1057. <https://doi.org/10.1111/mec.13536>.
- Pelosi, C., Taschen, E., Redecker, D., Blouin, M., 2024. Earthworms as conveyors of mycorrhizal fungi in soils. *Soil Biol. Biochem.* 189, 109283. <https://doi.org/10.1016/j.soilbio.2023.109283>.
- Péres, G., Bellido, A., Curmi, P., Marmonier, P., Cluzeau, D., 2010. Relationships between earthworm communities and burrow numbers under different land use systems. *Pedobiologia* 54, 37–44. <https://doi.org/10.1016/j.pedobi.2010.08.006>.
- Posit team, 2023. *RStudio: Integrated Development Environment for R*.
- R Core Team, 2023. *R: A Language and Environment for Statistical Computing*. R Found. Stat. Comput. Vienna Austria.
- Rieke, E.L., et al., 2022. Evaluation of aggregate stability methods for soil health. *Geoderma* 428, 116156. <https://doi.org/10.1016/j.geoderma.2022.116156>.
- Rillig, M.C., Wright, S.F., Nichols, K.A., Schmidt, W.F., 2001. Large contribution of arbuscular mycorrhizal fungi to soil carbon pools in tropical forest soils. *Plant and Soil* 167–177.
- Rosinger, C., Bodner, G., Bernardini, L.G., Huber, S., Mentler, A., Sae-Tun, O., Scharf, B., Steiner, P., Tintner-Olifiers, J., Keiblinger, K., 2022. Benchmarking carbon sequestration potentials in arable soils by on-farm research on innovative pioneer farms. *Plant and Soil*. <https://doi.org/10.1007/s11104-022-05626-8>.
- Rosner, K., Bodner, G., Hage-Ahmed, K., Steinkellner, S., 2018. Long-term soil tillage and cover cropping affected arbuscular mycorrhizal Fungi, nutrient concentrations, and yield in sunflower. *Agron. J.* 110, 2664–2672. <https://doi.org/10.2134/ agronj2018.03.0177>.
- Sae-Tun, O., Bodner, G., Rosinger, C., Zechmeister-Boltenstern, S., Mentler, A., Keiblinger, K., 2022. Fungal biomass and microbial necromass facilitate soil carbon sequestration and aggregate stability under different soil tillage intensities. *Appl. Soil Ecol.* 179, 104599. <https://doi.org/10.1016/j.apsoil.2022.104599>.
- Saidy, A.R., Smernik, R.J., Baldock, J.A., Kaiser, K., Sanderman, J., 2015. Microbial degradation of organic carbon sorbed to phyllosilicate clays with and without hydrous iron oxide coating. *Eur. J. Soil Sci.* 66, 83–94. <https://doi.org/10.1111/ejss.12180>.
- Säle, V., Aguilera, P., Laczko, E., Mäder, P., Berner, A., Zihlmann, U., Van Der Heijden, M.G.A., Oehl, F., 2015. Impact of conservation tillage and organic farming on the diversity of arbuscular mycorrhizal fungi. *Soil Biol. Biochem.* 84, 38–52. <https://doi.org/10.1016/j.soilbio.2015.02.005>.
- Schrader, S., Zhang, H., 1997. Earthworm casting: stabilization or destabilization of soil structure? *Soil Biol. Biochem.* 29, 469–475. [https://doi.org/10.1016/S0038-0717\(96\)00103-4](https://doi.org/10.1016/S0038-0717(96)00103-4).
- Singh, A.K., Rai, A., Singh, N., 2016. Effect of long-term land use systems on fractions of glomalin and soil organic carbon in the Indo-Gangetic plain. *Geoderma* 277, 41–50. <https://doi.org/10.1016/j.geoderma.2016.05.004>.
- Singh, A.K., Zhu, X., Chen, C., Wu, J., Yang, B., Zakari, S., Jiang, X.J., Singh, N., Liu, W., 2022. The role of glomalin in mitigation of multiple soil degradation problems. *Crit. Rev. Environ. Sci. Technol.* 52, 1604–1638. <https://doi.org/10.1080/10643389.2020.1862561>.
- Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* 32, 2099–2103. [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6).
- Six, J., Bossuyt, H., Degryse, S., Denef, K., 2004. A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil Tillage Res.* 79, 7–31. <https://doi.org/10.1016/j.still.2004.03.008>.

- Sokol, N., Sandermann, J., Bradford, M., 2019. Pathways of mineral-associated soil organic matter formation: integrating the role of plant carbon source, chemistry, and point of entry. *Glob. Change Biol.* 25, 12–24. <https://doi.org/10.1111/gcb.14482>.
- Stroud, J.L., Irons, D., Watts, C.W., Whitmore, A.P., 2016. *Lumbricus terrestris* abundance is not enhanced after three years of compost amendments on a reduced tillage wheat cultivation conversion. *Appl. Soil Ecol.* 98, 282–284. <https://doi.org/10.1016/j.apsoil.2015.10.007>.
- Thomopoulos, S., Elsgaard, L., Munkholm, L.J., Ravnkov, S., 2023. Evaluation of the relation between soil biomass of arbuscular mycorrhizal fungi and glomalin-related soil protein in conservation agriculture. *Soil Biol. Biochem.* 187, 109222. <https://doi.org/10.1016/j.soilbio.2023.109222>.
- Trouvelot, A., Kough, J., Gianinazzi-Pearson, V., 1986. Mesure du taux de mycorrhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. In: Gianinazzi-Pearson, V., Gianinazzi, S. (Eds.), *Physiological and Genetical Aspects of Mycorrhizae*. INRA, Paris, pp. 217–221.
- Ussiri, D.A.N., Johnson, C.E., 2004. Sorption of organic carbon fractions by Spodosol mineral horizons. *Soil Sci. Soc. Am. J.* 68, 253–262. <https://doi.org/10.2136/sssaj2004.2530>.
- van Groenigen, K.-J., Bloem, J., Bååth, E., Boeckx, P., Rousk, J., Bodé, S., Forristal, D., Jones, M.B., 2010. Abundance, production and stabilization of microbial biomass under conventional and reduced tillage. *Soil Biol. Biochem.* 42, 48–55. <https://doi.org/10.1016/j.soilbio.2009.09.023>.
- Vierheilig, H., Coughlan, A.P., Wyss, U., Piché, Y., 1998. Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Appl. Environ. Microbiol.* 64, 5004–5007. <https://doi.org/10.1128/AEM.64.12.5004-5007.1998>.
- Walkley, A., 1947. A critical examination of a rapid method for determining organic carbon in soils —effect of variations in digestion conditions and of inorganic soil constituents. *Soil Sci.* 63, 251–264. <https://doi.org/10.1097/00010694-194704000-00001>.
- Weidhuner, A., Hanauer, A., Krausz, R., Crittenden, S.J., Gage, K., Sadeghpour, A., 2021. Tillage impacts on soil aggregation and aggregate-associated carbon and nitrogen after 49 years. *Soil Tillage Res.* 208, 104878. <https://doi.org/10.1016/j.still.2020.104878>.
- Weninger, T., Kreiselmeier, J., Chandrasekhar, P., Julich, S., Feger, K.-H., Schwärzel, K., Bodner, G., Schwen, A., 2019. Effects of tillage intensity on pore system and physical quality of silt-textured soils detected by multiple methods. *Soil Res.* 57, 703. <https://doi.org/10.1071/SR18347>.
- Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York.
- WRB, 2014. *World Reference Base for Soil Resources 2014: International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*. FAO, Rome.
- Wright, S.F., Upadhyaya, A., 1996. Extraction of an abundant and unusual protein from soil and comparison with hyphal protein of arbuscular mycorrhizal fungi. *Soil Sci.* 161, 575–586. <https://doi.org/10.1097/00010694-199609000-00003>.
- Wright, S.F., Upadhyaya, A., 1998. A survey of soils for aggregate stability and glomalin, a glycoprotein produced by hyphae of arbuscular mycorrhizal fungi. *Plant and Soil* 97–107. <https://doi.org/10.1023/A:1004347701584>.
- Zicsi, A., 1994. Die Regenwürmer Österreichs (Oligochaeta: Lumbricidae) mit Bestimmungstabellen der Arten. *Verh. Zool.-Bot. Ges. Österr.* 37–74.