

Central Lancashire Online Knowledge (CLoK)

Title	The palaeoenvironmental potential of bioarchaeological isotope data
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/56162/
DOI	https://doi.org/10.1038/s43247-025-02507-7
Date	2025
Citation	Depaermentier, Margaux L. C., Kempf, Michael, Vander Linden, Marc, Britton, Kate, Madgwick, Richard, Büntgen, Ulf, Drucker, Dorothée G., Jones, Jennifer, Snoeck, Christophe et al (2025) The palaeoenvironmental potential of bioarchaeological isotope data. Communications Earth & Environment, 6 (1).
Creators	Depaermentier, Margaux L. C., Kempf, Michael, Vander Linden, Marc, Britton, Kate, Madgwick, Richard, Büntgen, Ulf, Drucker, Dorothée G., Jones, Jennifer, Snoeck, Christophe and Stevens, Rhiannon E.

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1038/s43247-025-02507-7>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Central Lancashire Online Knowledge (CLoK)

Title	The palaeoenvironmental potential of bioarchaeological isotope data
Type	Article
URL	https://clok.uclan.ac.uk/id/eprint/56162/
DOI	https://doi.org/10.1038/s43247-025-02507-7
Date	2025
Citation	Depaermentier, Margaux L. C., Kempf, Michael, Vander Linden, Marc, Britton, Kate, Madgwick, Richard, Büntgen, Ulf, Drucker, Dorothée G., Jones, Jennifer R., Snoeck, Christophe et al (2025) The palaeoenvironmental potential of bioarchaeological isotope data. Communications Earth & Environment, 6 (1). ISSN 2662-4435
Creators	Depaermentier, Margaux L. C., Kempf, Michael, Vander Linden, Marc, Britton, Kate, Madgwick, Richard, Büntgen, Ulf, Drucker, Dorothée G., Jones, Jennifer R., Snoeck, Christophe and Stevens, Rhiannon E.


It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1038/s43247-025-02507-7>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clok.uclan.ac.uk/policies/>



The palaeoenvironmental potential of bioarchaeological isotope data



Margaux L. C. Depaermentier ¹✉, Michael Kempf ^{2,3}, Marc Vander Linden ⁴, Kate Britton ⁵, Richard Madgwick ⁶, Ulf Büntgen ^{3,7,8}, Dorothée G. Drucker ⁹, Jennifer R. Jones ¹⁰, Christophe Snoeck ¹¹ & Rhiannon E. Stevens ¹²

Bioarchaeology not only provides insights into human, animal, and environmental ecology, but also generates huge amounts of stable and radiogenic isotope data that are not well recognised by other disciplines. Here, we present potential avenues for the integration and interpretation of archaeological isotope data into environmental studies. We emphasise the large spatio-temporal scales on which isotope patterns can be observed, for example using isoscapes, the limitations and potential pitfalls that come with isotope data from archaeological research, and future cross-disciplinary collaborations between bioarchaeology and other palaeo-disciplines.

Isotope analyses were introduced in archaeological research in the late 1970s, when J. C. Vogel and N. J. van der Merwe applied carbon isotope analyses to investigate maize cultivation in North America¹. In 1985, J. E. Ericson² provided the basis for mobility studies using strontium isotopes. A few decades later, isotope analyses of carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$), oxygen ($\delta^{18}\text{O}$), radiogenic strontium ($^{87}\text{Sr}/^{86}\text{Sr}$)^{3–5} and increasingly sulphur ($\delta^{34}\text{S}$)⁶, hydrogen (δD)⁷, stable strontium ($\delta^{88}\text{Sr}$)⁸, zinc ($\delta^{66}\text{Zn}$)⁹, lead ($^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, $^{208}\text{Pb}/^{204}\text{Pb}$)¹⁰, neodymium ($^{143}\text{Nd}/^{144}\text{Nd}$)¹¹ and also compound specific approaches¹² have become common tools in archaeological research.

In bioarchaeology, i.e. the study of biological remains including humans, faunal and flora (but not restricted to osteoarchaeology), isotope sampling is limited to a fraction of sites (depending on preservation and access) and the research questions are formulated by archaeological interest. The resulting archives consist of individual high-quality records (Supplementary Note 1) in which $^{87}\text{Sr}/^{86}\text{Sr}$, $^{208}\text{Pb}/^{206}\text{Pb}$, and more recently $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes are mostly dedicated to mobility studies^{10,11,13}, $\delta^{18}\text{O}$ and δD to mobility and climatic reconstruction^{14,15} and $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^2\text{H}$, $\delta^{66}\text{Zn}$ ¹⁶ and $\delta^{34}\text{S}$ isotopes are applied to investigate dietary, social, environmental, climatic and agricultural changes^{17–19}. The sample size strongly varies depending on the goals of the study: a single individual can represent an entire case study, whereas community-level targeted investigations can

comprise sample sizes of many tens of individuals²⁰. Various isotope systems are moreover increasingly combined into multi-proxy and multi-tissue approaches to improve the measured signals^{11,19,21}, to contrast differing signals related to skeletal remodelling and isotope turnover processes²², and to minimise the negative effects of destructive sampling^{22–24}. To our knowledge no study has yet integrated $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, $\delta^{34}\text{S}$, $^{87}\text{Sr}/^{86}\text{Sr}$, and $^{208}\text{Pb}/^{206}\text{Pb}$, although protocols exist to enable this²⁵.

Several initiatives have started to create regional, temporal and thematic datasets to gather and centralise these fragmented datasets^{26–30}. Moreover, some archaeological studies have taken advantage of such huge datasets to track large-scale cultural^{31,32}, ecological^{33–36}, climatic^{37,38} or environmental^{39,40} patterns. However, the scientific potential of this archive remains comparatively untapped, especially in the perspective of multi-disciplinary research. This implies that palaeo-environmental, -ecological and -climatic research is restricted to areas where proxies such as ice cores, speleothems, tree-rings, molluscs or lake/marine sediments are available, which cannot necessarily be easily connected to human activity, while bioarchaeological samples have a nearly global coverage and intrinsically enable to investigate human-environment interactions. This paper hence suggests how bioarchaeological isotope data can be integrated into palaeo-studies as they provide complementary environmental and climatic information that is not available from other archives.

¹Department of Ancient Civilizations, University of Basel, Basel, Switzerland. ²Quaternary Geology, Department of Environmental Sciences, University of Basel, Basel, Switzerland. ³Department of Geography, University of Cambridge, Cambridge, UK. ⁴Institute for the Modelling of Socio-Environmental Transitions, Bournemouth University, Fern Barrow, Poole, Bournemouth, UK. ⁵Department of Archaeology, University of Aberdeen, Aberdeen, Scotland, UK. ⁶School of History, Archaeology and Religion, Cardiff University, Cardiff, UK. ⁷Department of Geography, Masaryk University, Brno, Czech Republic. ⁸CzechGlobe, Global Change Research Institute CAS, Brno, Czech Republic. ⁹Senckenberg Centre for Human Evolution and Palaeoenvironment an der Universität Tübingen, Tübingen, Germany. ¹⁰Research Centre for Field Archaeology and Forensic Taphonomy, School of Law and Policing, University of Central Lancashire, Preston, UK. ¹¹Archaeology, Environmental changes and Geo-Chemistry, Vrije Universiteit Brussel, Brussels, Belgium. ¹²UCL Institute of Archaeology, University College London, London, UK. ✉e-mail: m.depaermentier@unibas.ch

Recycling bioarchaeological isotope data in environmental research

Bioarchaeological studies have contributed to palaeoecological research by revealing migratory and dietary behaviour of domesticated (affected by agropastoral and husbandry practices) and wild species—including now extinct ones^{36,41,42}—and by highlighting their interaction with humans^{43–45}. In most cases, these isotope data are biased towards human activity (see Supplementary Note 1)—with the exception of studies conducted in collaboration with palaeontology^{46–48}. But more importantly, isotope ratios from bioarchaeological tissues can serve as palaeoclimatic and palaeoenvironmental proxies, which is not recognised in other palaeo-disciplines. This section presents the palaeoclimatic and palaeoenvironmental potential of bioarchaeological isotope data using several examples and emphasising both the spatial and the temporal dimensions.

Environmental aspects

The $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values of animal tissues are ultimately related to the $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values of consumed environmental water, and thus reflect, e.g. latitude and longitude, altitude, distance to the moisture source and seasonality (see a review in O⁴⁹ and H⁷). Other isotope systems can be targeted in faunal tissues reflecting other aspects of local palaeoclimatic conditions (see Stevens et al in press⁵⁰ for a review). For example, plant $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$ and $\delta^{66}\text{Zn}$ values are affected by local variability in humidity and aridity, salinity, daylight, or forest cover versus open landscapes. These ‘signatures’

are transferred through the food chain so that the isotopic composition of animal and human tissues reflect the organisms dietary choices and contain information about the palaeoenvironmental and palaeoecological context of their habitat^{9,49–51}—provided that the archaeozoological material is not biased towards grazers or browsers only, or that mixed-feeding herbivores are available from the site. Such isotope data can thus be used to retrace palaeoenvironmental diversity at a given archaeological site during a specific time frame^{21,38}, or to reconstruct surrounding ecosystems, for example in coastal areas^{52,53}, wetlands⁵⁴, or steppe²¹ (Fig. 1). This approach also had particular success for reconstructing environmental change, for example associated with MIS 3, when the Neanderthals became extinct⁵⁵, or the Last Glacial Maximum (LGM) and refugia regions^{34,56}.

Several examples thus underline the ecological and palaeoenvironmental potential of bioarchaeological isotope data. In C_3 -dominated ecosystems under peri-arctic, boreal and temperate conditions, contrast in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values reflect the habitat and foraging preferences of herbivores. A striking example of niche partitioning based on specialised diet is observed among large ungulates of the mammoth steppe during the Last Glacial Period, a pattern that persisted across broad temporal and geographical scales. Reindeer exhibit the highest $\delta^{13}\text{C}$ values due to their specific and substantial consumption of lichen, showing few if any overlap with horses, which have among the lowest $\delta^{13}\text{C}$ values³⁶. Mammoths show the highest $\delta^{15}\text{N}$ values, likely due to grazing on more mature plants and possibly coprophagy, while woolly rhinoceroses and large bovids present





Fig. 1 | Characteristics of bioarchaeological and environmental proxy archives. The bioarchaeological samples are (from left to right): hair, nail, tooth, bones with short or long turnover rates, bones from young animals/individuals, and charred grains. The temporal depth varies between proxies, yet the palaeontology and bioarchaeology hardly provide any continuous time series and the bioarchaeological sample frequency is considerably reduced towards the oldest samples. The dating uncertainty or precision can reach one calendar year for tree-rings, speleothems, varved lake's sediments, speleothems, young ice cores, corals and molluscs, but it can be decadal to millennial for most proxies depending on the temporal depth and the dating method. The proxy resolution varies from sub-annual to decadal for most proxies, it can be (multi-)centennial for marine sediments and other layers that form slowly. The global spatial distribution vary between proxies: ice cores depend on global glacier distribution (light blue signature; source: <https://wgms.ch/downloads/DOI->

[WGMS-FoG-2025-02b.zip](https://wgms.ch/downloads/DOI-)); speleothems depend on karst regions (light brown signature; source: https://download.bgr.de/bgr/grundwasser/whymap/shp/WHYMAP_WOKAM_v1.zip); corals are restricted to tropical areas (dark blue signature; source: <https://data.basin.org/datasets/b983863c0a1a41e8839383b40ade437d/>); molluscs are restricted to water bodies or archaeological sites, marine and lakes sediments originate from the respective water bodies, tree-rings depend on past forest covers and to some extent on archaeological sites (green signature: global tree ring record distribution; source: https://gis.ncdc.noaa.gov/kml/paleo_tree.kmz, last data access: 06th June 2025), palaeontological and bioarchaeological samples have a global extent but depend on past human activities, excavated sites and preservation conditions. The spatial scale reflected by each proxy goes from the local to the global scale, but even local proxies may inform about supra-regional processes when considered altogether. These characteristics are summarised in Supplementary Table 1. Figure © Michael Kempf 2025.

Fig. 2 | Scale dependency of ecological functionality. The upper panel shows decreasing climatic and ecological scales from GCMs (Global Climate Models) to supraregional ecozones, regional habitats, and local and micro niches (realised niches). The lower panel shows increasing spatial operations, including small-scale fragmentation, simplification such as regional aggregation and binning, and estimation processes such as interpolation. Figure © Michael Kempf 2025.



intermediate $\delta^{15}\text{N}$ values^{50,57}. Change in the $\delta^{15}\text{N}$ baseline is associated with complex, though sometimes decipherable, environmental conditions. In particular, the increase in $\delta^{15}\text{N}$ across Europe during the Lateglacial to early Holocene transition may be linked indirectly to rising temperatures^{35,58–61}. Geographical variation in $\delta^{15}\text{N}$ values of large ungulates during the Lateglacial period is best explained by local environmental conditions, through the lingering effects of permafrost on N-cycling in soils^{35,59,62}. A decline in isotopic niche partitioning reflects the changing post-LGM conditions, as seen in the ^{15}N -depleted mammoths in Central Europe⁶³ and the increasing overlap in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values among large ungulates^{62,64}.

From a palaeoclimatic perspective, archaeological charcoals and molluscan shells represent further well-suited materials for investigating seasonal patterns and climate change on the site scale⁶⁵. At the global scale, bioarchaeological proxies further allow for tracking larger climatic processes such as El Niño⁶⁶ or variability in the monsoon⁶⁷. When considered separately, archaeological sites thus offer site-specific or micro-regional insights into the effects of large-scale phenomena, while the accumulation of bioarchaeological data from one or several regions provides itself a record of large-scale phenomena (see Figs. 1, 2 and Section 3.2). Moreover, isotope data from diverse species further reflect different trends depending on their individual niches³⁴, while palaeoenvironmental changes are not always reflected across all niches, such as the impact of LGM on ibex, but not on red deer dietary ecology in Northern Spain⁵⁶, or the impact of the Heinrich 4 (~39/40 Ka BP) event on reindeer feeding ecology at Les Cottés (France) relative to other species³⁶. Bioarchaeological isotope data are therefore complementary datasets to fill gaps among proxies from the environmental sciences.

This also applies in terms of geographical distribution of the records, as (zoo-)archaeological remains often survive when other environmental proxies do not. Stable isotope analysis of faunal remains provides a proxy for local environmental conditions, when many other proxies such as ice cores, sea cores, and speleothems are often distant from these locations and/or do not necessarily reflect local climatic conditions⁶⁸. Zooarchaeological remains that originate from human activity can even provide contrasting evidence to other site-specific proxies that may not be contemporary to human site-use. For example, $\delta^{18}\text{O}$ data from bovid teeth at the site of La Ferrassie, France, demonstrated that Neanderthals used the site during milder phases of a (broadly) cool period⁶⁹. The data contrasted with wider regional environmental records for the period, as well as certain sediment records at the site itself, which also attested to these cooler conditions. This highlights the diverse scales of the different proxies, and the unique insights afforded through the isotope analysis of anthropogenically-derived faunal remains in reconstructing climate at the scale of human behaviour⁶⁹. When combining and comparing different proxy datasets, the use of modern data is also important for testing the application of these proxies and feed current

models in terms of spatial representativity⁷⁰. This further offers the potential to use bioarchaeological isotope data to model phenomena for which proxies exist only in recent times⁷¹. Combining either contemporaneous records from different regions or diachronic records from a defined geographical area thus represent invaluable records for past processes that are otherwise recovered only in specific contexts or restricted geographical areas^{72,73}.

When working with archaeological datasets, wider consideration needs to be given to the impact of anthropogenic activity on the stable isotope record. Concerning hunter-gatherer populations, the faunal record is a product of hunting strategies employed, and reflect times when a site was occupied by human populations, potentially reflecting seasonal occupation preferences or occupation only during particular climatic conditions, as seen at the Middle Palaeolithic site of Axlor in Northern Spain⁷⁴. When working with material pertaining to (agro-)pastoralist populations the application of manure to fodder crops can impact on observed values^{75,76} as well as keeping animals in pens or enclosures⁷⁷, transhumance^{77–79} the provision of specific⁸⁰ or unusual fodder or graze such as salt-marsh, seaweed and marine resources^{81,82}, or the use of diverse or distinctive pasture locations such as forests⁸³ can all impact on the stable isotope record. In contrast to palaeontological material^{84–86} the potential of such bioarchaeological samples to reflect palaeoenvironmental conditions would be biased by the human impact on feeding habits. Small mammals such as rodents from the archaeological context are good alternatives to target the local environment^{87,88}. However, as explained above, the targeting of anthropogenically-derived remains of prey-species (specifically large obligate-drinking ungulates) can conversely be viewed as an inherent bias that is favourable, resulting in a climatic proxy that is directly linked to human site-use.

Moreover, a certain degree of mobility of humans, animals (e.g. for food, transport), food resources and animal fodder can always be expected, in particular for the historical periods^{18,89}. Likewise, trading of resources between rural and urban locations and between central places and hinterlands⁹⁰ can result in the displacement of remains. In such cases, the isotope composition of bioarchaeological material would not be representative for local conditions and hence could not be used for reconstructing the local palaeoenvironment. It must be further kept in mind that regardless of the economy (hunter-gatherer, pastoralists, farmer), the material found at an archaeological site results from human selection and is biased towards what survived until today due to cultural choices or preservation issues. It might therefore not represent all biomes from the past local ecosystem—which is, however, a problem known from modern samples as well⁹¹. Working with archaeologists who understand such a record can help to ensure that consideration is given to some of these nuances and how they might be influencing any trends observed.

Temporal and seasonal aspects

Isotope analysis of incrementally developed structures such as speleothems^{68,73}, tree-rings^{72,92}, shells⁹³, and stratified sequences like lake sediments⁹⁴, provide multifaceted archives of past climatic and environmental conditions. Such records are valuable taken separately or combined, producing complementary proxy datasets^{95,96}. Data from these archives cover different chronological and geographical scales on varying temporal resolutions with proxy-inherent limitations (Fig. 1). The analysis of speleothems and tree rings, for example, captures two different seasonal signals: mean annual or cooler season (fall to spring) precipitation in speleothems^{68,73} and the growing season signal in tree rings⁹². Climatic variability deriving from one or the other is a characterisation of a specific period of the year and an enhanced drought signal in one proxy does not necessarily occur in the other—triggering an overestimation of climatic extremes across a multi-annual period.

In comparison, the measured isotopic composition of bioarchaeological samples informs on contexts as old as several millions of years, with a resolution varying from a few weeks to several decades depending on the tissue and the sampling method (Table 1 and Supplementary Note 1). For example, $\delta^2\text{H}$ isotope analysis of bone collagen can provide insight into time-averaged (i.e. supra-annual) precipitation⁹⁷. Similarly, $\delta^{18}\text{O}$ values from 'bulk' enamel samples (when undertaken to ensure a full year of growth is sampled) can allow the reconstruction of mean annual temperatures^{98,99}. On the other hand, high-resolution intra-tooth micro-sampling on animal teeth offers continuous data for the period during which a particular tooth formed, covering both the fall/winter and the spring/summer seasons¹⁰⁰, thus permitting the reconstruction of seasonal palaeoclimatic conditions^{14,38} over a full year. This approach has been applied at a number of Neanderthal and early modern human sites in Europe. For example, at Ilsehöhle in Ranis, Germany, intra-tooth sampling and O isotope analysis of horses (combined with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis of faunal collagen) revealed decreasing temperatures throughout the sequence but also that extreme seasonality and coolest winter temperatures coincided with earliest modern human use of the site (the initial Upper Palaeolithic)²¹.

However, such approaches rely on dating and clear stratigraphic association, and the precision of the sample's absolute dating depends on the period considered, and on the method used, e.g. radiocarbon dating¹⁰¹, luminescence¹⁰², dendrochronology⁹², stratigraphy, typology¹⁰³. In contrast to the possibility of dating tree-rings, ice cores or speleothems to one specific calendar year, the associated chronological uncertainty among bioarchaeological samples (despite radiocarbon dating, for example) can vary dramatically and stratigraphic (or other) relationships at sites remain pivotal. The duration of the occupation might further be difficult to estimate¹⁰⁴. The absolute dating resolution of bioarchaeological samples may be as large as several hundreds of years in the case of the oldest samples, when there is a prolonged lack of change in archaeological material culture, or when the calibration curve for radiocarbon dates hits a plateau^{105–108}, while it can reach a few years or decades when methods are combined and/or when the archaeological context provides a high-resolution chronology such as during the Early Middle Ages^{109,110}. Nevertheless, bioarchaeological research can provide time sequences in the same way as ice cores or lake varves.

Following the tree-ring approach by Loader and colleagues¹¹¹ and by Black et al. on molluscs¹¹² it may be possible to compare large sequential isotope datasets from animal teeth to, e.g. tree-ring and/or speleothem isotope data, and thus to integrate these bioarchaeological records into existing time series via cross-dating. A machine learning approach at the global scale and over long chronologies¹¹³, would facilitate linking of bioarchaeological sequential isotope data to palaeoclimatic and palaeoenvironmental parameters and completion or testing of existing models. This will imply further treatment of the data (i.e. interpolation, imputation, averaging), creating a loss of information (see also Fig. 2). Introducing various modelling procedures represents a first step to deal with these issues.

Following metadata guidelines provided by environmental sciences⁹⁶ may further facilitate the integration of bioarchaeological isotope data into

existing datasets such as the global Iso2k database, expanding the latter to the BCE time scale (see also ref. 26 and the *Pandora initiative* since 2023). Despite being discontinuous and uneven sequences, they would fill temporal and seasonal gaps and provide new insights and dimensions on phenomena recorded elsewhere. Yet a preliminary cross-disciplinary discussion is required to make sure that workflows align not only within⁷² but also between disciplines. Targeted sampling strategies within an area where comparable proxy data from speleothems, tree rings or others are available—and where a satisfying criterion regarding sample size can be met—would allow testing of this approach.

Isoscapes and the spatial dimension of isotope variability

The production of isotope data covering a wide geographical area provides the potential for generating isoscapes, which are detailed maps of isotope distributions across landscapes. These are fundamental isotope baselines for archaeological and ecological research, food sciences, and forensics in order to investigate mobility patterns and provenance. This is crucial when the isotope composition varies considerably over space in a predictable way according to known factors^{10,114–116}. In parallel, some isotope systems are particularly sensitive to environmental and climatic changes and/or to human activities: their isoscapes thus allow for tracking spatial variations in environmental settings at a specific time—or over time^{61,117}. In this context, bioarchaeological research has strongly contributed to the production of isoscapes, as O, S, Sr, and Pb isoscapes are purposely created to interpret the bioarchaeological isotope data in terms of mobility patterns^{5,10,117,118}. This unique and extended dataset can be used in the past as it is in ecological research, since it provides a reference framework for several regions of the world, enabling the investigation of animal ecology and migration patterns^{119,120}. It may also contribute to baselines used to establish the authenticity of product origins^{121,122}. As a by-product of isotope data accumulation in bioarchaeology, isoscapes for C and N isotopes can now also be produced, enabling to track spatial and temporal variations in environmental parameters³⁵. Yet the anthropocentric questions triggering the production of these bioarchaeological isoscapes and the diversity of approaches applied in this making process need to be considered to facilitate the reusability of isoscapes generated in bioarchaeological research.

Archives for targeted isoscapes

The development of isoscapes relies on primary or secondary data that can be anchored to a specific region, terrain or environment. The breadth of potential archives for these data across the lithosphere, biosphere, hydrosphere and atmosphere is almost limitless, yet bioarchaeological research targets specific samples to meet specific needs, as the modern ecosystem may differ considerably from the past. It is not only modern chemical contamination (which is a key part of the provenancing process¹²³), but also natural chemical cycling and environmental processes that cause these chronological shifts in certain isotope proxies (e.g. $\delta^{15}\text{N}$ ⁶¹, $^{87}\text{Sr}/^{86}\text{Sr}$ ¹²⁴, $\delta^{34}\text{S}$ ⁴⁰, and Stevens et al.⁵⁰ in for a general review). The most work on this issue relates to $^{87}\text{Sr}/^{86}\text{Sr}$ analysis (see summary in Holt et al.¹¹⁵), but most considerations are applicable across isotope systems.

The impact of these processes varies across isotope systems, and archives for bioarchaeological research are selected accordingly. For example, to overcome the problem of Pb pollution in the modern environment—whose impact is noticeable since the advent of metallurgy¹⁰ and in particular from the Roman time¹²⁵—base geological samples are generally favoured in Pb isoscapes instead of environmental samples (e.g. plants, soils and water), with prehistoric faunal samples providing an additional useful source¹⁰. Environmental samples are of greater value for $^{87}\text{Sr}/^{86}\text{Sr}$, but the bioarchaeological approach prefers samples from 'pristine' landscapes, away from roads, urban areas or arable agriculture to avoid modern contamination¹²⁶. Even traditional agricultural practices, such as the use of seaweed as a fertiliser, may shift biosphere signatures—i.e. closer to marine values in Sr¹²⁷ and S¹²⁸.

Table 1 | Overview of the main bioarchaeological samples and analysed tissues and their potential with respect to diverse isotope systems (‘: When considering no particular diet such as low- or high-protein diet)

Sample type	Tissue type	Tissue properties	Maximal age of sample	Individuals lifespan represented	Susceptibility to diagenesis	Measurements	fractionation between diet and tissue	part of the diet ¹ (mainly) reflected
Dental enamel bulk sample	Biomineral	Incremental formation, acellular, avascular, no remodelling or regeneration, high density, low porosity, high crystallinity, high-order prismatic structure, low organic matter content, great post-mortem stability	Millions of years	Discrete growth period (from in utero time up to ca. 20 years of age for the third molar), representing an average of the several years of enamel formation + may be affected by ‘old-Sr’ available in the body (reservoir effect)	Relatively immune	$\delta^{13}\text{C}$	Approx. +14‰	Whole diet i.e. all dietary macronutrients/energy components (incl. components without proteins)
						$\delta^{18}\text{O}$	Variable depending on boby mass, physiological processes, and drinker type	Drinking and food water
						$^{87}\text{Sr}/^{86}\text{Sr}$	Negligible	Plant food and water; food high in Ca and/or Sr
						$\delta^{13}\text{C}$	Approx. +14‰	Whole diet i.e. all dietary macronutrients/energy components (incl. components without proteins)
Dental enamel sub-sample				Discrete growth period of a tooth layer: depending on the method: a few to a couple of months (if assuming that there is no or few impact of old Sr) or even weekly resolution		$\delta^{18}\text{O}$	Variable depending on boby mass, physiological processes, and drinker type	Drinking and food water
						$^{87}\text{Sr}/^{86}\text{Sr}$	Negligible	Plant food and water; food high in Ca and/or Sr
						$\delta^{13}\text{C}$	Approx. +14‰	Non-calciend: maybe none, maybe only reflects burial soil or local biosphere; calciend: not preserved
						$\delta^{18}\text{O}$	Variable depending on boby mass, physiological processes, and drinker type	Non-calciend: maybe none, maybe only reflects burial soil or local biosphere; calciend: not preserved
Bone bioapatite	Biomineral	High porosity, low crystallinity, highly substituted, high organic matter content, slow formation and remodelling turnover (rate depending on the bone type, might be much shorter than for collagen: maybe seasonal)	(millions of years in arid environments)	Either a long-term average of the last several years or decades of life (depending on age at death) or no lifespan at all (in case of contamination)	Strongly affected (although regarding $\delta^{18}\text{O}$, bioapatite phosphate is much more resistant than carbonate)	$\delta^{13}\text{C}$	Approx. +14‰	Non-calciend: maybe none, maybe only reflects burial soil or local biosphere; calciend: not preserved
						$\delta^{18}\text{O}$	Variable depending on boby mass, physiological processes, and drinker type	Non-calciend: maybe none, maybe only reflects burial soil or local biosphere; calciend: not preserved
						$^{87}\text{Sr}/^{86}\text{Sr}$	Negligible	Non-calciend: reflects mixed, undifferentiable signal of burial soil and plant food and water; calciend: plant food and water; food high in Ca and/or Sr
Dentine collagen	Proteinaceous, robust biomolecule	Grows incrementally, negligible remodelling, but secondary and tertiary dentine form later and dentinal tubules keep providing nutrients	Over 100,000 years	Discrete growth period (from in utero time up to ca. 20 years of age for the third molar) representing an average of the few years of dentine formation + issue of secondary and tertiary dentine	Affected	$\delta^{13}\text{C}$	Up to +5‰ diet-to-collagen, ca. 1‰ collagen-to-collagen	Protein components of the diet
						$\delta^{15}\text{N}$	Between +3 and +5‰ for each trophic level	
						$\delta^{18}\text{O}$	Variable depending on boby mass, physiological	Structural oxygen in food, food water and drinking water

Table 1 (continued) | Overview of the main bioarchaeological samples and analysed tissues and their potential with respect to diverse isotope systems (‘: When considering no particular diet such as low- or high-protein diet)

Sample type	Tissue type	Tissue properties	Maximal age of sample	Individuals lifespan represented	Susceptibility to diagenesis	Measurements	Fractionation between diet and tissue	part of the diet ¹ (mainly reflected)
Bone collagen	Proteinaceous, robust biomolecule	High organic matter content, slow formation and remodelling turnover (rate depending on the bone type: few years up to several decades)	Over 100,000 years	Long-term average of the last several years or decades of life (depending on age at death)	Relatively immune	$\delta^{34}\text{S}$	processes, and drinker type Between -1.8 and $+0.7\%$	Protein components of the diet (especially the methionine, i.e. only a small section of the larger amino-acids pool)
						$\delta^{13}\text{C}$	Up to $+5\%$ diet-to-collagen, ca. 1%	Protein components of the diet
						$\delta^{15}\text{N}$	Between $+3$ and $+5\%$ for each trophic level	
						$\delta^{18}\text{O}$	Variable depending on body mass, physiological processes, and drinker type	Structural oxygen in food, food water and drinking water
						$\delta^{34}\text{S}$	Between -1.8 and $+0.7\%$	Protein components of the diet (especially the methionine, i.e. only a small section of the larger amino-acids pool)

Whenever the site location and its surroundings are still accessible, plant sampling from landscapes with little anthropogenic impact has become the mainstay for $^{87}\text{Sr}/^{86}\text{Sr}$ isoscapes built on primary data^{115,129}. Environmental alternatives for $^{87}\text{Sr}/^{86}\text{Sr}$ include surface water sources¹³⁰, soils and soil leachates¹³¹ and land molluscs⁶⁵. However, concerns have been raised in relation to contamination and the degree to which these archives are consistently representative of the bioavailable Sr of the local area¹³². As the base of the food chain, and since their local provenance can be assured, plants have many benefits for isoscape production¹¹⁵. However, they do not provide a homogenised signature for an area and are susceptible to extreme values, which are generally most likely in shallow rooted plants. Therefore, dense sampling is advantageous and assessing results from shallow, medium and deep-rooted plants is a good approach to characterise variability¹³³. Homogenising plant samples from the same locality provides an alternative, cost-effective approach¹³⁴, but reduces resolution to some degree.

Some diversity in taxa is of value due to plant-specific environmental processes that can impact local values (e.g. the forest-effect on $^{87}\text{Sr}/^{86}\text{Sr}$). Plants are also used for $\delta^{34}\text{S}$ mapping¹¹⁴, but can often be subject to the impact of modern pollution¹³⁵, though this problem is much reduced in many areas due to clean air legislation⁵⁴. Faunal sources such as sheep wool¹³⁶ or archaeological bone⁵⁴ provide alternatives. Archaeological fauna is a good alternative in areas where pollution remains an issue as long as there is strong evidence for local origin. Therefore, small-bodied wild mammals (such as small rodents) with a limited home range are a good choice¹³⁷, though these are generally sparse in faunal assemblages. Because the currently only global $\delta^{34}\text{S}$ isoscape is based on bioarchaeological remains¹¹⁸, which may show the limits mentioned above, it is advisable to complete it with other samples such as plants¹¹⁴ to test for the local reliability—although the non-local and diagenetically altered samples were sorted out before producing the isoscape.

Using larger mammals, whether modern or ancient, provides homogenised values and helps to characterise a landscape rather than pinpointing extreme values¹³⁸. They moreover sometimes represent the only available resource, especially at sites within modern urban areas or which are not accessible anymore due to building activities or some other restrictions. However, using this material requires a dialogue with experts on this archaeo(zoo)logical context, as animals may have had substantial grazing ranges, or may have been involved in transhumance practices and carcass parts found on sites may have been transported from elsewhere by humans, meaning they would not represent local bioavailability. In this context, using environmental analyses to model the site's catchment area is an efficient way to determine the extent of environmental similarities or differences represented by the faunal sample compared to other proxies^{24,139}.

Using bone or dentine in archaeological animals for $^{87}\text{Sr}/^{86}\text{Sr}$ is further ill-advised as these results represent a blend of biogenic (life) and diagenetic (burial) and are therefore non diagnostic¹⁴⁰. In bioarchaeology, infants also have some potential for providing homogenised local values¹⁴¹, provided that early mobility can be excluded. $\delta^{18}\text{O}$ isoscapes are invariably generated on baseline water sources, typically precipitation and/or groundwater^{116,142} as these sources dominate the O in organisms and the complexities of variable fractionation mean other archives are problematic. However, these are also subject to environment-specific impacts, for example through variable fractionation effects associated with evaporation depending on the depth of surface waters¹⁴², and the results may vary depending on the conversion equation used to compare tissue, drinking, and environmental water¹⁴³.

Bioarchaeological isoscapes in ecological and palaeo-research

The geographical scale of isoscapes generated in bioarchaeological research depends strongly on the scope of the study, spanning from the site-specific or micro-regional^{154,139} to the regional and supra-regional^{10,129,144} or the global scale¹⁴⁵. In addition, regional Sr isoscapes are getting increasingly produced with the only goal of serving as isotope background for future research, reflecting the main geological and environmental units of the area¹²⁹, yet remaining a first step for bioarchaeological—and ecological or forensic—

investigations. Depending on the research question, denser sampling might be required to complete this first approximation of the overall isotope diversity within the targeted area.

The geographical scale has in turn an impact on the resolution of the isoscape. At a reduced geographical scale, the isotope diversity of the local environment can be as detailed as to represent every ecological unit within a site's catchment area using an important diversity of samples and sampling locations^{53,146}—although bioarchaeological isoscapes often target places suitable for anthropogenic land-use^{17,139}. On the other hand, large-scale isoscapes rather interpolate more scattered measurements based on the geological background, topography, hydrology, distance to the coast, or any relevant variable^{10,144,145}. Importantly, local isotope values do not necessarily match the values predicted by global isoscapes (e.g. Ireland^{129,145}), because increasing the geographical area often leads to a loss of resolution among the parameters used for the interpolation, while the relevant environmental parameters vary between ecosystems and local specificities thus cannot be considered at the large scale (Fig. 2). In this context, bioarchaeology contributes significantly to characterising the isotope variability in context- and region-specific environmental conditions. In this context, it is essential to create models that integrate multiple environmental and climatic variables to account for all factors influencing isotope variability at the investigated scale^{17,139,147}. For more transparency in the interpolation process, isoscapes (of mean or median values) are meanwhile frequently published together with a map showing the error ranges¹²⁹.

By integrating bioarchaeological isoscape data in palaeo-research, the discrepancy in terms of geographical and temporal resolution of the various proxies may require the use of categorised data or average values, which necessarily leads to a loss of information. In this context, it is worth engaging a cross-disciplinary discussion regarding targeted processes and gaps that need to be filled in terms of isoscapes, to determine which proxies and sample types are required and suitable. Yet with the large diversity of proxies derived from all disciplines as well as their great temporal and geographical dimension, it may be possible to disentangle locally from globally driven triggers of isotope variability within various ecological processes. It would be also important to discuss issues related to disturbed environments and their impact on local isotope composition.

For example, in temperate zones such as the UK areas of high precipitation in the west, the bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ is shifted from the geological value towards the value of rainfall¹⁴⁶. Similarly, contexts of marine inundation, seaspray (causing increasing values) and waterlogging (generally reducing values), altering bioavailable $\delta^{34}\text{S}$, $\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ ^{52–54}, require specific attention. And since the deposition of aeolian dust can have a major impact in altering bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ away from underlying lithology¹³², bioarchaeological samples may track previous stages of the landscape. Such an attempt has been done in the Peak District, England, where the considerable change in $^{87}\text{Sr}/^{86}\text{Sr}$ over time may be due to the expansion of blanket peat and/or of leaching¹⁴⁸. A bioarchaeological study in Sweden further demonstrated the potential variation of $^{87}\text{Sr}/^{86}\text{Sr}$ over time in areas affected by melting glaciers¹⁴⁹.

Formal isoscapes tend not to be developed for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ due to the manifold factors dictating variability. However, there is evidence that cultural practices such as dietary habits (e.g. marine versus terrestrial food, C_3 versus C_4 crops, high versus low-protein diet) or agricultural strategies (e.g. manuring) as well as geographical, environmental and climatic settings such as latitude, lithology or moisture have an impact on these isotope ratios, with archaeological fauna providing a useful archive to explore large-scale patterns^{35,37,61,150}. The exponential amount of isotope data produced in bioarchaeology thus allows the investigation of changes in ecological, geological, and geomorphological processes in the long-term and recent studies have taken advantage of both the diachronic perspective and the large geographical distribution of bioarchaeological isotope data to determine new indicators for large-scale evolution of environmental conditions^{35,40,151}. By combining bioarchaeological isotope data with further approaches and proxies^{152,153}, there might be, for instance, potential to investigate changes in forest cover using the canopy effect¹⁵⁴.

This opportunity to translate the huge amount of isotope data into isoscapes is further useful for the investigation of habitats and animal ecologies¹⁵⁵. It can also be used to track mobility patterns, because specific geographically constrained environmental conditions, agricultural practices, or dietary habits produce typical isotopic signatures—especially when considering multiple isotopic systems^{150,156}. It may be a challenge, however, to correlate changes in isotope baselines with changes in environmental conditions, especially when one needs to consider and quantify a potential anthropogenic impact on the isotope variability – including manuring fields or consuming boiled, stewed or fermented food and drinks, which can considerably alter the original local stable isotope ratio^{75,157}. A first step in this direction would be to constrain the research area in terms of period and region, in which we can rule out some factors of variability (targeting especially periods at which the human impact was absent or negligible), to focus on the herbivores (to avoid animals with more flexible dietary habits), and to integrate different proxies, although it remains key to consider the potential effect of both mobility and seasonality on isotope variability¹⁵⁸ as well as the time lag and varying isotope accumulation rate related to tissue formation¹⁵⁹.

Conclusions and perspectives

Bioarchaeological research has achieved considerable advances in the understanding of past human and animal behaviours and how this has shaped or impacted on the environment. It hence actively contributed to palaeoenvironmental and ecological research, yet this contribution is mainly restricted to the acknowledgment of resulting narratives and created knowledge. On the contrary, the tremendous amount of isotope data derived from a large diversity of bioarchaeological samples and contexts remains underexploited. With this paper, we present potential and limits of integrating bioarchaeological isotope data into other palaeo-disciplines and identify future directions and strategies enabled by a cross-disciplinary collaboration, starting from the research designs and models.

The huge diversity of sampled tissues and materials provides complementary information about various aspects of past ecologies and environments and can contribute to track processes such as precipitation patterns, climate change, or the evolution of palaeoenvironmental conditions in regions where other proxies are not available. In this context, there is an urgent need to start integrating these data into existing models to test how they can be used, for instance, in predictive/inductive models. More importantly, the large geographical distribution of bioarchaeological data and their deep chronology allow for tracking large-scale and long-term processes. In turn, the diversity and density of the existing dataset make it possible to question the impact of large-scale processes as well as of short-term or long-term changes at the local or regional scale. As the data is easily related to human activity, one can address questions related to human-environment interactions over the long term.

In this context, it is paramount to recognise the complexity derived from bioarchaeological research, and to consider the different meanings of the various samples, to accurately deal with uncertain or imprecise chronologies, and to acknowledge the bias in the sample related to archaeological research procedures. This paper offers a wide spectrum of answers to these issues. It would, however, be of great value for greater engagement between bioarchaeologists and other palaeo-scientists to foster collaboration and design joint research programs that benefit from analytical advances and interdisciplinary methodological transfer. There is a great potential in targeting common research areas of interest, identifying gaps in terms of chronology, location or types of data/proxies (i.e. the types of sampled tissue/material), creating joint monitoring programs, and testing for instance physiological, climatic, ecological, and environmental models. Based on this, one could adapt future sampling procedures and hence produce data and proxies as meaningful and reusable as possible for the various disciplines.

Received: 27 March 2025; Accepted: 20 June 2025;

Published online: 01 July 2025

References

1. Vogel, J. C. & van der Merwe, N. J. Isotopic evidence for early maize cultivation in New York State. *Am. Antiq.* **42**, 238–242 (1977).
2. Ericson, J. E. Strontium isotope characterization in the study of prehistoric human ecology. *J. Hum. Evol.* **14**, 503–514 (1985).
3. Ambrose, S. H. Isotopic analysis of paleodiets: methodological and interpretative considerations. In *Investigations of Ancient Human Tissue. Chemical Analyses in Anthropology* (ed. Sandford, M.K.) 59–130 (Gordon and Breach, 1993).
4. Lee-Thorp, J. A. On isotopes and old bones. *Archaeometry* **50**, 925–950 (2008).
5. Bentley, R. A. Strontium Isotopes from the Earth to the archaeological skeleton: a review. *J. Archaeol. Method Theory* **13**, 135–187 (2006).
6. Nehlich, O. The application of sulphur isotope analyses in archaeological research: a review. *Earth Sci. Rev.* **142**, 1–17 (2015).
7. Reynard, L. M. & Hedges, R. Stable hydrogen isotopes of bone collagen in palaeodietary and palaeoenvironmental reconstruction. *J. Archaeol. Sci.* **35**, 1934–1942 (2008).
8. Guiserix, D. et al. Simultaneous analysis of stable and radiogenic strontium isotopes in reference materials, plants and modern tooth enamel. *Chem. Geol.* **606**, 121000 (2022).
9. Junqueira, T. P. et al. Applications of zinc stable isotope analysis in environmental and biological systems: a review. *GEEA* **24**; <https://doi.org/10.1144/geochem2024-003> (2024).
10. Evans, J. A. et al. Applying lead (Pb) isotopes to explore mobility in humans and animals. *PLoS ONE* **17**, e0274831 (2022).
11. Plomp, E. Neodymium isotopes in modern human dental enamel: an exploratory dataset for human provenancing. *Data Brief* **38**, 107375 (2021).
12. Evershed, R. P., Dudd, S. N., Copley, M. S. & Mutherjee, A. Identification of animal fats via compound specific $\delta^{13}\text{C}$ values of individual fatty acids: assessments of results for reference fats and lipid extracts of archaeological pottery vessels. *Doc. Praeh.* **29**, 73–96 (2002).
13. Depaermentier, M. L. C., Kempf, M., Bánffy, E. & Alt, K. W. Tracing mobility patterns through the 6th–5th millennia BC in the Carpathian Basin with strontium and oxygen stable isotope analyses. *PLoS ONE* **15**, e0242745 (2020).
14. Reade, H., O'connell, T. C., Barker, G. & Stevens, R. E. Increased climate seasonality during the late glacial in the Gebel Akhdar, Libya. *Quat. Sci. Rev.* **192**, 225–235 (2018).
15. France, C. A., Qi, H. & Kavich, G. M. Combined influence of meteoric water and protein intake on hydrogen isotope values in archaeological human bone collagen. *J. Archaeol. Sci.* **96**, 33–44 (2018).
16. Jaouen, K., Beasley, M., Schoeninger, M., Hublin, J.-J. & Richards, M. P. Zinc isotope ratios of bones and teeth as new dietary indicators: results from a modern food web (Koobi Fora, Kenya). *Sci. Rep.* **6**, 26281 (2016).
17. Depaermentier, M. L. C., Kempf, M., Bánffy, E. & Alt, K. W. Cultural diversity shaped neolithic subsistence in the Carpathian Basin. *Sci. Rep.* **15**, 4281 (2025).
18. Alexander, M. M., Gerrard, C. M., Gutiérrez, A. & Millard, A. R. Diet, society, and economy in late medieval Spain: stable isotope evidence from Muslims and Christians from Gandía, Valencia. *Am. J. Phys. Anthropol.* **156**, 263–273 (2015).
19. van der Sluis, L. G., Reimer, P. J. & Ogle, N. Adding hydrogen to the isotopic inventory—combining $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^2\text{H}$ stable isotope analysis for palaeodietary purposes on archaeological bone. *Archaeometry* **61**, 720–749 (2019).
20. Pearson, J. & Grove, M. Counting sheep: sample size and statistical inference in stable isotope analysis and palaeodietary reconstruction. *World Archaeol.* **45**, 373–387 (2013).

21. Pederzani, S. et al. Stable isotopes show *Homo sapiens* dispersed into cold steppes ~45,000 years ago at Ilsehöhle in Ranis, Germany. *Nat. Ecol. Evol.* **8**, 578–588 (2024).
22. Gruchy et al. Bone of contention: intra-element variability in remodelling of human femora based on histomorphometric and isotope analyses. *PLoS ONE* **19**, e0305089 (2024).
23. Leggett, S., Rose, A., Praet, E. & Le Roux, P. Multi-tissue and multi-isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$ and 87/86 Sr) data for early medieval human and animal palaeoecology. *Ecology* **102**, e03349 (2021).
24. Depaermentier, M. L. C. et al. Bioarchaeological analyses reveal long-lasting continuity at the periphery of the late antique Roman Empire. *iScience* **26**, 107034 (2023).
25. Esposito, C. et al. NOthing goes to WASTE (NOWA): a protocol to optimise sampling of ancient teeth. *J. Archaeol. Sci.* **171**, 106087 (2024).
26. Pilaar Birch, S. E. & Graham, R. W. A Stable isotope data repository as part of neotoma, a paleoecological database. *BioScience* **65**, 953 (2015).
27. Salesse, K. et al. IsoArch.eu: an open-access and collaborative isotope database for bioarchaeological samples from the Graeco-Roman world and its margins. *J. Archaeol. Sci. Rep.* **19**, 1050–1055 (2018).
28. Shipley, O. N. et al. Design, development, and implementation of IsoBank: a centralized repository for isotopic data. *PLoS ONE* **19**, e0295662 (2024).
29. Ebert, C. E. et al. The Caribbean and Mesoamerica Biogeochemical Isotope Overview (CAMBIO). *Sci. Data* **11**, 349 (2024).
30. Billings, T. N. et al. The North American repository for archaeological isotopes. *Sci. Data* **12**, 50 (2025).
31. Wang, T. et al. Tianshanbeilu and the Isotopic millet road: reviewing the late neolithic/bronze age radiation of human millet consumption from north China to Europe. *Natl. Sci. Rev.* **6**, 1024–1039 (2019).
32. Leggett, S. A hierarchical meta-analytical approach to western European dietary transitions in the first millennium AD. *Eur. J. Archaeol.* 1–21; <https://doi.org/10.1017/ea.2022.23> (2022).
33. van Klinken, G. J., Richards, M. P. & Hedges, R. E. M. An overview of causes for stable isotopic variations in past European human populations. environmental, ecophysiological, and cultural effects. In *Biogeochemical Approaches to Paleodietary Analysis. Advances in Archaeological and Museum Science* (eds Ambrose, S.H. & Katzenberg, M.A.) 39–63 (New York, London, 2002).
34. Drucker, D. G. et al. Ecology of large ungulates in the northeastern Iberian Peninsula during the Upper Palaeolithic through stable isotopes and tooth wear analysis. *Quat. Environ. Hum.* **2**, 100011 (2024).
35. Reade, H. et al. Nitrogen palaeo-isoscapes: changing spatial gradients of faunal $\delta^{15}\text{N}$ in late Pleistocene and early Holocene Europe. *PLoS ONE* **18**, e0268607 (2023).
36. Britton, K. et al. Multi-isotope analysis of bone collagen of Late Pleistocene ungulates reveals niche partitioning and behavioural plasticity of reindeer during MIS 3. *Sci. Rep.* **13**, 15722 (2023).
37. van Klinken, G. J., van der Plicht, J. & Hedges, R. E. M. Bone $^{13}\text{C}/^{12}\text{C}$ ratios reflect (palaeo) climatic variations. *Geophys. Res. Lett.* **21**, 445–448 (1994).
38. Pederzani, S. et al. Subarctic climate for the earliest *Homo sapiens* in Europe. *Sci. Adv.* **7**, eabi4642 (2021).
39. Drucker, D. G., Bocherens, H. & Billiou, D. Evidence for shifting environmental conditions in Southwestern France from 33,000 to 15,000 years ago derived from carbon-13 and nitrogen-15 natural abundances in collagen of large herbivores. *Earth Planet. Sci. Lett.* **216**, 163–173 (2003).
40. Stevens, R. et al. Major excursions in sulfur isotopes linked to permafrost change in Eurasia during the last 50,000 Years. *Research Square*; <https://doi.org/10.21203/rs.3.rs-2556240/v1> (2023).
41. Crowley, B. E. & Godfrey, L. R. Strontium isotopes support small home ranges for extinct lemurs. *Front. Ecol. Evol.* **7**; <https://doi.org/10.3389/fevo.2019.00490> (2019).
42. Drucker, D. G. et al. Tracking possible decline of woolly mammoth during the Gravettian in Dordogne (France) and the Ach Valley (Germany) using multi-isotope tracking (^{13}C , ^{14}C , ^{15}N , ^{34}S , ^{18}O). *Quat. Int.* **359–360**, 304–317 (2015).
43. Pilaar Birch, S. E., Miracle, P. T., Stevens, R. E. & O'Connell, T. C. Late Pleistocene/Early Holocene migratory behavior of ungulates using isotopic analysis of tooth enamel and its effects on forager mobility. *PLoS ONE* **11**, e0155714 (2016).
44. Baumann, C. et al. Dietary niche partitioning among Magdalenian canids in southwestern Germany and Switzerland. *Quat. Sci. Rev.* **227**, 106032 (2020).
45. Kveiborg, J. et al. Paleodiet reconstructions and human utilization of middle Holocene *Equus ferus* in northwest Europe. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **649**, 112334 (2024).
46. Richards, M. P. et al. Isotopic evidence for omnivory among European cave bears: late Pleistocene *Ursus spelaeus* from the Peștera cu Oase, Romania. *PNAS* **105**, 600–604 (2008).
47. Bocherens, H. et al. Isotopic evidence for dietary ecology of cave lion (*Panthera spelaea*) in North-Western Europe: prey choice, competition and implications for extinction. *Quat. Int.* **245**, 249–261 (2011).
48. Robinson, J. R., Rowan, J., Barr, W. A. & Sponheimer, M. Intrataxonomic trends in herbivore enamel $\delta^{13}\text{C}$ are decoupled from ecosystem woody cover. *Nat. Ecol. Evol.* **5**, 995–1002 (2021).
49. Pederzani, S. & Britton, K. Oxygen isotopes in bioarchaeology: principles and applications, challenges and opportunities. *Earth Sci. Rev.* **188**, 77–107 (2019).
50. Stevens, R. E., Pederzani, S., Britton, K. & Wexler, S. Bones and teeth isotopes as archives for palaeoclimatic, palaeoenvironmental and palaeoecological data. *Quat. Sci. Rev.* **357**, 109320 (2025).
51. Hedges, R. E., Stevens, R. E. & Richards, M. Bone as a stable isotope archive for local climatic information. *Quat. Sci. Rev.* **23**, 959–965 (2004).
52. Göhring, A., Hölzl, S., Mayr, C. & Strauss, H. Identification and quantification of the sea spray effect on isotopic systems in α -cellulose ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$), total sulfur ($\delta^{34}\text{S}$), and 87Sr/86Sr of European beach grass (*Ammophila arenaria*, L.) in a greenhouse experiment. *Sci. Total Environ.* **856**, 158840 (2023).
53. Göhring, A., Hölzl, S., Mayr, C. & Strauss, H. Multi-isotope fingerprints of recent environmental samples from the Baltic coast and their implications for bioarchaeological studies. *Sci. Total Environ.* **874**, 162513 (2023).
54. Lamb, A. L., Chenery, C. A., Madgwick, R. & Evans, J. A. Wet feet: developing sulfur isotope provenance methods to identify wetland inhabitants. *R. Soc. Open Sci.* **10**, 230391 (2023).
55. Bocherens, H., Drucker, D. G. & Madelaine, S. Evidence for a (^{15}N) positive excursion in terrestrial foodwebs at the middle to upper Palaeolithic transition in south-western France: implications for early modern human palaeodiet and palaeoenvironment. *J. Hum. Evol.* **69**, 31–43 (2014).
56. Jones, J. R., Marín-Arroyo, A. B., Straus, L. G. & Richards, M. P. Adaptability, resilience and environmental buffering in European Refugia during the late Pleistocene: insights from La Riera Cave (Asturias, Cantabria, Spain). *Sci. Rep.* **10**, 1217 (2020).
57. Drucker, D. G. The isotopic ecology of the mammoth steppe. *Annu. Rev. Earth Planet. Sci.* **50**, 395–418 (2022).
58. Drucker, D. G., Kind, C.-J. & Stephan, E. Chronological and ecological information on Late-glacial and early Holocene reindeer from northwest Europe using radiocarbon (^{14}C) and stable isotope (^{13}C , ^{15}N) analysis of bone collagen: case study in southwestern Germany. *Quat. Int.* **245**, 218–224 (2011).

59. Stevens, R. E. et al. Nitrogen isotope analyses of reindeer (*Rangifer tarandus*), 45,000 BP to 9,000 BP: palaeoenvironmental reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **262**, 32–45 (2008).
60. Drucker, D. G., Madelaine, S. & Morala, A. Les derniers rennes de Dordogne. *Paleo*, 85–100; <https://doi.org/10.4000/paleo.2087> (2011).
61. Stevens, R. E. & Hedges, R. E. Carbon and nitrogen stable isotope analysis of northwest European horse bone and tooth collagen, 40,000BP–present: Palaeoclimatic interpretations. *Quat. Sci. Rev.* **23**, 977–991 (2004).
62. Drucker, D. G., Bridault, A. & Cupillard, C. Environmental context of the Magdalenian settlement in the Jura Mountains using stable isotope tracking (^{13}C , ^{15}N , ^{34}S) of bone collagen from reindeer (*Rangifer tarandus*). *Quat. Int.* **272–273**, 322–332 (2012).
63. Drucker, D. G. et al. Collagen stable isotopes provide insights into the end of the mammoth steppe in the central East European plains during the Epigravettian. *Quat. Res.* **90**, 457–469 (2018).
64. Reiss, L. et al. Changing food webs before and during the Last Glacial Maximum based on stable isotopes of animal bone collagen from Lower Austria. *J. Quat. Sci.* **38**, 1337–1356 (2023).
65. Prendergast, A. L. & Stevens, R. E. Molluscs (isotopes): analyses in environmental archaeology. In *Encyclopedia of Global Archaeology* (ed. Smith, C.) 7332–7340 (Springer International Publishing, 2020).
66. Sandweiss, D. H. et al. Archaeological climate proxies and the complexities of reconstructing Holocene El Niño in coastal Peru. *PNAS* **117**, 8271–8279 (2020).
67. Vaiglova, P. et al. Past rainfall patterns in Southeast Asia revealed by microanalysis of $\delta^{18}\text{O}$ values in human teeth. *J. Archaeol. Sci.* **162**, 105922 (2024).
68. Tian, Y. et al. Holocene climate change in southern Oman deciphered by speleothem records and climate model simulations. *Nat. Commun.* **14**, 4718 (2023).
69. Pederzani, S. et al. Reconstructing Late Pleistocene paleoclimate at the scale of human behavior: an example from the Neandertal occupation of La Ferrassie (France). *Sci. Rep.* **11**, 1419 (2021).
70. Luterbacher, J. & Zorita, E. Analysis and Interpretation: spatial climate field reconstructions. In *The Palgrave Handbook of Climate History* (eds White, S., Pfister, C. & Mauelshagen, F.) 131–139 (Palgrave Macmillan UK, 2018).
71. Suraprasit, K. et al. Long-term isotope evidence on the diet and habitat breadth of Pleistocene to Holocene Caprines in Thailand: implications for the extirpation and conservation of Himalayan Goats. *Front. Ecol. Evol.* **8**; <https://doi.org/10.3389/fevo.2020.00067> (2020).
72. Büntgen, U. et al. The influence of decision-making in tree ring-based climate reconstructions. *Nat. Commun.* **12**, 3411 (2021).
73. Fleitmann, D. et al. Palaeoclimatic interpretation of high-resolution oxygen isotope profiles derived from annually laminated speleothems from Southern Oman. *Quat. Sci. Rev.* **23**, 935–945 (2004).
74. Pederzani, S. et al. Late Pleistocene Neandertal exploitation of stable and mosaic ecosystems in northern Iberia shown by multi-isotope evidence. *Quat. Res.* **116**, 108–132 (2023).
75. Larsson, M., Bergman, J. & Olsson, P. A. Soil, fertilizer and plant density: exploring the influence of environmental factors to stable nitrogen and carbon isotope composition in cereal grain. *J. Archaeol. Sci.* **163**, 105935 (2024).
76. Bogaard, A., Heaton, T., Poulton, P. & Merbach, I. The impact of manuring on nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and crop management practices. *J. Archaeol. Sci.* **34**, 335–343 (2007).
77. Shahack-Gross, R., Simons, A. & Ambrose, S. H. Identification of pastoral sites using stable nitrogen and carbon isotopes from bulk sediment samples: a case study in modern and archaeological pastoral settlements in Kenya. *J. Archaeol. Sci.* **35**, 983–990 (2008).
78. Diaz, L. Unraveling Ancient Andean Camelid Herding Strategies Using Multi-Isotope Analyses. In *Arts & Sciences Electronic Theses and Dissertations*. Vol. 3335, <https://doi.org/10.7936/ktkp-h909> (2024).
79. Melton, M. A. et al. Reconstructing middle horizon camelid diets and foddering practices: microbotanical and isotope analyses of dental remains from Quilcapampa, Peru. *Lat. Am. Antiq.* **34**, 783–803 (2023).
80. Somerville, A. D. Human-animal interactions in the pre-colonial Americas: insights from stable carbon isotope analysis. In *Exploring Human Behavior Through Isotope Analysis* (eds Beasley, M.M. & Somerville, A.D.) 181–205 (Springer International Publishing, 2023).
81. Britton, K., Müldner, G. & Bell, M. Stable isotope evidence for salt-marsh grazing in the Bronze Age Severn Estuary, UK: implications for palaeodietary analysis at coastal sites. *J. Archaeol. Sci.* **35**, 2111–2118 (2008).
82. Jones, J. R. & Mulville, J. Isotopic and zooarchaeological approaches towards understanding aquatic resource use in human economies and animal management in the prehistoric Scottish North Atlantic Islands. *J. Archaeol. Sci. Rep.* **6**, 665–677 (2016).
83. Gillis, R. E. et al. Diverse prehistoric cattle husbandry strategies in the forests of Central Europe. *Nat. Ecol. Evol.* **9**, 87–98 (2025).
84. Szabó, P. et al. Pliocene—early Pleistocene continental climate and vegetation in Europe based on stable isotope compositions of mammal tooth enamel. *Quat. Sci. Rev.* **288**, 107572 (2022).
85. Nguy, W. H. & Secord, R. Middle Miocene paleoenvironmental reconstruction in the central Great Plains, USA, from stable carbon isotopes in ungulates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **594**, 110929 (2022).
86. Cullen, T. M., Zhang, S., Spencer, J. & Cousens, B. Sr-O-C isotope signatures reveal herbivore niche-partitioning in a Cretaceous ecosystem. *Palaeontology* **65**; <https://doi.org/10.1111/pala.12591> (2022).
87. Samec, C. T. & Yacobaccio, H. D. Chinchillidae exploitation during the first half of the Holocene in the Argentinian Puna: a contribution from zooarchaeology and stable isotope analysis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **648**, 112298 (2024).
88. Berto, C., Arnaud, J., López-García, J. M., Luzi, E. & Arzarello, M. Analysis of the early Pleistocene small mammals from Pirro Nord 13 (Apricena, southern Italy) and their implications for reconstructing the palaeoenvironment of the early human occupation in Europe. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **647**, 112251 (2024).
89. Chidimuro, B. et al. North and South: exploring isotopic analysis of bone carbonates and collagen to understand post-medieval diets in London and northern England. *Am. J. Biol. Anthropol.* **182**, 126–142 (2023).
90. Stevens, R. E., Lightfoot, E., Hamilton, J., CUNLIFFE, B. & Hedges, R. E. Stable isotope investigations of the Danebury hillfort pit burials. *Oxf. J. Archaeol.* **29**, 407–428 (2010).
91. Norwood, A. L., Wang, B. & Kingston, J. D. Linking African herbivore community enamel isotopes and environments: challenges, opportunities, and paleoecological implications. *Oecologia* **204**, 467–489 (2024).
92. Fritts, H. C. *Tree Rings and Climate* (Elsevier, 1976).
93. Zong, X. et al. Climatic significance of modern minute land snail shells $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ on the Chinese Loess Plateau. *Ecol. Indic.* **145**, 109733 (2022).
94. Leng, M. J. & Marshall, J. D. Palaeoclimate interpretation of stable isotope data from lake sediment archives. *Quat. Sci. Rev.* **23**, 811–831 (2004).
95. Dee, S. et al. Water isotopes, climate variability, and the hydrological cycle: recent advances and new frontiers. *Environ. Res. Clim.* **2**, 22002 (2023).

96. Konecny, B. L. et al. The Iso2k database: a global compilation of paleo- $\delta^{18}\text{O}$ and $\delta^2\text{H}$ records to aid understanding of common era climate. *Earth Syst. Sci. Data* **12**, 2261–2288 (2020).
97. Reynard, L. M. et al. Mediterranean precipitation isoscape preserved in bone collagen $\delta^2\text{H}$. *Sci. Rep.* **10**, 8579 (2020).
98. Britton, K. et al. Oxygen isotope analysis of Equus teeth evidences early Eemian and early Weichselian palaeotemperatures at the Middle Palaeolithic site of Neumark-Nord 2, Saxony-Anhalt, Germany. *Quat. Sci. Rev.* **226**, 106029 (2019).
99. Pryor, A. J., Stevens, R. E., O'connell, T. C. & Lister, J. R. Quantification and propagation of errors when converting vertebrate biomineral oxygen isotope data to temperature for palaeoclimate reconstruction. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **412**, 99–107 (2014).
100. Sharp, Z. D. & Cerling, T. E. Fossil isotope records of seasonal climate and ecology: straight from the horse's mouth. *Geology* **26**, 219 (1998).
101. Hajdas, I. Radiocarbon dating and its applications in quaternary studies; <https://doi.org/10.23689/FIDGEO-988> (2008).
102. Gliganic, L. A., McDonald, J. & Meyer, M. C. Luminescence rock surface exposure and burial dating: a review of an innovative new method and its applications in archaeology. *Archaeol. Anthropol. Sci.* **16**; <https://doi.org/10.1007/s12520-023-01915-0> (2024).
103. Grechko, D. Western frontier of the Archaic Scythia: typo-chronology vs radiocarbon dating. *Spraw. Arch.* **75**, 405–436 (2023).
104. Rodríguez Ramos, R., Rodríguez López, M. & Pestle, W. J. Revision of the cultural chronology of precolonial Puerto Rico: a Bayesian approach. *PLoS ONE* **18**, e0282052 (2023).
105. Wu, X. et al. Early pottery at 20,000 years ago in Xianrendong Cave, China. *Science* **336**, 1696–1700 (2012).
106. Shennan, S. (ed.) *The First Farmers of Europe* (Cambridge University Press, 2018).
107. Potter, B. A. et al. Current understanding of the earliest human occupations in the Americas: evaluation of Becerra-Valdivia and Higham (2020). *PaleoAmerica* **8**, 62–76 (2022).
108. Hajdas, I. et al. Radiocarbon dating. *Nat. Rev. Methods Primers* **1**; <https://doi.org/10.1038/s43586-021-00058-7> (2021).
109. Hines, J. & Bayliss, A. (eds) *Anglo-Saxon Graves and Grave Goods of the 6th And 7th centuries AD: A Chronological Framework* (Society for Medieval Archaeology, 2013).
110. Jacobsson, P. et al. Refining the Hallstatt Plateau: short-term ^{14}C variability and small scale offsets in 50 consecutive single tree-rings from Southwest Scotland Dendro-dated to 510–460 BC. *Radiocarbon* **60**, 219–237 (2018).
111. Loader, N. J. et al. Tree ring dating using oxygen isotopes: a master chronology for central England. *J. Quat. Sci.* **34**, 475–490 (2019).
112. Black, B. A. et al. The revolution of crossdating in marine palaeoecology and palaeoclimatology. *Biol. Lett.* **15**, 20180665 (2019).
113. Savard, M. M. & Daux, V. An overview on isotopic divergences — causes for instability of tree-ring isotopes and climate correlations. *Climate* **16**, 1223–1243 (2020).
114. Tarrant, D. & Richards, M. P. Modern plants and sulfur isoscapes - a review, discussion, and construction of a pilot $\delta^{34}\text{S}$ isoscape for mobility and provenance studies. *Rapid Commun. Mass Spectrom.* **38**, e9908 (2024).
115. Holt, E., Evans, J. A. & Madgwick, R. Strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) mapping: a critical review of methods and approaches. *Earth Sci. Rev.* **216**, 103593 (2021).
116. Bowen, G. J. & Revenaugh, J. Interpolating the isotopic composition of modern meteoric precipitation. *Water Resour. Res.* **39**; <https://doi.org/10.1029/2003WR002086> (2003).
117. Bowen, G. J. Isoscapes: Spatial pattern in isotopic biogeochemistry. *Annu. Rev. Earth Planet. Sci.* **38**, 161–187 (2010).
118. Bataille, C. P. et al. Triple sulfur-oxygen-strontium isotopes probabilistic geographic assignment of archaeological remains using a novel sulfur isoscape of western Europe. *PLoS ONE* **16**, e0250383 (2021).
119. Holder, P. W. et al. Isotopes and trace elements as natal origin markers of *Helicoverpa armigera*-an experimental model for biosecurity pests. *PLoS ONE* **9**, e92384 (2014).
120. Brlik, V. et al. Animal tracing with sulfur isotopes: spatial segregation and climate variability in Africa likely contribute to population trends of a migratory songbird. *J. Anim. Ecol.* **92**, 1320–1331 (2023).
121. Rossmann, A. et al. The potential of multielement stable isotope analysis for regional origin assignment of butter. *Eur. Food Res. Technol.* **211**, 32–40 (2000).
122. Almeida, C. M. & Vasconcelos, M. T. S. D. ICP-MS determination of strontium isotope ratio in wine in order to be used as a fingerprint of its regional origin. *J. Anal. Spectrom.* **16**, 607–611 (2001).
123. Durante, C. et al. An analytical approach to Sr isotope ratio determination in Lambrusco wines for geographical traceability purposes. *Food Chem.* **173**, 557–563 (2015).
124. Johnson, L., Evans, J., Montgomery, J. & Chenery, C. The forest effect: biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ shifts due to changing land use and the implications for migration studies. *Sci. Total Environ.* **839**, 156083 (2022).
125. Silva-Sánchez, N. & Armada, X.-L. Environmental impact of roman mining and metallurgy and its correlation with the archaeological evidence: an European perspective. *Environ. Archaeol.* 1–25; <https://doi.org/10.1080/14614103.2023.2181295> (2023).
126. Thomsen, E. & Andreassen, R. Agricultural lime disturbs natural strontium isotope variations: implications for provenance and migration studies. *Sci. Adv.* **5**, eaav8083 (2019).
127. Blanz, M. et al. Seaweed fertilisation impacts the chemical and isotopic composition of barley: implications for analyses of archaeological skeletal remains. *J. Archaeol. Sci.* **104**, 34–44 (2019).
128. Blanz, M., Gröcke, D. R., Martin, P. & Church, M. J. The effect of seaweed fertilisation on sulfur isotope ratios ($\delta^{34}\text{S}$) and grain size in barley: implications for agronomy and archaeological research. *Front. Environ. Archaeol.* **3**; <https://doi.org/10.3389/fearc.2024.1465082> (2024).
129. Snoeck, C. et al. Towards a biologically available strontium isotope baseline for Ireland. *Sci. Total Environ.* **712**, 136248 (2020).
130. Frei, K. M. & Frei, R. The geographic distribution of strontium isotopes in Danish surface waters – a base for provenance studies in archaeology, hydrology and agriculture. *Appl. Geochem.* **26**, 326–340 (2011).
131. Hoogewerff, J. A. et al. Bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ in European soils: a baseline for provenancing studies. *Sci. Total Environ.* **672**, 1033–1044 (2019).
132. Hartman, G. & Richards, M. Mapping and defining sources of variability in bioavailable strontium isotope ratios in the Eastern Mediterranean. *Geochim. Cosmochim. Acta* **126**, 250–264 (2014).
133. Sengeløv, A. et al. From plants to patterns: constructing a comprehensive online strontium isoscape for Belgium (IsoBel) using high density grid mapping. *Geoderma* **453**, 117123 (2025).
134. Macheridis, S. et al. Preliminary strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$) baselines for the Bjäre Peninsula and Halland in southern Sweden. *Front. Environ. Archaeol.* **3**; <https://doi.org/10.3389/fearc.2024.1379055> (2024).
135. Zhao, F. J., Spiro, B., Poulton, P. R. & McGrath, S. P. Use of sulfur isotope ratios to determine anthropogenic sulfur signals in a grassland ecosystem. *Environ. Sci. Technol.* **32**, 2288–2291 (1998).
136. Zazzo, A. et al. Isotopic composition of sheep wool records seasonality of climate and diet. *Rapid Commun. Mass Spectrom.* **29**, 1357–1369 (2015).
137. Perry, M. A., Coleman, D. & Delhopital, N. Mobility and exile at 2nd century A.D. khirbet edh-dharih: strontium isotope analysis of

- human migration in western Jordan. *Geoarchaeology* **23**, 528–549 (2008).
138. Bentley, R. A. & Knipper, C. Geographical patterns in biologically available strontium, carbon and oxygen isotope signatures in prehistoric SW Germany. *Archaeometry* **47**, 629–644 (2005).
 139. Depaermentier, M. L. C., Kempf, M., Bánffy, E. & Alt, K. W. Modelling a scale-based strontium isotope baseline for Hungary. *J. Archaeol. Sci.* **135**, 1–16 (2021).
 140. Madgwick, R., Lewis, J., Grimes, V. & Guest, P. On the hoof: exploring the supply of animals to the Roman legionary fortress at Caerleon using strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) isotope analysis. *Archaeol. Anthropol. Sci.* **11**, 223–235 (2019).
 141. Esposito, C. et al. Intense community dynamics in the pre-Roman frontier site of Fermo (ninth–fifth century BCE, Marche, central Italy) inferred from isotopic data. *Sci. Rep.* **13**, 3632 (2023).
 142. Darling, W. G., Bath, A. H. & Talbot, J. C. The O and H stable isotope composition of freshwaters in the British Isles. 2. Surface waters and groundwater. *Hydrol. Earth Syst. Sci. Discuss.* **7**, 183–195 (2003).
 143. Pellegrini, M., Lee-Thorp, J. A. & Donahue, R. E. Exploring the variation of the $\delta^{18}\text{O}_\text{p}$ and $\delta^{18}\text{O}_\text{c}$ relationship in enamel increments. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **310**, 71–83 (2011).
 144. Serna, A., Prates, L., Valenzuela, L. O. & Salazar-García, D. C. Back to the bases: building a terrestrial water $\delta^{18}\text{O}$ baseline for archaeological studies in North Patagonia (Argentina). *Quat. Int.* **548**, 4–12 (2020).
 145. Bataille, C. P., Crowley, B. E., Wooller, M. J. & Bowen, G. J. Advances in global bioavailable strontium isoscapes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **555**, 109849 (2020).
 146. Montgomery, J. Passports from the past: investigating human dispersals using strontium isotope analysis of tooth enamel. *Ann. Hum. Biol.* **37**, 325–346 (2010).
 147. Jones, J. & Britton, K. Multi-scale, integrated approaches to understanding the nature and impact of past environmental and climatic change in the archaeological record, and the role of isotope zooarchaeology. *J. Archaeol. Sci. Rep.* **23**, 968–972 (2019).
 148. O'Regan, H. J., Wilkinson, D. M., Wagner, D. & Evans, J. Why so high? Examining discrepancies between the Sr biosphere map and archaeological tooth data from the Peak District, England. *J. Archaeol. Sci.* **157**, 105826 (2023).
 149. Boethius, A., Kielman-Schmitt, M. & Robson, H. K. Mesolithic Scandinavian foraging patterns and hunting grounds targeted through laser ablation derived $^{87}\text{Sr}/^{86}\text{Sr}$ ratios at the Early-Mid Holocene site of Huseby Klev on the west coast of Sweden. *Quat. Sci. Rev.* **293**, 107697 (2022).
 150. Hülsemann, F. et al. Global spatial distributions of nitrogen and carbon stable isotope ratios of modern human hair. *Rapid Commun. Mass Spectrom. RCM* **29**, 2111–2121 (2015).
 151. McAllister, M. S., Morley, M. W., Tyler, J. J., McInerney, F. A. & Blyth, A. J. Investigating the palaeoenvironmental context of Late Pleistocene human dispersals into Southeast Asia: a review of stable isotope applications. *Archaeol. Anthropol. Sci.* **14**; <https://doi.org/10.1007/s12520-022-01540-3> (2022).
 152. Vitali, V. et al. High-frequency stable isotope signals in uneven-aged forests as proxy for physiological responses to climate in Central Europe. *Tree Physiol.* **41**, 2046–2062 (2021).
 153. Di Matteo, G., Nardi, P. & Fabbio, G. On the use of stable carbon isotopes to detect the physiological impact of forest management: the case of Mediterranean coppice woodland. *For. Ecol. Manag.* **389**, 158–166 (2017).
 154. Drucker, D. G., Bridault, A., Hobson, K. A., Szuma, E. & Bocherens, H. Can carbon-13 in large herbivores reflect the canopy effect in temperate and boreal ecosystems? Evidence from modern and ancient ungulates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **266**, 69–82 (2008).
 155. Hobson, K. A. & Kardynal, K. J. Multi-isotope ($\delta^2\text{H}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) feather profiles and morphometrics inform patterns of migratory connectivity in three species of North American swallows. *Mov. Ecol.* **11**, 48 (2023).
 156. Knipper, C. et al. Coalescing traditions-Coalescing people: community formation in Pannonia after the decline of the Roman Empire. *PLoS ONE* **15**, e0231760 (2020).
 157. Brettell, R., Montgomery, J. & Evans, J. Brewing and stewing: the effect of culturally mediated behaviour on the oxygen isotope composition of ingested fluids and the implications for human provenance studies. *J. Anal. At. Spectrom.* **27**, 778 (2012).
 158. Balasse, M. Reconstructing dietary and environmental history from enamel isotopic analysis: time resolution of intra-tooth sequential sampling. *Int. J. Osteoarchaeol.* **12**, 155–165 (2002).
 159. Balasse, M., Ambrose, S. H., Smith, A. B. & Price, T. D. The seasonal mobility model for prehistoric herders in the south-western Cape of South Africa assessed by isotopic analysis of sheep tooth enamel. *J. Archaeol. Sci.* **29**, 917–932 (2002).

Acknowledgements

M.L.C.D. thanks the Basel Doctoral Program in Ancient Civilisations (DBAW) for financing the workshop at the University of Basel during which the concept and preparation of this paper were initiated. M.K.'s research is funded by the Swiss National Science Foundation (SNSF/SNF): Project EXOCHAINS - Exploring Holocene Climate Change and Human Innovations across Eurasia (SNSF grant number: TMPFP2_217358). K.B. thanks ERC-selected/UKRI-funded project EP/Y023641/1 for salary support during production of this manuscript. U.B. received funding from the Czech Science Foundation (# 23-08049S; Hydro8), the ERC Advanced Grant (# 882727; Monostar), and the ERC Synergy Grant (# 101118880; Synergy-Plague). We further thank Judith Sealy, Noreen Tuross, Chris Baumann, Michelle Alexander, Claudia Gerling, Dominik Fleitmann, Stéphane Affolter, Ansgar Kahmen, and Daniel Nelson for their input during the related workshop. Open Access publication is supported by the Swiss National Science Foundation (SNF) and the EXOCHAINS project awarded to MK.

Author contributions

Margaux L. C. Depaermentier: concept, writing, and revision. Michael Kempf: concept, writing, revision and visualisation. Marc Vander Linden: concept and writing. Kate Britton: concept, writing and revision. Richard Madgwick: concept and writing, Ulf Büntgen: writing and revision. Dorothée G. Drucker: concept, writing and revision. Jennifer R. Jones: concept, writing and revision. Christophe Snoeck: concept and writing. Rhiannon E. Stevens: concept, writing and revision.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s43247-025-02507-7>.

Correspondence and requests for materials should be addressed to Margaux L. C. Depaermentier.

Peer review information *Communications Earth and Environment* thanks Penny Bickle and the other anonymous reviewer(s) for their contribution to the peer review of this work. Primary handling editor: Aliénor Lavergne. [A peer review file is available.]

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2025