



Evaluating rehabilitation and return to play procedures in male professional football: A narrative review

Ben Dixon^{1,2} , Jill Alexander¹  and Damian Harper¹ 

Abstract

Injuries in male professional football present a multi-faceted and complex challenge for practitioners with significant consequences across performance, psychosocial and financial domains. Medical and performance staff encounter considerable pressure to maximise player availability and advance rehabilitation timeframes, whilst minimising the risk of subsequent injury. Decisions involving increased risk promise, potentially, higher rewards and may significantly influence a player's career and team performance. Therefore, it is crucial that rehabilitation and return to play (RTP) procedures are evidence-informed and strategically designed to mitigate subsequent injury risk. Due to the complex nature of injuries, and the multidisciplinary approach required during the rehabilitation process, a wide range of knowledge and skills are essential to inform a shared decision-making process and successful RTP. To support the understanding of the competencies required, this narrative review gives an overview of existing aetiological models, decision-making frameworks and RTP approaches within the current literature. The current role of criteria-based progressions in rehabilitation frameworks are evaluated and potential future research, that could improve rehabilitation procedures and inform better RTP decision-making and post-RTP care in the professional football environment, is highlighted.

Keywords

Aetiological models, decision-making, injury, multidisciplinary, soccer, sports medicine

Introduction

The implications of injury within professional football are multi-faceted, complex and can have significant performance, psychosocial, and financial implications.^{1,2} Significant pressure is placed upon medical and performance departments to maximise player availability and accelerate rehabilitation. Usually, riskier options promise higher returns and the decision to progress or delay a player's return to play (RTP) may have a significant impact on a player's career and a team's future performances.³ Therefore, it is vital that the support given to a player during rehabilitation, and the RTP process, is well informed, timely and minimises the risk of subsequent injury. This dilemma challenges the practitioner to balance multiple processes, influential contextual factors and specific individual needs, by blending 'art' and 'science' through effectively applying current research into practice.⁴

Several studies have been conducted globally on injury epidemiology in male professional football.^{5–13} Reportedly, teams competing at the highest level in Europe typically suffer 50 injuries per season.⁵ Injury incidence studies across male professional football reports ranges between 6.5–9.1/1000hours^{5–12,14} with injury incidence rates similar between

professional leagues and countries.¹⁰ The variation in reported overall incidence may be due to studies reporting single season estimates⁹ compared to others using inter-seasonal differences.^{5,7,8,11} Inter-seasonal changes give important information on injury trends and are critical in the development of injury prevention strategies.¹¹ Furthermore, dense fixture schedules, alongside the increasing physical and psychological demands of the game,^{15–17} add complexity to the rehabilitation process, to ensure a player is fully prepared for RTP.

Reviewer: João Gustavo Claudino (State University of Campinas, Brazil)

¹Football Performance Hub, Institute of Coaching and Performance, School of Health, Social Work and Sport, University of Lancashire, Preston, UK

²Performance and Medical Department, Millwall Football Club, London, UK

Corresponding author:

Ben Dixon, Football Performance Hub, Institute of Coaching and Performance, School of Health, Social Work and Sport, University of Lancashire, Preston, UK.

Emails: bjdixon@lancashire.ac.uk; bdixon@millwall.co.uk

Match injuries have a higher incidence (20.9–58/1000 h), compared to training (2.8–6.8/1000 h).^{5,8–12} Lower extremity injury incidence was most frequent (6.8/1000 h), with muscle and tendon injuries the most frequent type (4.6/1000 h).¹⁰ It is important to consider the severity of injury and associated time-loss. Severe injuries (>28 days lost) accounted for between 15–23% of injuries sustained with moderate injuries (8–28 days lost) between 33–47%.^{5,8–11} On average, each player misses 37 days per season meaning approximately 12% of the season is lost due to severe injury,⁵ with the most severe injuries resulting in longer rehabilitation periods and potential threats to player's careers.¹⁸

Injury severity can strongly impact the physical and psychosocial characteristics of the player demonstrating the need for a holistic approach to rehabilitation.^{19,20} High severity of injury can negatively affect individual player performance, across mental, physical and technical domains^{18,21,22} with injury burden known to negatively affect team performance and results.^{1,2} Reportedly, lower injury incidence was correlated with higher league position, more games won, more goals scored and greater total points.² Consequently, this has the potential to influence job security of the head coach, with the average tenure reported to be less than 16 months,²³ or the surrounding multidisciplinary team (MDT).²⁴ The financial implications are significant. For English Premier League teams, an average cost of approximately £45 million per season per team²⁵ has been reported, in Australian football costs range between AUD\$ 187,990 to 332,680 annually,²⁶ for a Brazilian club hamstring injuries alone reportedly caused a potential loss of \$43.2 million USD in a single season²⁷ and the cost of muscle injuries, for Spanish First Division teams, equates to a total monthly cost of €7.3 million.²⁶ This provides professional football clubs with a strong economic incentive to invest and improve injury prevention and rehabilitation processes.

An index injury is defined as the first injury to occur or any subsequent injury that is clinically unrelated to the previous injury.²⁸ Reinjury is defined as “an injury of the same type and at the same site as an index injury and which occurs after a player's return to full participation”.²⁵ Incidence of reinjury is relatively low (1.3/1000 h), however, they reportedly account for 6.1–18.8% of injuries.^{5,10,29} High recurrence rates have been reported for hamstring (12–43%), groin injuries (31–50%), knee sprains (30–40%) and lateral ankle sprains (17%).^{6,7,14,30} These findings demonstrate that some reinjuries are higher risk, however, could be attributable to inadequate rehabilitation or premature RTP.¹⁴ Most identified studies recognise the longer timeframe of rehabilitating a reinjury causing a higher burden than initial injuries.^{5,8–11,31} Further considerations are the increased time loss suffered and the psychological impact, including player mental health, faith in the rehabilitation process and the trust between player and practitioner.³² Reinjury anxiety had been identified as a mediating factor in psychological readiness to return.^{33–35} Therefore, the process from initial injury

assessment to final return to performance (RTPerf) is crucial to promote a reduction in subsequent injury and to maintain player performance.

Epidemiological data emphasises the importance of effective injury rehabilitation programming, informed through a multidisciplinary approach,^{19,36,37} and the need for evidence informed return to training (RTT) and RTP protocols.^{5–11} Several rehabilitation and RTP frameworks have been developed in male professional football including the “control-chaos continuum”,^{38–40} a five stage on-field program⁴¹ and a criteria-based return to performance pathway.⁴² To continually improve and inform rehabilitation procedures, and RTP decision-making, further development and understanding of objective criteria is vital to minimise the risk of further injury. The factors described provide a clear rationale for the continued advancement of injury prevention and rehabilitation strategies³⁹ to ensure player reintegration post-injury is optimised and reinjury risk minimised.

The aim of this narrative review is to give an overview of aetiological models, decision-making frameworks and return to sport (RTS) approaches reported within the current literature. The role of criteria-based progressions in rehabilitation frameworks, and their application within male professional football, are evaluated through the identification of potential future research, that could improve rehabilitation procedures and inform better RTP decision-making and post-RTP care. Common phrases used to describe the ‘return’ of a player during different phases of rehabilitation in football are defined in Table 1 and used within the subsequent sections of this review.

Aetiological models

An understanding of injury prevention and aetiological models can assist practitioners in discussing possible causes of injury whilst appreciating the complex nature of the event. A mechanism of injury refers to the method by which damage or trauma occurs, and can be defined as the process of considering the forces involved in an injury-causing event.⁴⁵ Injuries in sport are related to some form of overload, either involving direct trauma, mechanical failure through ‘stress’ or ‘strain’ or a combination of both.⁴⁶ When the ‘stress’ (i.e., the internal forces experienced by a structure defined as force per unit of area) and/or ‘strain’ (i.e., the amount of deformation or length change in the direction of the applied force)⁴⁶ exceeds the maximal strength or failure strain capacity of a particular tissue type, this results in injury.^{46–48} Traditional injury prevention models have been reductionist in an attempt to simplify multifaceted components, to identify relationships or a sequence of events, but fail to acknowledge injuries involve complex interactions between numerous factors.⁴⁹ Aetiological models have cited the interaction between both intrinsic and extrinsic factors such as age, injury history, neural inhibition, fascicle length alteration, strength deficiencies, neuromuscular fatigue, movement efficiency, training load and

Table 1. Descriptions of different phases of 'return' of a player during different phases of the rehabilitation process.

Phase Term	Abbreviation	Description
Return to Participation	-	The player is participating in rehabilitation, training (modified or unrestricted), or in sport at a lower level than their RTS goal. The player is physically active, but not yet 'ready' (medically, physically and/or psychologically) for full RTS. ⁴³
Return to Sport	RTS	The player has returned to their sport but is not performing at their desired performance level. ⁴³
Return to Training	RTT	The player is gradually exposed to different aspects of team training. ⁴² Further RTT subphases include: <ul style="list-style-type: none"> • Partial Team Interaction: introduce the player to partial elements of team training and the club-specific training structure with restrictions agreed.⁴ • Team Interaction: involve player in modified team training, modification of drills/game-based training and training parameters in line with team training content/club-specific structure and in season micro-cycle with restrictions agreed. There is a high level of adaptability in terms of the level of integration and aims to challenge the player before resuming full team training.⁴ • Non-contact Modification: the player is re-integrated into non-contact team training.⁴⁴ • Contact Modification: the player is progressively integrated into team training and exposed to contact.⁴⁴ • Full Integration: the player is fully integrated into team training.⁴⁴ <p>Subphases and level of modification are influenced by injury type, severity, length of absence and potential load increases.⁴</p>
Return to Play	RTP	The player is gradually exposed to competitive match-play. ⁴²
Return to Performance	RTPerf	The player has returned to match-play and is performing at or above their preinjury level. ^{42,43}

competition schedule amongst others.^{50,51} Therefore a multidisciplinary, holistic approach is vital to accommodate the complex and individual nature of injury and rehabilitation.⁴⁹ An overview of the development of aetiological models is presented in Table 2.

Reviewing, understanding and applying aetiological models, can help practitioners to build injury mitigation strategies,^{50,71} analyse load/response^{75,76} and inform RTP with the aim of reducing injury risk. It is important to acknowledge the dynamic, recursive and complex nature of interacting internal (e.g., age, sex, physical fitness, psychological factors) and external (e.g., sports rules, environment) risk factors alongside the situational patterns and mechanisms that can lead to injury.⁶⁵ Furthermore, contextual factors, such as team dynamics, coaching philosophy, game model, positional switching, season phase and fixture congestion may impact injury occurrence and rehabilitation processes.⁷⁷ For example, studies have shown, during congested fixture schedules, overall injury incidence to increase from 15.6 to 33.7 per 1000 h and match injury incidence from 44.8 to 50.3 per 1000 h.⁷⁸ The application of aetiological models into injury prevention frameworks⁷¹ is, therefore, an important consideration for practitioners to understand. It is also important to acknowledge the concept of quaternary prevention⁷⁹ and integrate this into prevention programming and contemporary prevention models. Quaternary prevention aims to protect athletes from interventions that may cause more harm than good, e.g., overdiagnosis (identifying problems that were never going to cause harm), overtreatment (providing unnecessary medical

interventions), and overmedication (prescribing too much or unnecessary medication).⁷⁹ Furthermore, it is proposed that rehabilitation programming should implement the principles of quaternary prevention to protect athletes from excessive interventions and use reliable, validated tools and testing protocols to accurately evaluate and help make informed RTP decisions.⁷⁹ Future research into injury associations and causal pathways, acknowledging the complex nature of injury, can further inform the progression of these models, and their role within injury prevention frameworks.

Decision-Making and multidisciplinary approaches to return to play procedures

Given the dynamic, recursive nature of injuries, the multiple stakeholders involved, and the financial implications within the rehabilitation process, debate around who is responsible in making the final RTP decision and how that decision is made within a complex environment remains a contemporary topic. Traditionally, medical practitioners have been accountable for injury records within clubs, however, evidence suggests an association with head coach leadership styles and the quality of internal communication.^{32,80–83} Other important factors include the growing influence and size of medical and performance departments,²⁴ the dynamics of the performance structure, sharing individual expertise, psychologically safe environments, clearly defining problems and applying solutions, the need for fast, intuitive

Table 2. An overview of the development of aetiological models.

Study	Year	Model of Injury Causation	Type	Focus
Ettema ⁵²	1973	Stress-Capacity Model	Balance scale model: stress and capacity seen as static entities and must be balanced to prevent injuries.	<ul style="list-style-type: none"> Developed from preventative medicine identifying risk factors as external (environmental) and internal (capacity by the state of the personal). Stress is determined by external factors and capacity by internal factors. Highlights the potential to manipulate the external environment to reduce risk and the influence of psychosocial factors.
van Mechelen et al. ⁵³	1987	Sequence of Prevention Model	Four-step dynamic loop: <ol style="list-style-type: none"> Establishing the extent of the sports injury problem. Establishing aetiology and mechanism of injuries. Introducing preventive measures Assessing their effectiveness by repeating step 1. 	<ul style="list-style-type: none"> Evaluates the effectiveness of injury prevention interventions. One of the first models to identify injury aetiology as a key component of the process. Influential in the development of sports injury research framework, TRIPP⁵⁴ and was developed to incorporate a socioecological view and addressing the injury in context⁵⁵.
Andersen & Williams ⁵⁶	1988	A Model of Stress and Athletic Injury	Interactional multi-component model of injury presenting the cognitive, physiological, attentional, behavioural, intrapersonal, social, and stress history variables that may influence injury occurrence and prevention.	<ul style="list-style-type: none"> Built around the stress response with four central components: 1) potentially stressful athletic situation, 2) cognitive appraisal of the situation, 3) physiological and attentional responses, 4) potential injury outcome. Above the stress response are three contributing areas: 1) personality factors, 2) history of stressors, 3) coping resources. Model is predicated on the assumption that two mechanisms behind the stress-injury relationship are increases in general muscle tension and deficits in attention during stress. To reduce the impact of these mechanisms two groups of interventions are presented addressing either the cognitive appraisal or the physiological/attentional aspects.
van Dijk et al. ⁵⁷	1990	Stress-Strain-Capacity Model	Time-based three phase sequential model: <ol style="list-style-type: none"> Stress: external load and personal control. Strain: all acute and short-term effects, e.g., acute sports injuries. Long term and permanent effects, e.g., overuse injuries. <p>Capacity is a dynamic value that is the sum of all intrinsic factors allowing an athlete to perform.</p>	<ul style="list-style-type: none"> Considers the effects of training, overload and behavioural aspects over an extended time period. The model acknowledges that stress and capacity are dynamic units, and an athlete can actively affect stress and strain. An injury is the result of a complex set of interactions of internal and external risk factors.

(continued)

Table 2. Continued.

Study	Year	Model of Injury Causation	Type	Focus
van Mechelen et al. ⁵⁸ (Modified from Backx et al., 1990 ⁵⁹ , Bol et al., 1991 ⁶⁰ , Kok & Bouter, 1990 ⁶¹)	1992	Sports Behaviour Model	Dynamic model on the interrelation between risk factors, the occurrence of injury and subsequent preventative measures.	<ul style="list-style-type: none"> • Illustrates the impact behaviour may have on injury risk. • Determinants are influenced by internal/ external factors, personal equipment and load. • Three factors considered determinants of behaviour: <ol style="list-style-type: none"> 1. Attitude: the knowledge and beliefs of a person concerning the consequences of a certain form of behaviour. 2. Social Influence: influence by others. 3. Self-efficacy-cum-barriers: whether one is able to perform the desired behaviour.
Meeuwisse ⁶²	1994	Multifactorial Model	Epidemiological, multifactorial model: Proposes the model should be used to examine various factors and their interrelationships.	<ul style="list-style-type: none"> • Formed from the assessment of causal and associative factors. • Accepts previously identified intrinsic and extrinsic risk factors and adds an inciting event as the final link to injury occurring. • Suggests a different relative contribution of intrinsic and extrinsic factors for acute and overuse injuries.
Williams & Andersen ⁶³	1998	Revised Model of Stress and Athletic Injury	Development on the interactional model of stress-injury.	<ul style="list-style-type: none"> • Highlighted the bidirectional nature directly between personality traits and coping resources and interaction between the three areas of stress response. • Removed medication as a proposed intervention. • Expanded the peripheral narrowing and distractibility variables to include audition.
McIntosh ⁶⁴	2005	Multifactorial and Biomechanical Model	Multifactorial model with a biomechanical focus on tissue properties and injury.	<ul style="list-style-type: none"> • Incorporates multiple intrinsic and extrinsic inputs, and accounts for acute and repeated events. • Introduces the effect of increased or decreased mechanical load alongside the individual load response. • Acknowledges the importance of movement co-ordination and efficiency, and that injury interventions focus on internal and external load modification by reducing load or improving capacity.
Bahr & Krosshaug ⁶⁵	2005	Comprehensive Model	Comprehensive model combining epidemiological and biomechanical approaches with a focus on the specific sport.	<ul style="list-style-type: none"> • The importance of injury mechanism and the language used to describe injury inciting events is highlighted. • The key component added is a precise, multi-level description of the inciting event. • The different levels are categorised: <ol style="list-style-type: none"> 1. Playing situation. 2. Player/opponent behaviour. 3. Gross biomechanical description (i.e., whole body). 4. Detailed biomechanical description (i.e., specific joint).
Meeuwisse et al. ⁶⁶	2007	Dynamic, Recursive Model	Cyclical model based upon a contact injury model ⁶⁷ and incorporating the consequences of repeated sport participation.	<ul style="list-style-type: none"> • Evolved the concept that injuries are dynamic and recursive and changes in injury risk with training exposure result in constant variations in injury susceptibility. • Ever-changing risk factors are important to acknowledge in injury prevention.

(continued)

Table 2. Continued.

Study	Year	Model of Injury Causation	Type	Focus
Appaneal & Perna ⁶⁸	2014	Biopsychosocial Model of Stress and Athletic Injury and Health (BMSAIH)	Interactional model proposed as an independent extension to the revised model of stress and athletic injury.	<ul style="list-style-type: none"> • Focuses on the dynamic interactions between physiological, psychological and social elements and the collective influence on stress, injury and health. • Expands on previous models in three ways: 1) it clarifies mediating physiological pathways between athletes' stress response and adverse health outcomes, 2) it considers other health outcomes and behavioural factors that impact sport participation, 3) it integrates the impact of training upon athlete health. • The central component is psychosocial distress (e.g., negative life events) may act synergistically with training-related stress increasing susceptibility.
Bittencourt et al. ⁶⁹	2016	Complex Systems Model	Applies a complex systems approach, to evaluate risk factors, identify interactions and how these interactions contribute to injury, within a dynamic, recursive aetiological model.	<ul style="list-style-type: none"> • Proposes sports injuries will be better understood by recognising frequent patterns of interaction among multilevel risk factors. • These risk factors are unpredictable interacting units within a complex system resulting in a web of determinants. • These interactions form regularities which emerge as either injury or adaptation. This constrains the interactions creating a recursive loop and dynamically shaping further interactions. • Risk factors have a different weighting for different physical demands and an individual's tolerance, therefore, looking for existing patterns and identifying relationships is necessary to allow the development of effective interventions to specific risk profiles. • Suggests research should move from using isolated risk factors to injury pattern recognition by identifying the complex pattern of interactions.
Windt & Gabbett ⁷⁰	2017	The Workload-Injury Model	Recursive model incorporating workload within the causal chain for injury.	<ul style="list-style-type: none"> • This model was the first to explicitly incorporate workloads within the causal chain for injury. • The dynamic nature of injuries is acknowledged, and the concept that application of workload is the primary process whereby an athlete is exposed to external risk factors and potential inciting events is presented. • Proposes a conceptual framework for 'why' workloads are strongly associated with injuries by continually modifying injury predisposition through either positive or negative physiological adaptations.
O'Brien et al. ⁷¹	2019	The Team-sport Injury Prevention Cycle (TIP)	Three-step continual cycle model: 1. (Re)Evaluate: evaluating current injuries and analysing current prevention strategies being used.	<ul style="list-style-type: none"> • Considers the application of aetiological models into injury prevention strategies. • TIP reflects the cyclical nature of injury prevention and considers a team's unique/specific context and challenges as part of the process. • A multidisciplinary approach is highlighted, and practitioners can assess injury risk whilst

(continued)

Table 2. Continued.

Study	Year	Model of Injury Causation	Type	Focus
			<ol style="list-style-type: none"> 2. Identify: exploring the risk factors and mechanisms underpinning the injuries identified during the evaluation phase. 3. Intervene: planning both the content and delivery of injury prevention strategies. <p>Ongoing re-evaluation and modification is required as part of a dynamic, cyclical process.</p>	identifying barriers and facilitators that can impact the success of prevention strategies.
Kalkhoven et al. ⁴⁶	2020	Model for Athletic Injury and Framework for Stress-related, Strain-related and Overuse Injury	<p>Conceptual model formed with six interacting levels:</p> <ol style="list-style-type: none"> 1. Causal contextual factors. 2. Individual physiological profile, functioning, mechanical properties and force applied to the body. 3. Load tolerance of structures and loading applied. 4. Specific tissues internal stress and strain experienced. 5. Timeframe of stress/strain application (immediate/repetitive). 6. Injury occurrence. 	<ul style="list-style-type: none"> • Considers the interaction between physiological and mechanical factors and highlights possible causal pathways. • Attempts to establish direct injury causation by investigating how risk factors contribute to injury occurrence. • Considers how individual components interact with other factors that underpin the load tolerance of different structures/tissues, and the loads applied. • Introduces the consideration of different loading patterns resulting in various injury outcomes and defining these as acute stress-related, strain-related, or overuse. • A holistic approach using concepts of load tolerance and application, physiological and mechanical properties of specific tissues and attempting to identify causal pathways had not been previously established.
Liveris ⁷²	2025	System Dynamics Model	<p>Computational modelling method to simulate dynamic, complex systems.</p>	<ul style="list-style-type: none"> • Initially uses Causal Loop Modelling (CLD) to evaluate intrinsic, extrinsic and institutional risk factors interrelationships that lead to injury. • CLD shows a perspective of risk factors' dynamic, nonlinear association, the connections between elements having a positive or negative effect, and the links create loops characterised as reinforcing or balancing.⁷³ • Further employs Partial Least Squares Structural Equation Modelling to quantitatively assess intrinsic risk factors' interrelationships and their effects on hamstring injuries and frequency of non-contact lower limb injuries.⁷⁴

Legend: BMSAIH, Biopsychosocial Model of Stress and Athletic Injury and Health; CLD, Causal Loop Modelling; TIP, Team-sport Injury Prevention Cycle; TRIPP, Translating Research into Injury Prevention Practice framework.

decision-making and the quality of internal communication within the MDT and across departments.^{83–85} Therefore, it is important that practitioners develop an understanding of decision-making frameworks, the importance of multidisciplinary approaches and have clear communication procedures to create an informed, shared decision-making process and develop purposeful application of critical thinking skills.⁸³ Table 3 presents an overview of decision-making frameworks that are applicable in a sporting environment to guide the RTP decision-making process.

Arden et al.⁸⁶ state that, ideally, decisions are guided by accepted clinical criteria, however, many have little empirical evidence and there is a lack of agreement on the best criteria. It was proposed, a logical step would be a consensus statement on RTS and to ascertain agreement in key areas.^{43,86} The consensus statement on RTS presented by Arden et al.⁴³ is divided into 4 main sections: 1) definitions related to RTS, 2) models to guide RTS, 3) evidence to inform decision-making and 4) future research. A key concept highlighted is advocating an athlete-centred approach, placing them in the position of active decision-maker, alongside other relevant stakeholders, as part of a shared decision-making process.^{43,87} The authors cited three models to guide the RTS process: 1) the StARRT framework,⁸⁸ 2) the biopsychosocial model²⁰ and 3) optimal loading.⁸⁹ The authors state that load progression is key and maintaining 'optimal' loading, by gradually increasing workload and avoiding 'spikes',⁹⁰ are important clinical considerations.⁴³ The importance of biological, psychological and social factors influencing decision-making are presented and the role of 'load' in rehabilitation and post-RTS were defined.⁴³ The lack of consensus and scientific evidence for RTS criteria for the majority of sport injuries is highlighted and a future priority was identified to address the identification of positive and negative predictive (or prognostic) factors for RTS outcomes.⁴³

Shared decision-making

Dijkstra et al.⁹² acknowledged multiple individuals are involved in the decision-making process, and including the athlete in the process fulfils psychosocial needs and supports an athlete-centred approach.⁹⁵ The fluid nature of decision-making is acknowledged, however, only three stakeholders (healthcare professional, athlete and coach) are highlighted. Within the context of professional football, medical and performance departments, the player, technical coaches and directors are also involved.⁹⁶ Furthermore, with significant quantities of subjective and objective data available, and influential contextual factors (e.g., player status, match importance), informing RTT and RTP decision-making a flexible structure may be required. Developing a flexible structure, specific to each unique club environment, allows practitioners to incorporate different individual experiences and skillsets, be guided by subjective and

objective data and consider the context of the decision to allow an informed, shared approach.

Within professional football RTP practices have been investigated and the translation of research into practice has been highlighted to enhance the process.⁹⁷ It was reported that a shared decision-making process was used by 80% of teams although there was a lack of agreement in the level of involvement across phases by different practitioners.⁹⁷ This could be accounted for by potential bias, however, the inconsistency found in the composition raises some potential concerns about the specific dynamics of the communication among staff.⁹⁷ Shared decision-making was demonstrated across the phases of rehabilitation with different weighting of importance given as rehabilitation progressed, however, challenges were acknowledged relating to team hierarchy, particularly as players returned to matches.⁹⁷ This demonstrates a positive use of shared decision-making in football, however, further research into MDT dynamics has demonstrated uncertainty around who makes decisions and how they are made can be an issue.⁸⁴ Additionally, department growth, resulting in more lines of communication, and modernising methods of communication (i.e., mobile phones, email, social media rather than face to face conversations)²⁴ further complicate the shared decision-making process. With leadership styles and communication potentially impacting injury,^{80–82} clearer processes, communication pathways and RTP decision strategies are evidently important and required.⁸³

Criterion-Based decision-making

Whilst respecting biological healing timeframes, a paradigm shift towards criterion-based approach to RTS is evident and studies have shown a reduction in reinjury rates when individuals have passed objective criteria before RTS.⁹⁸ It is evident that practitioners are utilising evidence-based research and testing procedures throughout the rehabilitation process, informing decisions and adapting decision-making frameworks, although further research is required into RTP criteria in practice.^{43,99} There is consensus for criteria-based RTP factors that a range of clinical, physical, functional and psychological measures should be used.^{43,100} These influence and inform decision-making, and several studies have investigated their use and role within injury, rehabilitation and decision-making frameworks.^{43,97–104} In a survey by Dunlop et al.,⁹⁷ the use of criteria-based evaluations and clinical criteria were reported during early phases, with sport-specific functional criteria (e.g., acceleration/deceleration, maximal sprints) assessed across all phases and a greater focus placed on psychological readiness in the latter stages. Despite the translation into practice no information was given to advance knowledge on specific metrics or thresholds to be used within the professional football environment.

Table 3. An overview of decision-making frameworks that can be used to guide return to play decision-making processes.

Study	Year	Decision-making Framework	Overview	Process
Creighton et al. ⁹¹	2010	Decision-Based Return to Play Model (RTP)	<ul style="list-style-type: none"> Attempts to clarify the processes and provide a structure with a logical rationale when making an RTP decision. Uses an influence diagram incorporating the states of nature elements, decision elements and the contributing information between elements Was the first model to attempt to provide a foundation for the integration of individual factors and components for evidence-based rationale for RTP decision-making. 	<ol style="list-style-type: none"> Evaluation of health status: the stage of healing of the injury. Evaluation of participation risk: the specific demands of the sport and injury. Decision modification: further potential influential factors.
Shrier ⁸⁸	2015	Strategic Assessment of Risk and Risk Tolerance (StARRT)	<ul style="list-style-type: none"> Considers RTP decision-making as a risk assessment. Assessed risk is compared to the risk tolerance of the individual. If assessed risk is greater than the risk tolerance, RTP should not be allowed. The key difference is StARRT identifies tissue health and stresses as decision elements, the role of this in injury and the factors that influence each step. 	<ol style="list-style-type: none"> Assessment of health risk: the stress the tissue can tolerate, i.e., the health of the tissue. Assessment of activity risk: the stress that will be applied during the sporting activity. Assessment of risk tolerance: the threshold for acceptable risk and factors that affect this.
Ardern et al. ⁸⁶	2016	Rearranged StARRT Model	<ul style="list-style-type: none"> The StARRT framework is a biopsychosocial model that highlights key elements in RTP decisions, however, the decision itself is not solved within the model. The StARRT model sits at the intersection of the evidence-based practice pillars. The practitioner integrates evidence with experience and patient/player preference when making decisions. The definition of a successful RTP is considered and, therefore, the decision should be shared between the practitioner and athlete. Key roles of the healthcare professional, athlete and coach are highlighted with different contributions depending upon the factors influencing the decision. The size and impact of each individual contribution is variable depending on the status or phase of injury. The importance of effective, consistent and ongoing communication to achieve trust and to make quality RTP decisions is highlighted. 	<p>Three evidence-based pillars:</p> <ol style="list-style-type: none"> Evidence: functional tests, physical exam, injury/medical history. Clinician: mask injury, liability, seriousness, ethics/conflict of interest, (re)injury prevention. Patient: competitive level, symptoms, sport, values, position, psychological state, demographics, limb dominance, values.
Dijkstra et al. ⁹²	2017	Shared Decision-making Process	<ul style="list-style-type: none"> Key roles of the healthcare professional, athlete and coach are highlighted with different contributions depending upon the factors influencing the decision. The size and impact of each individual contribution is variable depending on the status or phase of injury. The importance of effective, consistent and ongoing communication to achieve trust and to make quality RTP decisions is highlighted. 	<ol style="list-style-type: none"> Choice: making the athlete and coach aware that reasonable options exist. Option: providing more detailed information about the different options. Decision: guiding the athlete and coach to consider their preference and decide what is best.
Yung et al. ^{93,94}	2022	Complex Systems Perspective	<ul style="list-style-type: none"> Decision support systems can be developed using AI algorithms and machine learning as potential tools that can consider dynamic interaction at multiple levels simultaneously. This may allow practitioners to compare potential outcomes of different decisions (e.g. reinjury likelihood), increase decision efficiency, and identify data patterns that may cause a certain outcome to compliment human decision-making. Evaluating the quality of decision-making is an important aspect and differentiating between the decision and the outcome is important, 	<ol style="list-style-type: none"> Choose appropriate RTP test and synthesise the data in a meaningful way. Understand decision-making theories. Use shared decision-making to eliminate contextual 'blind spots' and improve decision quality.

(continued)

Table 3. Continued.

Study	Year	Decision-making Framework	Overview	Process
			<p>a high-quality decision does not necessarily warrant a good outcome due to uncertainties.</p> <ul style="list-style-type: none"> Based upon an adapted decision analysis model the authors outline a three-step framework to make systematic and objective RTP decisions. 	

Legend: AI, Artificial intelligence; RTP, return to play; StARRT, Strategic Assessment of Risk and Risk Tolerance.

The development of injury specific criteria is important.^{43,100,104,105} With particular reference to hamstring injuries in professional football five core domains were identified as part of criteria-based rehabilitation progression presented by Zambaldi et al.:¹⁰⁴ 1) functional performance, 2) strength, 3) flexibility, 4) pain and 5) player's confidence. Following this a RTS continuum was proposed for professional athletes in which measures of progression should be tested to progress through six categories: 1) movement and core, 2) strength and endurance, 3) power, 4) general and sport conditioning, 5) load performance testing and 6) self-reported outcome.⁹⁸ Criteria-based progressions assist practitioners in being transparent in the decision-making process, however, sporting environments are complex adaptive systems extending into injury and rehabilitation protocols whereby contextual factors play an important role in the development of effective RTS frameworks.

The complex and multifactorial nature of RTS decisions provides a significant challenge to practitioners with potential impacts on the athlete's well-being and performance, and the overarching performance of the team. With multiple stakeholders involved it is important to have a clear shared decision-making process with players and coaches active in the process. This may allow for important balance between intuitive and analytical processes, therefore, practitioners are encouraged to adopt models that suit the context and environment to enhance the quality of decision-making.¹⁰⁶ The development of artificial intelligence (AI), data driven algorithms and machine learning techniques, to inform clinical decision-making, is a prominent, and evolving, area of interest in sports medicine. Machine learning models, such as decision trees, Markov processes and neural networks, analyse multidimensional data to identify subtle patterns and potential signs of injury risk and tracking algorithms provide information on body movements in real-time allowing the identification of biomechanical inefficiencies¹⁰⁷ which may prove crucial as part of the rehabilitation process. Due to methodological limitations and high risk of bias, current models cannot be recommended to be used in practice,¹⁰⁸⁻¹¹⁰ however, once these limitations have been addressed, machine learning could provide enhanced systematic analysis that could provide future solutions to the training load/injury paradox.¹¹⁰⁻¹¹² Further advancement of the evidence base supporting injury prediction models, RTS and rehabilitation protocols, may improve the ability of practitioners to make high quality, evidence informed decisions to manage training load and reduce injury-risk.

Return to sport models and on-field rehabilitation

It is widely accepted the use of competency-based criteria is vital in progressing through different phases of rehabilitation and several factors, such as muscle strength,

psychological readiness, sport specific load and cardiovascular fitness, contribute to an athlete's readiness for RTS.^{19,30,38,42,43,98–100,102,104,105,113–115} The traditional notion that RTS is a single decision at a point in time has developed into the concept of an evolving continuum supporting the athlete from the onset of injury to full RTP, whilst mitigating the risk of further injury.^{42,98,113} The identified multidisciplinary input throughout the process can result in differing timeframes and objective criteria being recommended depending on the practitioner's specialist area^{87,99} increasing the relevance of having a reliable shared decision-making process in place.

Return to sport clearance Continuum (Draovitch et al., 2022)

To foster efficient progression, Draovitch et al.,⁹⁸ proposed a 'Return to Sport Clearance Continuum' (RTSCC). This continuum consists of 5 sequential phases comprising: 1) repair phase (to minimise swelling, increase range of motion and develop muscle activation), 2) rehabilitation and recovery phase (restoring normal arthrokinematics), 3) reconditioning phase (focusing on skill and force development and load-volume tolerance), 4) performance phase (transition to full training), and 5) preseason/training camp phase (to be properly managed for an upcoming season).⁹⁸ The RTSCC states training load should be monitored throughout, to avoid overloading tissues, and testing criteria is suggested to ensure objective progress through the phases.⁹⁸ The scope of practice for staff involved at each phase is not considered as part of the continuum, despite the different skills required. Specialised personnel and team work between the medical and performance departments is important,¹⁹ yet the specific involvement is often unclear.⁹⁷ The RTSCC provided a general framework for RTS processes and decision-making to be developed, although more detail of practitioner responsibilities may be useful to develop an understanding of the specialist skills and knowledge required throughout different phases of rehabilitation.

The return to performance pathway (Mitchell and Gimpel, 2024)

A football specific framework has been initially proposed by Mitchell et al.¹¹³ and developed by Mitchell and Gimpel.⁴² The 'Return to Performance Pathway' includes eleven phases from a diagnosis and planning phase, into an acute phase and then progression through multiple gym and grass phases. A key component of the pathway is the objective exit criteria for each phase in an attempt to ensure safe progression.⁴² Mitchell and Gimpel's⁴² framework demonstrates the importance of a progressive multistage rehabilitation process and has adapted previous frameworks for the end-stage rehabilitation phases known as on-field rehabilitation (OFR).^{38,41,42,116} The RTPerf Pathway

framework⁴² provides a comprehensive overview of the rehabilitation process by introducing acute and gym phases prior to starting OFR, whereas previous frameworks had focused primarily on OFR.^{38,41,116–119} These final phases are highlighted as being important to prepare the athlete for re-entry into sport and where the overlap between rehabilitation and RTS processes occurs. This overlap requires specialised personnel, and teamwork between the medical, performance and coaching departments, to transition the athlete and bridge the gap between rehabilitation and sports specific training.^{19,36} OFR represents these final phases as the athlete needs to be physically prepared for the demands of their sport, from both an injury specific and a general conditioning perspective. A stronger focus is required in these phases to prepare the athlete as there is a lack of validated competency criteria for final RTP protocols, particularly for moderate-severe (>14 days) injuries.^{31,36,49,116,120} Risk of subsequent injury and inter-injury relationships need to be evaluated and specific player monitoring post-RTP is not considered. In the period following RTP a 'one month risk decay' of subsequent non-contact injuries has been reported.¹²¹ Following return, initial risk of non-contact subsequent injury was about two times higher than baseline. This risk diminished by half after approximately 25 days and levels off afterwards.¹²¹ The severity of injury should be considered with severe injuries showing an increasing injury risk within the first ten days and remaining relatively high thereafter.¹²¹ The continuous hazard curve of non-contact injuries shows a decline towards four weeks post RTP¹²¹ indicating additional phases may be required to assess exposure to pre- and post-RTT loads and to further inform tertiary injury mitigation strategies.¹²²

On-Field rehabilitation

Table 4 presents conceptual models specific to OFR that have been developed^{38,41,116–119} and re-evaluated to ensure rehabilitation processes are representative of most recent evidence and practice design.^{39,40} The frameworks and studies presented demonstrate the strength of progressive OFR protocols and the impact on reinjury risk and subsequent physical performance. Cited limitations include the prescriptive nature of programmes and the individual nature of injury and responses to load interventions.⁴⁹ These conceptual frameworks can offer increased flexibility to support decision-making and allow for individual criteria-based progression throughout different stages of rehabilitation. This empowers practitioners to continually evolve their practice and understanding of the process.⁴⁹ These frameworks provide a solid conceptual guide for practitioners to utilise during rehabilitation, however, they are based upon expert opinion, inductive reasoning and case study applications.^{123–125} There is a need for validation through experimental evidence and testing alongside practical insights into how OFR is currently executed.^{49,126}

Developing on-field rehabilitation

The concept of 'load' has been highlighted through the aetiological, decision-making, RTS and OFR frameworks previously described.^{38,41,43,46,57,64,70,105,115,117-119} A survey of practitioners has shown 'training load' was one of the most frequently used criteria when progressing through the final stages of rehabilitation⁹⁷ and specific metrics are available, although, more evidence is needed for the practical application of certain thresholds.^{36,97,126,127} This has been further highlighted in a survey by Armitage et al.³⁶ regarding OFR practices within professional football. During OFR, wearable technology, such as global positioning systems (GPS) and heart rate monitors,¹²⁸ was ranked as the most popular monitoring technique and is used daily in 97% of cases. Furthermore, wearable technology was ranked the second most important factor for decision-making between sessions and, as rehabilitation progresses, monitoring techniques shifted from subjective to objective measures with GPS deemed most important.³⁶ This demonstrates the influence of GPS within rehabilitation, particularly during the late OFR phases before RTP, however, there is a lack of consensus with regards to specific metrics and thresholds used. Some practitioners using targets that "are arbitrary (set at a squad level), relative (set at an individual level) or a combination of both".³⁶ In some cases, there was a change from using absolute to relative units and different thresholds were reported for acceleration and deceleration metrics.³⁶ Despite this, it is widely accepted that systematic load progression/management and the development of tissue tolerance of an injury site is vital alongside restoring sport specific qualities.^{38,41,42,49,116}

Previous methods of measuring load progressions within rehabilitation have been demonstrated to be ineffective.¹²⁹⁻¹³¹ As part of a progressive rehabilitation, acute loads are 'high' due to injured players having no chronic load, therefore, models such as acute:chronic ratio¹³² or percentage change are insufficient due to this divergence.^{129,131,133} Based on preinjury data, current running loads and previous injury 'benchmarks', chronic load can be developed, taking player tolerance into consideration, with appropriate planning and incremental load progressions.⁷⁷ Lolli et al.¹²⁹ states acute load alone could be a useful predictor of injury in absolute terms and may not require normalisation for chronic load through different statistical approaches, i.e., analysing the actual amount of measurable acute load (e.g., total distance, sprint distance or total accelerations/decelerations) without normalising or comparing it relative to any other factors (e.g., chronic load) may be suitable to assess injury risk. Improving our understanding of 'load' within each phase of rehabilitation, and potential progression targets, would aid in the development of RTS frameworks.⁴⁹ In an applied setting, practitioners, through progressive overload, should aim to increase chronic load

depending on the severity of the injury and with consideration towards the tissue type and healing process.

Research has shown a focus placed on psychological readiness during the latter stages of rehabilitation,⁹⁷ yet, with the reported shift of monitoring from subjective to objective measures through the rehabilitation phases,³⁶ practitioners must be cautious not to neglect subjective feedback as the player transitions through RTT, RTP and post-rehabilitation phases. Studies have acknowledged the importance of blending subjective and objective measures to monitor training and rehabilitation loads,^{36,134} however, physical and psychological readiness may not coincide³⁵ and, therefore, may be overlooked, misinterpreted or misunderstood. Despite the acknowledged importance of psychological readiness there is limited research into the efficacy and effectiveness of psychological strategies specifically facilitating readiness to return.³⁵ Furthermore, with the potential for injury severity to significantly impact mental health,¹³⁵ with anxiety and reinjury concerns being reported as key factors when players are close to RTP,³²⁻³⁵ subjective measures may prove invaluable. Competence (i.e., sense of proficiency in their sporting capabilities), autonomy (i.e., a sense of control over their desired RTS trajectory) and relatedness (i.e., feelings of social connection and affiliation) have been shown to be key components influencing the final stages of rehabilitation,³⁵ therefore, monitoring and strategies to develop these skills are recommended. As sport-specific training is progressed visual-cognitive demands (e.g., high speed decision-making, dual-task processing) increase³⁹ leading to a subsequent increase in mental fatigue. Mental fatigue has been shown to impact physical and technical performance,^{136,137} therefore, progressive exposure, resilience and recovery from mental fatigue may be important to consider.

Several studies cite the importance of progressive sport-specific training in the late stages of rehabilitation and as a key component of OFR frameworks.^{4,38,39,41,42,44,49,97,98,113,116,123,138} Furthermore, the understanding of the physical and cognitive demands of the sport is key in individualising injury risk management strategies throughout different stages of the season.⁵⁰ Within professional football the demands of match-play are well documented, and studies show a variation in demands depending on a number of factors, such as positional differences, formational changes and player status.¹³⁹⁻¹⁴⁶ With studies showing high numbers of matches and increasing intensity the physical and mental load on players is significant.^{15,16,147} Furthermore, high-intensity actions (i.e., accelerations, decelerations, sprinting) have been shown to be important in match defining events (i.e., scoring goals) and winning games.¹⁴⁸⁻¹⁵¹ With constantly evolving demands, requiring players to perform more frequent high intensity actions,¹⁵⁻¹⁷ consistent re-evaluation is required to ensure the best support to players during the rehabilitation process. Less fit players have been shown to

Table 4. An overview of conceptual models for on-field rehabilitation in football.

Study	Year	On-Field Rehabilitation Framework	Overview
Taberner et al. ³⁸⁻⁴⁰	2019	Control-Chaos Continuum	<ul style="list-style-type: none"> Introduces five stages of OFR, moving through 1) high control, 2) moderate control, 3) control to chaos, 4) moderate chaos and 5) high chaos. The stages interlink GPS variables, progressing volume and intensity, sport-specific conditioning, technical and movement qualities, while progressively increasing perceptual and reactive neurocognitive demands. The application of this framework has been illustrated across different case studies depending on the specific needs of the injury.^{44,124,125} Updated model of the control-chaos continuum, utilising research in injury neurophysiology, that integrates practice design and physical-cognitive interactions. The updated framework incorporates elements of visual cognition, attentional challenges, decision-making, and progressive representation of the game model to enhance sport-specific preparation for returning to sport. Increased emphasis on progressively challenging neurophysiological capabilities, skill redevelopment and reducing cognitive demands on RTS by embedding tactical principles and positional responsibilities of the game model earlier in the rehabilitation process. Emphasizes the design and delivery of progressive training in increasingly 'chaotic' conditions with the goal to align session and drill objectives to what the player experiences during competition. Adaptable for injury severity, and integrates physical-cognitive load monitoring, and strength and power diagnostics to enhance decision-making throughout RTS. Provides examples of how to adapt the 'high chaos' element to facilitate the transition to RTT and return to competition.
Buckthorpe et al. ^{41,116}	2019	On-field Rehabilitation Part 1: 4 Pillars of High Quality On-field Rehabilitation	<ul style="list-style-type: none"> Divided OFR planning into a two-part series within the context of ACL rehabilitation, however, the frameworks are designed to be applied to severe (>28 days) injuries. The 'Four Pillars' of high-quality OFR planning are 1) movement quality, 2) physical conditioning, 3) sport-specific skills and 4) training load. The sequential process begins with restoring movement patterns before gradually increasing metabolic and mechanical load and then integrating specific neurocognitive and perceptual demands. The four pillars contribute across the five stage OFR framework (Part 2). The stages proposed are 1) linear movement, 2) multidirectional movement, 3) football-specific 4) technical skills and 5) practice simulation. Each stage has specific entry criteria to ensure safe progression through the process while progressively increasing the athlete's exposure to volume and intensity. Highlights criteria for full return should include clinical, functional, biomechanical, psychological and sport-specific factors. Developed and validated a functional OFR programme for hamstring strain injury. Presents a thirteen item OFR programme arranged in order of increasing complexity. Upon completion player is declared fit to train with the team. A panel of 15 experts deemed all 13 items of the programme to be valid. 19 players who had suffered a grade two hamstring strain injury returned to play
Jiménez-Rubio et al. ^{117,118}	2019	Rehabilitation and Readaptation Program for Hamstring Injury	<ul style="list-style-type: none"> 19 players who had suffered a grade two hamstring strain injury returned to play

(continued)

Table 4. Continued.

Study	Year	On-Field Rehabilitation Framework	Overview
	2021	Indoor and On-field Reconditioning Program for Adductor Longus Injury	<p>competitively in 22.42 (2.32) days after undergoing the program, and, reportedly, did not suffer a reinjury in a 6 month follow up period.</p> <ul style="list-style-type: none"> Developed and validated an indoor and OFR programme for adductor longus injury. Presents a 16-item indoor programme and 4-item OFR programme arranged in increasing complexity. All 20 items were determined to be valid by a panel of 16 experts. The programme was implemented on 12 players and no reinjuries were reported in the 15-week follow up period.
Stathas et al. ¹¹⁹	2024	On Fi.RE Framework	<ul style="list-style-type: none"> Evaluated an accelerated OFR framework, defined as early initiation of OFR, compared to a traditional framework in male professional football. In the On Fi.RE, protocol the criteria for progression, in the early phases of rehabilitation, were higher: pain thresholds and ROM targets to initiate on-field training earlier in the process. The same criteria were used in the later stages of rehabilitation for both protocols. The On Fi.RE protocol reported a reduced time of 6.5 days to RTT and a reduced risk of subsequent injuries.

Legend: ACL, anterior cruciate ligament; GPS, global positioning system; OFR, on-field rehabilitation; On Fi.Re, On-Field Rehabilitation; RTP, return to play; RTS, return to sport; RTT, return to training.

perform fewer high-intensity actions and a quicker fatigue progression impacting technical performance^{143,144,152} and potential injury-risk.¹⁵³ This has motivational connotations for a rehabilitating player and should be a consideration for practitioners when returning players, ensuring an adequate fitness level.

There is a significantly increased injury risk in the first match after RTT and ensuring sufficient loads have been achieved and tolerated is an important factor.^{42,89,154,155} Research in Australian rules football has shown subsequent injuries to a different site or tissue were more common than reinjury^{155,156} and it has been reported that, in professional football, subsequent and recurrent injuries could account for 14% of injuries.¹⁵⁷ The reporting of subsequent injuries in sport is inconsistent and there is a paucity of data regarding severity, type and associated risk factors.^{158,159} This is an issue for injury prevention strategies, which rely on accurate injury surveillance data, as it does not provide an adequate understanding of injury risk.^{157,159} It is proposed the mechanism could be a general state of under-loading that predisposes the athlete to subsequent injury.^{155,156} This raises the question are athletes receiving enough 'general load' prior to RTT or are practitioners placing too much focus solely on 'injury-specific' rehabilitation?

Stares et al.¹⁵⁵ demonstrated higher running loads, achieved during rehabilitation, delayed RTP timeframes, however, moderate to high sprint loads could be protective against subsequent injury. The authors proposed that the accumulation of sprint loads increased chronic training load facilitating adaptation and minimising workload spikes, although chronic load was not identified as a significant factor in subsequent injury.¹⁵⁵ This demonstrates the complexity of RTP decision-making as accelerating rehabilitation protocols may not allow sufficient time for progressive overload, however, extending RTP timeframes will result in additional missed games potentially affecting team success.^{1,2,155} With medical and performance departments under pressure to ensure efficient RTP, whilst trying to minimise further injury risk, further investigation into accelerated protocols, workload progressions and thresholds is required. The high physical demands of professional football, and the ever-increasing intensity and density of these demands, are important factors in rehabilitation programming. There is a pressing need for further investigation of the final phases of rehabilitation to ensure sufficient conditioning, and to develop specific objective thresholds, to inform progression and the final RTP decision.

Future research suggestions

Research acknowledges the importance and development of progressive rehabilitation pathways. Current frameworks,^{38,39,41,42,116} utilised in professional football, are presented in this review and provide a stepwise progression for practitioners with evidence suggesting their influence in

contemporary practice,³⁶ however, a multicomponent model adopting a non-linear, concurrent approach has been proposed for a general population with potential applications for sport models.¹⁶⁰ There has been a paradigm shift towards a criteria-based approach supporting progression and it has been stated an absence of valid objective criteria risks inadequate rehabilitation.^{42,43,98,100,104} Specific practitioner competencies and knowledge, to provide optimal support, throughout the phases is still unclear and requires further evaluation.^{19,97} There is a lack of empirical evidence and knowledge of causality between 'load' application and successful outcomes in RTP and,⁴⁹ given the protective effects of moderating physical capacities (i.e., developing force-velocity profiles, energy system capacity, movement quality), further investigation is required throughout the rehabilitation process. There are significant challenges for research due to the heterogeneity of injuries, multiple, interacting moderators of risk and limited accessible sample sizes reducing statistical power,¹⁶¹ particularly with athlete populations. Despite this, conceptual frameworks are essential to aid practice¹²⁶ and development, understanding and evaluation of objective criteria informing progression is an important next step. Further informing and validating frameworks^{49,126} may improve the decision-making process for athletes, practitioners and coaches.

External load metrics for return to play

Within current OFR frameworks external load metrics are acknowledged as an important objective criterion,^{38,41,42,100,125} with metrics of interest emerging, such as very high speed running, peak speed, acceleration and deceleration distance,^{162,163} for specific injury pathologies. However, caution is advised as the metrics suggested lack empirical evidence, are based on anecdotal experiences or there is an absence of post-OFR information or pre-injury data.^{36,49,138,163} When examining the relationship between 'training load' and injury risk it is important to acknowledge this is likely to be associative and not causal,^{49,126,164} a clear aetiology has not been established¹⁶⁵ and many studies have used inadequate analysis.¹²⁶ Insights into applied practice and expertise are crucial in evolving evidence-based practices.^{36,127,166} A survey of practitioners suggests that research exploring "reduced risk of reinjury based on achieving specific markers/thresholds" would be beneficial when considered within the complex nature of injury³⁶ and an association with workload and injury risk has been demonstrated in non-injured players.^{75,167} Although 'training load' is associative, these metrics may assist in setting targets for successful RTP and support practitioners and athletes by providing objective information to make informed decisions.¹⁶⁸

GPS monitoring is widely used to quantify external load^{36,138} and theoretical justification for load progression and management within rehabilitation is strong, although

the optimal strategy is unknown.¹³⁸ Despite issues within football stadia potentially affecting signal quality,¹⁶⁹ GPS monitoring has been shown to be a reliable and validated measurement tool for evaluating movement demands within sports.^{170–172} That said, due to the rapid nature of high-intensity actions (i.e., accelerations and decelerations), the validity of these has been questioned, with sampling rate, minimal effort duration^{173,174} and signal-filtering techniques important factors to consider.^{172,175–177} Given the high musculoskeletal forces that can accompany accelerations and decelerations^{178,179} and the critical importance of these actions for player performance and injury resilience,^{142,180} further research is necessary to investigate effective rehabilitation assessment, monitoring and programming (i.e., progression of exercises) to ensure players are optimally prepared to meet the demands of these specific actions during RTT and RTP. Furthermore, the role injury site and severity may have on acceleration and deceleration mechanics and the effect this may have on altering tissue forces and properties during, and following, the rehabilitation process requires investigation.

There is consensus in using match-specific GPS targets to inform RTP decisions.⁹⁹ Given the high demands a player is returning to, and the variation in physical demands due to playing position, there may be value in targeting individual within-session and acute (7 days) loads prior to RTT. Alongside this, individualised response to ‘load’ (i.e., subjective and objective measures of internal / external load and ‘fatigue monitoring’) and movement quality are important factors alongside training quantity.^{36,116} Training load has only recently been incorporated into models of injury aetiology^{70,103} despite being a risk factor for injury within a web of determinants.^{43,69} Future research should investigate this factor, acknowledging the role of ‘load’ within the complex nature of injuries, and further informing decision-making for RTT in an applied setting.

Within professional football, alongside GPS, the external load from specific technical actions can be quantified.¹⁸¹ This allows for the analysis of technical actions within rehabilitation in combination with physical output. Further research is required to assess physical and technical outputs, through drill level analysis and positional specific training demands, to help inform the training reintegration process. Achieving load through sport-specific exposure may be beneficial from a physical perspective,¹⁵⁵ improve physical-cognitive capabilities,³⁹ and impact the psychology/perception of the player’s readiness to return, potentially allowing an earlier RTT and RTP.

Injury surveillance and subsequent injury

An issue within applied rehabilitation is the lack of consensus and discrepancies in injury tracking/surveillance.^{157,159} These inconsistencies in reporting make it difficult to

research and understand subsequent relationships between injuries. Subsequent injury is a poorly defined and reported area within rehabilitation.¹⁵⁸ Further research is required to understand the inter-injury relationship and inform tertiary prevention programming.^{157,159,182} Furthermore, workload monitoring to assess a potential association between post-RTT load, player status (i.e., key, squad or development player) and subsequent injury risk would be beneficial. This may help inform practitioners as to possible optimal loading strategies post-RTT. Understanding subsequent injury risk may provide a timeframe for a ‘post-rehabilitation phase’ based on the severity of the index injury and improve the development of injury specific monitoring strategies for both reinjury, and possible subsequent injuries, to alternative sites.¹²² Furthermore, mechanism of subsequent injury is a further factor to investigate. Establishing a potential pattern of mechanisms may be beneficial to guide injury prevention programming. Previous injury and subsequent injury associations are rarely considered, despite previous injury modifying the complex interactions between determinants of injury.¹⁵⁸ Further studies are required into current practices within subsequent injury surveillance and specific athlete support post-RTT.

Player perceptions of rehabilitation processes

The role of multiple stakeholders within rehabilitation has been widely acknowledged,^{81,183} however, despite being part of the decision-making process,^{92,95} player perception and role within rehabilitation is not thoroughly discussed. A lack of consensus between players and coaches regarding the RTP decision has been previously noted.¹⁸⁴ Players are a key stakeholder in rehabilitation and anxiety, the need for a plan, evaluating the risk of early return, demonstrating progress and social support are important factors in the process.^{32,35} Furthermore, establishing athlete satisfaction post-injury requires further research.⁴³ Feedback of GPS training data within professional football has been shown to support the coaching process and understanding player feedback preference is key to elicit engagement.¹⁸⁵ However, little exists around RTP procedures and player perceptions regarding the use of specific GPS targets within the rehabilitation setting and feedback of this data.

Injury specific metrics

Different load metrics may have varying relationships to injury risk^{126,186–188} and the specific stress/strain capacity, resilience and adaptation of different tissues.^{46,165,189} In a survey of practitioners, “GPS metrics for certain injuries” was referred to as a useful future development in RTP protocols.³⁶ Quantifying the success of a rehabilitation process is an important consideration³⁶ and a continuum of importance for injury specific targets for different injury sites could be developed. These insights could give practitioners

greater understanding of key injury specific metrics for load management during rehabilitation and post-RTT, although this should be acknowledged within the complex nature of injury.

Limitations and perspectives

An acknowledged limitation of the article is the use of a narrative review and, therefore, a clear, systematic search methodology was not utilised resulting in potential bias.¹⁸⁶ The limitations of narrative reviews are documented,¹⁹⁰ however, using this method allowed for a wider search strategy exploring interdisciplinary topics and to reflect the complex nature of applied rehabilitation and decision-making for practitioners.

The review focuses on literature within the domain of male professional football, however, the frameworks presented may be further applicable to female professional football and developmental players. Research understanding the gender specific differences in injury and rehabilitation is growing,^{191,192} however, more is warranted specific to female football players and to academy players investigating the potential impact on long-term athlete development programming if an injury occurs. This may help to reduce disparity and improve understanding and application of the frameworks presented. Ultimately, a long-term objective of developing bespoke models for different sports, injury pathologies, age groups and genders, with multidisciplinary inputs, is crucial in future research and applied practices.

Conclusion

The purpose of this review was to give an overview of existing aetiological models, decision-making frameworks and RTS approaches within the current literature and evaluate the role of criteria-based progressions in rehabilitation frameworks and their application within male professional football. General sporting, and football specific models and frameworks are presented to demonstrate approaches and different interpretations of RTP and RTS terminology. Further consensus in terminology used to define phases, and subphases (i.e., RTT subphases), within the rehabilitation procedure is required to help guide intention and impact phase specific monitoring, and performance markers to inform safe return. By understanding injury type, mechanism, severity and location, the demands of the sport and multiple interacting risk factors, evidence-based RTP criteria can be used to guide decisions alongside practitioner intuitions and experiences. However, research of subsequent injury risk, and inter-injury relationships, is lacking in male professional football with current RTP frameworks using a stepwise progression through different rehabilitation phases and objective exit criteria to inform decision making. Development of flexible, bespoke frameworks, using validated objective criteria, is required and further

research is needed to ensure effective rehabilitation procedures, with successful RTP outcomes, to guide programming, and decision-making, that minimises subsequent injury risk. Through improved practitioner knowledge and understanding of aetiological models, decision-making and rehabilitation frameworks, and the future development of these topic areas, the support of injured athletes can be enhanced during, and following, the rehabilitation process.

Acknowledgements

Not applicable.

ORCID iDs

Ben Dixon  <https://orcid.org/0009-0009-0925-2304>

Jill Alexander  <https://orcid.org/0000-0002-6492-1621>

Damian Harper  <https://orcid.org/0000-0002-5430-1541>

Ethics considerations

Not applicable.

Consent to participate

Not applicable.

Consent for publication

Not applicable.

Author contributions

BD was responsible for the conception and writing of the review. DH and JA contributed and provided supervision and feedback throughout. All authors critically reviewed the manuscript and approved the final version.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Data availability

Not applicable.

References

1. Hägglund M, Waldén M, Magnusson H, et al. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA champions league injury study. *Br J Sports Med* 2013; 47: 38.
2. Eirale C, Tol JL, Farooq A, et al. Low injury rate strongly correlates with team success in Qatari professional football. *Br J Sports Med* 2013; 47: 807–808. 20120817.

3. McCall A, Lewin C, O'Driscoll G, et al. Return to play: the challenge of balancing research and practice. *Br J Sports Med* 2017; 51: 702–703.
4. Taberner M, Allen T and Cohen DD. Adapting the high chaos phase of the 'control-chaos continuum': a bridge to team training. *Sport Perform Sci Rep* 2020; 109: 1–6.
5. Ekstrand J, Häggglund M and Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011; 45: 53.
6. Häggglund M, Waldén M and Ekstrand J. Exposure and injury risk in Swedish elite football: a comparison between seasons 1982 and 2001. *Scand J Med Sci Sports* 2003; 13: 364–370.
7. Hawkins RD and Fuller CW. A prospective epidemiological study of injuries in four English professional football clubs. *Br J Sports Med* 1999; 33: 96.
8. Hawkins RD, Hulse MA, Wilkinson C, et al. The association football medical research programme: an audit of injuries in professional football. *Br J Sports Med* 2001; 35: 43–47.
9. Jones A, Jones G, Greig N, et al. Epidemiology of injury in English professional football players: a cohort study. *Phys Ther Sport* 2019; 35: 18–22.
10. López-Valenciano A, Ruiz-Pérez I, García-Gómez A, et al. Epidemiology of injuries in professional football: a systematic review and meta-analysis. *Br J Sports Med* 2020; 54: 711–718.
11. Palmer B, McBride M, Jones G, et al. Injury trends in men's English professional football: an 11-year case series. *J Elite Sport Perform* 2023; 3: 1–8.
12. Bengtsson H, Ortega Gallo PA and Ekstrand J. Injury epidemiology in professional football in South America compared with Europe. *BMJ Open Sport Exerc Med* 2021; 7: e001172.
13. Forsythe B, Knapik DM, Khazi-Syed D, et al. Analysis of injury epidemiology in soccer players in the 2019 confederation of north, Central America and Caribbean association football gold cup as reported by team physicians. *Arthrosc Sports Med Rehabil* 2025; 7: 101074.
14. Häggglund M, Waldén M and Ekstrand J. Previous injury as a risk factor for injury in elite football: a prospective study over two consecutive seasons. *Br J Sports Med* 2006; 40: 67.
15. Allen T, Taberner M, Zhilkin M, et al. Running more than before? The evolution of running load demands in the English Premier League. *Int J Sports Sci Coach* 2024; 19: 779–787.
16. Nassis GP, Massey A, Jacobsen P, et al. Elite football of 2030 will not be the same as that of 2020: preparing players, coaches, and support staff for the evolution. *Scand J Med Sci Sports* 2020; 30: 962–964.
17. Harper DJ, Sandford GN, Clubb J, et al. Elite football of 2030 will not be the same as that of 2020: what has evolved and what needs to evolve? *Scand J Med Sci Sports* 2021; 31: 493–494.
18. Della Villa F, Häggglund M, Della Villa S, et al. High rate of second ACL injury following ACL reconstruction in male professional footballers: an updated longitudinal analysis from 118 players in the UEFA elite club injury study. *Br J Sports Med* 2021; 55: 1350–1357.
19. Buckthorpe M, Frizziero A and Roi GS. Update on functional recovery process for the injured athlete: return to sport continuum redefined. *Br J Sports Med* 2019; 53: 265–267.
20. Ardern CL, Kvist J and Webster KE. Psychological aspects of anterior cruciate ligament injuries. *Oper Tech Sports Med* 2016; 24: 77–83.
21. Barth KA, Lawton CD, Touhey DC, et al. The negative impact of anterior cruciate ligament reconstruction in professional male footballers. *Knee* 2019; 26: 142–148.
22. Niederer D, Engeroff T, Wilke J, et al. Return to play, performance, and career duration after anterior cruciate ligament rupture: a case-control study in the five biggest football nations in Europe. *Scand J Med Sci Sports* 2018; 28: 2226–2233. 20180709.
23. Stone S. Average top-flight manager tenure across Europe less than 16 months, says Uefa, <https://www.bbc.co.uk/sport/football/66794395> (2023, accessed 4 September 2024).
24. Lewin G and Lewin C. The changing landscape of football medicine. *Aspetar Sports Med J* 2018; 7: 88–95.
25. Eliakim E, Morgulev E, Lidor R, et al. Estimation of injury costs: financial damage of English premier league teams' underachievement due to injuries. *BMJ Open Sport Exerc Med* 2020; 6: e000675. 20200520.
26. Nieto Torrejón L, Martínez-Serrano A, Villalón JM, et al. Economic impact of muscle injury rate and hamstring strain injuries in professional football clubs. *Evidence from LaLiga. PLOS ONE* 2024; 19: e0301498.
27. Oliveira-Júnior O, Gabbett TJ, Bittencourt NFN, et al. Potential financial loss and risk factors for hamstring muscle injuries in elite male Brazilian soccer players: a season-long prospective cohort pilot study. *Front Sports Act Living* 2024; 6: 1–10.
28. Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Clin J Sport Med* 2006; 16: 97–106.
29. Häggglund M, Waldén M and Ekstrand J. Injury recurrence is lower at the highest professional football level than at national and amateur levels: does sports medicine and sports physiotherapy deliver? *Br J Sports Med* 2016; 50: 51.
30. Flore Z, Hambly K, De Coninck K, et al. A rehabilitation algorithm after lateral ankle sprains in professional football (soccer): an approach based on clinical practice guidelines. *Int J Sports Phys Ther* 2024; 19: 910–922.
31. Ekstrand J, Krutsch W, Spreco A, et al. Time before return to play for the most common injuries in professional football: a 16-year follow-up of the UEFA elite club injury study. *Br J Sports Med* 2020; 54: 421–426.
32. Taberner M, Distain S, Chance O, et al. Clinical pearls on how practitioners can best support elite soccer players to

- return to peak performance after injury. *JOSPT Cases* 2021; 1: 126–128.
33. Forsdyke D, Gledhill A and Ardern CL. Psychological readiness to return to sport: three key elements to help the practitioner decide whether the athlete is REALLY ready? *Br J Sports Med* 2017; 51: 555–556.
 34. Forsdyke D, Madigan D, Gledhill A, et al. Perceived social support, reinjury anxiety, and psychological readiness to return to sport in soccer players. *J Sport Rehabil* 2022; 31: 749–755.
 35. Tranaeus U, Gledhill A, Johnson U, et al. 50 Years of research on the psychology of sport injury: a consensus statement. *Sports Med* 2024; 54: 1733–1748.
 36. Armitage M, McErlain-Naylor SA, Devereux G, et al. On-field rehabilitation in football: current practice and perceptions. A survey of the English premier league and football league. *Sci Med Footb* 2025; 9: 130–139.
 37. Van Melick N, Van Cingel REH, Brooijmans F, et al. Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus. *Br J Sports Med* 2016; 50: 1506–1515.
 38. Taberner M, Allen T and Cohen DD. Progressing rehabilitation after injury: consider the ‘control-chaos continuum’. *Br J Sports Med* 2019; 53: 1132.
 39. Taberner M, Allen T, O’Keefe J, et al. Evolving the control-chaos continuum: part 1 – translating knowledge to enhance on-pitch rehabilitation. *J Orthop Sports Phys Ther* 2025; 55: 78–88.
 40. Taberner M, Allen T, O’Keefe J, et al. Evolving the control-chaos continuum: part 2 – shifting ‘attention’ to progress on-pitch rehabilitation. *J Orthop Sports Phys Ther* 2025; 55: 162–172.
 41. Buckthorpe M, Della Villa F, Della Villa S, et al. On-field rehabilitation part 2: a 5-stage program for the soccer player focused on linear movements, multidirectional movements, soccer-specific skills, soccer-specific movements, and modified practice. *J Orthop Sports Phys Ther* 2019; 49: 570–575. 20190710.
 42. Mitchell A and Gimpel M. A return to performance pathway for professional soccer: a criteria based approach to return injured professional players back to performance. *JOSPT Open* 2024; 2: 166–178.
 43. Ardern CL, Glasgow P, Schneiders A, et al. 2016 Consensus statement on return to sport from the first world congress in sports physical therapy, Bern. *Br J Sports Med* 2016; 50: 853–864.
 44. Taberner M, Allen T, Constantine E, et al. From control to chaos to competition: building a pathway for return to performance following ACL reconstruction. *Aspetar Sports Med J* 2020; 9: 84–94.
 45. Woolley TGD, Dick S and Reid D. Mechanism of injury. In: P Lax (eds) *Textbook of acute trauma care*. Cham: Springer International Publishing, 2022, pp.421–433.
 46. Kalkhoven JT, Watsford ML and Impellizzeri FM. A conceptual model and detailed framework for stress-related, strain-related, and overuse athletic injury. *J Sci Med Sport* 2020; 23: 726–734.
 47. Edwards WB. Modeling overuse injuries in sport as a mechanical fatigue phenomenon. *Exerc Sport Sci Rev* 2018; 46: 224–231.
 48. Watson T. Soft tissue wound healing review. 2003.
 49. Armitage M, McErlain-Naylor SA, Devereux G, et al. On-field rehabilitation in football: current knowledge, applications and future directions. *Front Sports Active Living* 2022; 4: 970152.
 50. Roe M, Malone S, Blake C, et al. A six stage operational framework for individualising injury risk management in sport. *Inj Epidemiol* 2017; 4: 26.
 51. McCall A, Carling C, Nedelec M, et al. Risk factors, testing and preventative strategies for non-contact injuries in professional football: current perceptions and practices of 44 teams from various premier leagues. *Br J Sports Med* 2014; 48: 1352–1357. 20140516.
 52. Ettema JH. Het model belasting en belastbaarheid. *T Soc Geneesk* 1973; 51: 44–54.
 53. van Mechelen W, Hlobil H and Kemper HC. How can sports injuries be prevented? *Nationaal Instituut voor Sport-GezondheidsZorg publicatie* 1987; 25E.
 54. Finch CF. A new framework for research leading to sports injury prevention. *J Sci Med Sport* 2006; 9: 3–9; discussion 10. 20060417.
 55. Bolling C, Van Mechelen W, Pasman HR, et al. Context matters: revisiting the first step of the ‘sequence of prevention’ of sports injuries. *Sports Med* 2018; 48: 2227–2234.
 56. Andersen MB and Williams JM. A model of stress and athletic injury: prediction and prevention. *J Sport Exerc Psychol* 1988; 10: 294–306.
 57. van Dijk F, van Dormolen M, Kompier M, et al. Herwaardering model belasting-belastbaarheid. *Tijdschr Soc Gezondheidsz* 1990; 68: 3–10.
 58. van Mechelen W, Hlobil H and Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992; 14: 82–99.
 59. Backx F, Inklaar H, Koornneef M, et al. FIMS Position statement on the prevention of sports injuries. *Geneeskunde en Sport (Special Issue)* 1990; 23: 22–27.
 60. Bol E, Schmikli SL, Backx FJG, et al. Sportblessures onder de knie: NISGZ publicatie no. 38. In: 1991.
 61. Kok G and Bouter LM. On the importance of planned health education. *Am J Sports Med* 1990; 18: 600–605.
 62. Meeuwisse WH. Assessing causation in sport injury: a multifactorial model. *Clin J Sport Med* 1994; 4: 166–170.
 63. Williams JM and Andersen MB. Psychosocial antecedents of sport injury: review and critique of the stress and injury model’. *J Appl Sport Psychol* 1998; 10: 5–25.
 64. McIntosh AS. Risk compensation, motivation, injuries, and biomechanics in competitive sport. *Br J Sports Med* 2005; 39: 2–3.

65. Bahr R and Krosshaug T. Understanding injury mechanisms: a key component of preventing injuries in sport. *Br J Sports Med* 2005; 39: 24.
66. Meeuwisse WH, Tyreman H, Hagel BE, et al. A dynamic model of etiology in sport injury: the recursive nature of risk and causation. *Clin J Sport Med* 2007; 17: 215–219.
67. Gissane C, White J, Kerr K, et al. An operational model to investigate contact sports injuries. *Med Sci Sports Exerc* 2001; 33: 1999–2003.
68. Appaneal RN and Perna FM. Biopsychosocial model of injury. In: . *Encyclopedia of sport and exercise psychology*. Thousand Oaks: Sage Publications Inc, 2014, pp.74–77.
69. Bittencourt NFN, Meeuwisse WH, Mendonca LD, et al. Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition-narrative review and new concept. *Br J Sports Med* 2016; 50: 1309.
70. Windt J and Gabbett TJ. How do training and competition workloads relate to injury? The workload-injury aetiology model. *Br J Sports Med* 2017; 51: 28.
71. O'Brien J, Finch CF, Pruna R, et al. A new model for injury prevention in team sports: the team-sport injury prevention (TIP) cycle. *Sci Med Footb* 2019; 3: 77–80.
72. Liveris NI. Applying systems thinking approaches to investigate the complex interrelationships of risk factors affecting acute non-contact lower limb injuries in team sports (PhD Academy Award). *Br J Sports Med* 2025; bjsports-2025-2021: 683–684.
73. Liveris NI, Papageorgiou G, Tsepis E, et al. Towards the development of a system dynamics model for the prediction of lower extremity injuries. *Int J Exerc Sci* 2023; 16: 1052–1065.
74. Liveris NI, Tsarbou C, Papageorgiou G, et al. The complex interrelationships of the risk factors leading to hamstring injury and implications for injury prevention: a group model building approach. *Appl Sci* 2024; 14: 6316.
75. Malone S, Owen A, Mendes B, et al. High-speed running and sprinting as an injury risk factor in soccer: can well-developed physical qualities reduce the risk? *J Sci Med Sport* 2018; 21: 257–262.
76. Malone S, Roe M, Doran DA, et al. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite gaelic football. *J Sci Med Sport* 2017; 20: 250–254.
77. Taberner M, Allen T, O'Keefe J, et al. Contextual considerations using the 'control-chaos continuum' for return to sport in elite football - part 1: load planning. *Phys Ther Sport* 2022; 53: 67–74. 20211103.
78. Page RM, Field A, Langley B, et al. The effects of fixture congestion on injury in professional male soccer: a systematic review. *Sports Med* 2023; 53: 667–685.
79. Brito J, Mendes R, Figueiredo P, et al. Is it time to consider quaternary injury prevention in sports? *Sports Med* 2023; 53: 769–774.
80. Ekstrand J, Lundqvist D, Lagerback L, et al. Is there a correlation between coaches' leadership styles and injuries in elite football teams? A study of 36 elite teams in 17 countries. *Br J Sports Med* 2018; 52: 527–531. 20171022.
81. Ekstrand J, Lundqvist D, Davison M, et al. Communication quality between the medical team and the head coach/manager is associated with injury burden and player availability in elite football clubs. *Br J Sports Med* 2019; 53: 304–308.
82. Hassanmirzaei B, Schumacher Y, Tabben M, et al. Developing a data-driven multimodal injury and illness prevention programme in male professional football based on a risk management model: the IP2 NetWork. *BMJ Open Sport Exerc Med* 2024; 10: e002101.
83. King R, McHugh D, Alexander J, et al. Multi-disciplinary team practitioners working in high performance sport: skilled intuitive 'doers' or novel problem-solving innovators. *Eur J Sports Sci* 2024; 3: 15–26.
84. King R, Yiannaki C, Kiely J, et al. Multi-disciplinary teams in high performance sport, the what and the how: a utopian view or a darker reality. *J Expertise* 2024; 7: 149–174.
85. Ekstrand J, Häggglund M, Waldén M, et al. Higher level of communication between the medical staff and the performance staff is associated with a lower hamstring injury burden: a substudy on 14 teams from the UEFA elite club injury study. *BMJ Open Sport & Exercise Medicine* 2025; 11: e002182.
86. Ardern CL, Bizzini M and Bahr R. It is time for consensus on return to play after injury: five key questions. *Br J Sports Med* 2016; 50: 506–508.
87. Shrier I, Safai P and Charland L. Return to play following injury: whose decision should it be? *Br J Sports Med* 2014; 48: 94.
88. Shrier I. Strategic assessment of risk and risk tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015; 49: 1311–1315.
89. Blanch P and Gabbett TJ. Has the athlete trained enough to return to play safely? The acute:chronic workload ratio permits clinicians to quantify a player's risk of subsequent injury. *Br J Sports Med* 2016; 50: 71.
90. Gabbett TJ, Hulin BT, Blanch P, et al. High training workloads alone do not cause sports injuries: how you get there is the real issue. *Br J Sports Med* 2016; 50: 444–445.
91. Creighton DW, Shrier I, Shultz R, et al. Return-to-play in sport: a decision-based model. *Clin J Sport Med* 2010; 20: 379–385.
92. Dijkstra HP, Pollock N, Chakraverty R, et al. Return to play in elite sport: a shared decision-making process. *Br J Sports Med* 2017; 51: 19.
93. Yung KK, Ardern CL, Serpiello FR, et al. A framework for clinicians to improve the decision-making process in return to sport. *Sports Med - Open* 2022; 8: 52.
94. Yung KK, Ardern CL, Serpiello FR, et al. Characteristics of complex systems in sports injury rehabilitation: examples

- and implications for practice. *Sports Med - Open* 2022; 8: 24.
95. King J, Roberts C, Hard S, et al. Want to improve return to sport outcomes following injury? Empower, engage, provide feedback and be transparent: 4 habits!. *Br J Sports Med* 2019; 53: 26.
 96. Buchheit M, King R, Stokes A, et al. Return to play following injuries in pro football: insights into the real-life practices of 85 elite practitioners around diagnostics, progression strategies and reintegration processes. *Sport Perform Sci Rep* 2023; 1: 1–20.
 97. Dunlop G, Ardern CL, Andersen TE, et al. Return-to-play practices following hamstring injury: a worldwide survey of 131 Premier League football teams. *Sports Med* 2020; 50: 829–840.
 98. Draovitch P, Patel S, Marrone W, et al. The return-to-sport clearance continuum is a novel approach toward return to sport and performance for the professional athlete. *Arthrosc Sports Med Rehabil* 2022; 4: e93–e101.
 99. Van Der Horst N, Backx F, Goedhart EA, et al. Return to play after hamstring injuries in football (soccer): a worldwide Delphi procedure regarding definition, medical criteria and decision-making. *Br J Sports Med* 2017; 51: 1583–1591.
 100. Bisciotti GN, Volpi P, Alberti G, et al. Italian consensus statement (2020) on return to play after lower limb muscle injury in football (soccer). *BMJ Open Sport Exerc Med* 2019; 5: e000505.
 101. Gabbett TJ, Whyte DG, Hartwig TB, et al. The relationship between workloads, physical performance, injury and illness in adolescent male football players. *Sports Med* 2014; 44: 989–1003.
 102. Hickey JT, Timmins RG, Maniar N, et al. Criteria for progressing rehabilitation and determining return-to-play clearance following hamstring strain injury: a systematic review. *Sports Med* 2017; 47: 1375–1387.
 103. Taberner M, Cohen DD, Carter A, et al. ‘Where is the load?’ Revisiting the strategic assessment of risk and risk tolerance (StARRT) framework for return to sport by including an athlete’s sport-specific training capacity? *Br J Sports Med* 2022; 56: 832–834.
 104. Zambaldi M, Beasley I and Rushton A. Return to play criteria after hamstring muscle injury in professional football: a delphi consensus study. *Br J Sports Med* 2017; 51: 1221–1226. 20170228.
 105. Paton BM, Read P, Van Dyk N, et al. London International consensus and delphi study on hamstring injuries part 3: rehabilitation, running and return to sport. *Br J Sports Med* 2023; 57: 278–291.
 106. Yung KK, Ardern CL, Serpiello FR, et al. Judgement and decision making in clinical and return-to-sports decision making: a narrative review. *Sports Med* 2024; 54: 2005–2017.
 107. Mateus N, Abade E, Coutinho D, et al. Empowering the sports scientist with artificial intelligence in training, performance, and health management. *Sensors (Basel)* 2025; 25: 20241229.
 108. Bullock GS, Mylott J, Hughes T, et al. Just how confident can we be in predicting sports injuries? A systematic review of the methodological conduct and performance of existing musculoskeletal injury prediction models in sport. *Sports Med* 2022; 52: 2469–2482.
 109. Bullock GS, Hughes T, Sergeant JC, et al. Clinical prediction models in sports medicine: a guide for clinicians and researchers. *J Orthop Sports Phys Ther* 2021; 51: 517–525.
 110. Leckey C, Van Dyk N, Doherty C, et al. Machine learning approaches to injury risk prediction in sport: a scoping review with evidence synthesis. *Br J Sports Med* 2024; 59: 491–500.
 111. Majumdar A, Bakirov R, Hodges D, et al. Machine Learning for understanding and predicting injuries in football. *Sports Med - Open* 2022; 8: 73.
 112. Van Eetvelde H, Mendonça LD, Ley C, et al. Machine learning methods in sport injury prediction and prevention: a systematic review. *J Exp Orthop* 2021; 8: 27.
 113. Mitchell A, Waite O, Holding C, et al. The development of a return to performance pathway involving a professional soccer player returning from a multi-structural knee injury: a case report. *Int J Sports Phys Ther* 2023; 18: 450–466.
 114. Mendiguchia J, Martinez-Ruiz E, Edouard P, et al. A multi-factorial, criteria-based progressive algorithm for hamstring injury treatment. *Med Sci Sports Exerc* 2017; 49: 1482–1492.
 115. Estévez Rodríguez JL, Rivilla García J, Jiménez-Sáiz SL, et al. Consensus of return-to-play criteria after adductor longus injury in professional soccer. *Sports* 2025; 13: 34.
 116. Buckthorpe M, Della Villa F, Della Villa S, et al. On-field rehabilitation part 1: 4 pillars of high-quality on-field rehabilitation are restoring movement quality, physical conditioning, restoring sport-specific skills, and progressively developing chronic training load. *J Orthop Sports Phys Ther* 2019; 49: 565–569. 20190710.
 117. Jiménez-Rubio S, Estévez Rodríguez JL and Navandar A. Validity of a rehab and reconditioning program following an adductor longus injury in professional soccer. *J Sport Rehabil* 2021; 30: 1224–1229.
 118. Jimenez-Rubio S, Navandar A, Rivilla-Garcia J, et al. Validity of an on-field readaptation program following a hamstring injury in professional soccer. *J Sport Rehabil* 2019; 28: jsr.2018-0203.
 119. Stathas I, Kalliakmanis A, Kekelekis A, et al. Effectiveness of an on-field rehabilitation framework for return to sports in injured male professional football players: a single-blinded, prospective, randomised controlled trial. *BMJ Open Sport Exerc Med* 2024; 10: e001849. 20240119..
 120. Ekstrand J, Spreco A, Bengtsson H, et al. Injury rates decreased in men’s professional football: an 18-year prospective cohort study of almost 12 000 injuries sustained

- during 1.8 million hours of play. *Br J Sports Med* 2021; 55: 1084–1091.
121. Zhang G, Brink MS, Aus Der Fünten K, et al. The time course of injury risk after return-to-play in professional football (Soccer). *Sports Med* 2025; 55: 193–201.
 122. Dixon B, Alexander J and Harper D. ‘Post-rehabilitation phase’ in professional football: are we optimising player support after return to play? *Br J Sports Med* 2025; 59: 625–627.
 123. Taberner M and Cohen DD. Physical preparation of the football player with an intramuscular hamstring tendon tear: clinical perspective with video demonstrations. *Br J Sports Med* 2018; 52: 1275.
 124. Taberner M, O’Keefe J, Dunn A, et al. Return to sport and beyond following intramuscular tendon hamstring injury: a case report of an English Premier League football player. *Phys Ther Sport* 2022; 56: 38–47. 20220531.
 125. Taberner M, van Dyk N, Allen T, et al. Physical preparation and return to sport of the football player with a tibia-fibula fracture: applying the ‘control-chaos continuum’. *BMJ Open Sport Exerc Med* 2019; 5: e000639. 20191030.
 126. Impellizzeri FM, Ward P, Coutts AJ, et al. Training load and injury part 2: questionable research practices hijack the truth and mislead well-intentioned clinicians. *J Orthop Sports Phys Ther* 2020; 50: 577–584.
 127. Fanchini M, Steendahl IB, Impellizzeri FM, et al. Exercise-based strategies to prevent muscle injury in elite footballers: a systematic review and best evidence synthesis. *Sports Med* 2020; 50: 1653–1666.
 128. Benson LC, Räisänen AM, Volkova VG, et al. Workload a-WEAR-ness: monitoring workload in team sports with wearable technology. A scoping review. *J Orthop Sports Phys Ther* 2020; 50: 549–563.
 129. Lolli L, Batterham AM, Hawkins R, et al. The acute-to-chronic workload ratio: an inaccurate scaling index for an unnecessary normalisation process? *Br J Sports Med* 2019; 53: 1510–1512. 20180613.
 130. Impellizzeri FM, Woodcock S, Coutts AJ, et al. What role do chronic workloads play in the acute to chronic workload ratio? Time to dismiss ACWR and its underlying theory. *Sports Med* 2021; 51: 581–592.
 131. Impellizzeri FM, Woodcock S, Coutts AJ, et al. Acute to random workload ratio is ‘as’ associated with injury as acute to actual chronic workload ratio: time to dismiss ACWR and its components. SportRxiv 2020.
 132. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016; 50: 73.
 133. Coetzee D, Wall C, De Bruin M, et al. Return to play and performance guidelines in rugby union. *S Afr J Res Sport Phys Ed Recreat* 2023; 45: 39–61.
 134. Akenhead R and Nassis GP. Training load and player monitoring in high-level football: current practice and perceptions. *Int J Sports Physiol Perform* 2016; 11: 587–593.
 135. Pillay L, Van Rensburg DCJ, Ramkilawon G, et al. Don’t forget to mind the mind: a prospective cohort study over 12 months on mental health symptoms in active professional male footballers. *BMC Sports Sci Med Rehabil* 2024; 16: 214.
 136. Smith MR, Thompson C, Marcora SM, et al. Mental fatigue and soccer: current knowledge and future directions. *Sports Med* 2018; 48: 1525–1532.
 137. González-Villora S, Prieto-Ayuso A, Cardoso F, et al. The role of mental fatigue in soccer: a systematic review. *Int J Sports Sci Coach* 2022; 17: 903–916.
 138. Buckthorpe M. Optimising the late-stage rehabilitation and return-to-sport training and testing process after ACL reconstruction. *Sports Med* 2019; 49: 1043–1058.
 139. Bradley PS, Carling C, Archer D, et al. The effect of playing formation on high-intensity running and technical profiles in English FA Premier League soccer matches. *J Sports Sci* 2011; 29: 821–830.
 140. Bradley PS, Carling C, Gomez Diaz A, et al. Match performance and physical capacity of players in the top three competitive standards of English professional soccer. *Hum Mov Sci* 2013; 32: 808–821. 20130824.
 141. Di Mascio M and Bradley PS. Evaluation of the most intense high-intensity running period in English FA premier league soccer matches. *J Strength Cond Res* 2013; 27: 909–915.
 142. Harper DJ, Carling C and Kiely J. High-intensity acceleration and deceleration demands in elite team sports competitive match play: a systematic review and meta-analysis of observational studies. *Sports Med* 2019; 49: 1923–1947.
 143. Hostrup M and Bangsbo J. Performance adaptations to intensified training in top-level football. *Sports Med* 2023; 53: 577–594.
 144. Mohr M, Krustup P and Bangsbo J. Match performance of high-standard soccer players with special reference to development of fatigue. *J Sports Sci* 2003; 21: 519–528.
 145. Morgans R, Radnor J, Fonseca J, et al. Match running performance is influenced by possession and team formation in an English Premier League team. *Biol Sport* 2024; 41: 275–286. 20240212.
 146. Chaize C, Allen M and Beato M. Physical performance is affected by players’ position, game location, and substitutions during official competitions in professional championship English football. *J Strength Cond Res* 2024; 38: e744–e753.
 147. Bradley PS, Archer DT, Hogg B, et al. Tier-specific evolution of match performance characteristics in the English Premier League: it’s getting tougher at the top. *J Sports Sci* 2016; 34: 980–987.
 148. Faude O, Koch T and Meyer T. Straight sprinting is the most frequent action in goal situations in professional football. *J Sports Sci* 2012; 30: 625–631. 20120306.
 149. Sarajärvi J, Volossovitch A and Almeida CH. Analysis of headers in high-performance football: evidence from the

- English Premier League. *Int J Perform Anal Sport* 2020; 20: 189–205.
150. Rhodes D, Valassakis S, Bortnik L, et al. The effect of high-intensity accelerations and decelerations on match outcome of an elite English league two football team. *Int J Environ Res Public Health* 2021; 18: 9913.
151. Martínez-Hernández D, Quinn M and Jones P. Linear advancing actions followed by deceleration and turn are the most common movements preceding goals in male professional soccer. *Sci Med Footb* 2023; 7: 25–33.
152. Rampinini E, Impellizzeri FM, Castagna C, et al. Technical performance during soccer matches of the Italian Serie A league: effect of fatigue and competitive level. *J Sci Med Sport* 2009; 12: 227–233.
153. Eliakim E, Doron O, Meckel Y, et al. Pre-season fitness level and injury rate in professional soccer – a prospective study. *Sports Med Int Open* 2018; 02: E84–E90.
154. Bengtsson H, Ekstrand J, Walden M, et al. Few training sessions between return to play and first match appearance are associated with an increased propensity for injury: a prospective cohort study of male professional football players during 16 consecutive seasons. *Br J Sports Med* 2020; 54: 427–432. 20190829.
155. Stares J, Dawson B, Peeling P, et al. How much is enough in rehabilitation? High running workloads following lower limb muscle injury delay return to play but protect against subsequent injury. *J Sci Med Sport* 2018; 21: 1019–1024. 20180410.
156. Finch CF, Cook J, Kunstler BE, et al. Subsequent injuries are more common than injury recurrences: an analysis of 1 season of prospectively collected injuries in professional Australian football. *Am J Sports Med* 2017; 45: 1921–1927.
157. Pucciarelli A, Swain M, Lystad R, et al. Subsequent and recurrent injuries in football players: a systematic review. *J Sci Med Sport* 2023; 26: S107–S108.
158. Toohey LA, Drew MK, Cook JL, et al. Is subsequent lower limb injury associated with previous injury? A systematic review and meta-analysis. *Br J Sports Med* 2017; 51: 1670–1678. 20170807.
159. Bitchell CL, Varley-Campbell J, Robinson G, et al. Recurrent and subsequent injuries in professional and elite sport: a systematic review. *Sports Med - Open* 2020; 6: 58.
160. Cranswick I, Jones A, Brogden C, et al. Rehabilitation remodelled: a narrative review of injury rehabilitation models and proposal of a multi-component MSK rehabilitation model. *Res Sports Med* 2025: 1–29. doi: 10.1080/15438627.2025.2547191
161. Bleakley C, Klempel N, Wagemans J, et al. Statistical power in musculoskeletal research: A meta-review of 266 randomised controlled trials. *Sports Med - Open* 2025; 11: 134.
162. Estévez Rodríguez JL, Rivilla García J and Jiménez-Rubio S. Return to performance of a soccer player with an adductor longus injury: a case report. *Medicina (B Aires)* 2024; 60: 1998.
163. Picinini F, Della Villa F, Tallent J, et al. High return to competition rate after on-field rehabilitation in competitive male soccer players after ACL reconstruction: GPS tracking in 100 consecutive cases. *Orthop J Sports Med* 2025; 13. doi: 10.1177/23259671251320093
164. McCall A, Fanchini M and Coutts AJ. Prediction: the modern-day sport-science and sports-medicine “quest for the holy grail”. *Int J Sports Physiol Perform* 2017; 12: 704–706.
165. Kalkhoven JT, Watsford ML, Coutts AJ, et al. Training load and injury: causal pathways and future directions. *Sports Med* 2021; 51: 1137–1150.
166. Coutts AJ. Challenges in developing evidence-based practice in high-performance sport. *Int J Sports Physiol Perform* 2017; 12: 717–718.
167. Buchheit M, Settembre M, Hader K, et al. From high-speed running to hobbling on crutches: a machine learning perspective on the relationships between training doses and match injury trends. *Sport Perform Sci Rep* 2023; 216: 1–11.
168. Impellizzeri FM, Menaspà P, Coutts AJ, et al. Training load and its role in injury prevention, part i: back to the future. *J Athl Train* 2020; 55: 885–892.
169. Shergill AS, Twist C and Highton J. Importance of GNSS data quality assessment with novel control criteria in professional soccer match-play. *Int J Perform Anal Sport* 2021; 21: 820–830.
170. Theodoropoulos JS, Bettle J and Kosy JD. The use of GPS and inertial devices for player monitoring in team sports: a review of current and future applications. *Orthop Rev* 2020; 12: 7863.
171. Johnston RJ, Watsford ML, Kelly SJ, et al. Validity and interunit reliability of 10 Hz and 15 Hz GPS units for assessing athlete movement demands. *J Strength Cond Res* 2014; 28: 1649–1655.
172. Delaney JA, Wileman TM, Perry NJ, et al. The validity of a global navigation satellite system for quantifying small-area team-sport movements. *J Strength Cond Res* 2019; 33: 1463–1466.
173. Ellens S, Carey D, Gastin P, et al. Changing the criteria applied to acceleration and deceleration efforts changes the types of player actions detected. *Sci Med Footb* 2024; 8: 52–59.
174. Ellens S, Middleton K, Gastin PB, et al. Techniques to derive and clean acceleration and deceleration data of athlete tracking technologies in team sports: a scoping review. *J Sports Sci* 2022; 40: 1772–1800.
175. Buchheit M and Simpson BM. Player-tracking technology: half-full or half-empty glass? *Int J Sports Physiol Perform* 2017; 12: S2-35-S32-41.
176. Delaney JA, Cummins CJ, Thornton HR, et al. Importance, reliability, and usefulness of acceleration measures in team sports. *J Strength Cond Res* 2018; 32: 3485–3493.
177. Varley MC, Fairweather IH and Aughey RJ. Validity and reliability of GPS for measuring instantaneous velocity during acceleration, deceleration, and constant motion. *J Sports Sci* 2012; 30: 121–127. 20111129.

178. Fitzwilliam E, Steventon-Lorenzen N, Opar D, et al. Lower limb joint mechanics during maximal accelerative and decelerative running. *Med Sci Sports Exerc* 2024; 56: 1655–1663.
179. Verheul J, Harper D and Robinson MA. Forces experienced at different levels of the musculoskeletal system during horizontal decelerations. *J Sports Sci* 2024; 42: 2242–2253.
180. McBurnie AJ, Harper DJ, Jones PA, et al. Deceleration training in team sports: another potential ‘vaccine’ for sports-related injury? *Sports Med* 2022; 52: 1–12.
181. Lewis G, Towlson C, Roversi P, et al. Quantifying volume and high-speed technical actions of professional soccer players using foot-mounted inertial measurement units. *PLOS ONE* 2022; 17: e0263518.
182. Toohey LA, Drew MK, Fortington LV, et al. Comparison of subsequent injury categorisation (SIC) models and their application in a sporting population. *Inj Epidemiol* 2019; 6: 9.
183. Ekstrand J. Keeping your top players on the pitch: the key to football medicine at a professional level. *Br J Sports Med* 2013; 47: 23.
184. Loose O, Achenbach L, Fellner B, et al. Injury prevention and return to play strategies in elite football: no consent between players and team coaches. *Arch Orthop Trauma Surg* 2018; 138: 985–992.
185. Nosek P, Brownlee TE, Drust B, et al. Feedback of GPS training data within professional English soccer: a comparison of decision making and perceptions between coaches, players and performance staff. *Sci Med Footb* 2021; 5: 35–47.
186. Bowen L, Gross AS, Gimpel M, et al. Spikes in acute:chronic workload ratio (ACWR) associated with a 5–7 times greater injury rate in English Premier League football players: a comprehensive 3-year study. *Br J Sports Med* 2020; 54: 731–738.
187. Bowen L, Gross AS, Gimpel M, et al. Accumulated workloads and the acute:chronic workload ratio relate to injury risk in elite youth football players. *Br J Sports Med* 2017; 51: 452–459.
188. Jaspers A, Kuyvenhoven JP, Staes F, et al. Examination of the external and internal load indicators’ association with overuse injuries in professional soccer players. *J Sci Med Sport* 2018; 21: 579–585.
189. Gabbett TJ and Oetter E. From tissue to system: what constitutes an appropriate response to loading? *Sports Med* 2025; 55: 17–35.
190. Impellizzeri FM and Bizzini M. Systematic review and meta-analysis: a primer. *Int J Sports Phys Ther* 2012; 7: 493–503.
191. Abed V, Dupati A, Hawk GS, et al. Return to play and performance after anterior cruciate ligament reconstruction in the national women’s soccer league. *Orthop J Sports Med* 2023; 11: 232596712311649.
192. Hallén A, Tomás R, Ekstrand J, et al. UEFA Women’s elite club injury study: a prospective study on 1527 injuries over four consecutive seasons 2018/2019 to 2021/2022 reveals thigh muscle injuries to be most common and ACL injuries most burdensome. *Br J Sports Med* 2024; 58: 128–136.