

Central Lancashire Online Knowledge (CLOK)

Title	Using Visible Implant Elastomer to tag insects across life stages: a preliminary investigation with blow flies (Diptera: Calliphoridae)
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/8687/
DOI	https://doi.org/10.4039/tce.2013.18
Date	2013
Citation	Moffatt, Colin (2013) Using Visible Implant Elastomer to tag insects across life stages: a preliminary investigation with blow flies (Diptera: Calliphoridae). The Canadian Entomologist, 145 (-). pp. 466-470. ISSN 0008-347X
Creators	Moffatt, Colin

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.4039/tce.2013.18>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Using Visible Implant Elastomer to tag insects across life stages: a preliminary investigation with blow flies (Diptera: Calliphoridae)

Colin Moffatt

Abstract—Visible Implant Elastomer (VIE) has previously been used successfully to tag individuals in a variety of marine and amphibious animals, earthworms, and scorpions. Visible Implant Elastomer tags were injected into third instars of the blow fly *Calliphora vicina* Robineau-Desvoidy (Diptera: Calliphoridae) to test its compatibility and retention across life stages. Injecting into the dorsal midline of the 11th segment (seventh abdominal segment) produced survival rates of 80%, with no significant difference in the subsequent rate of development ($z = 0.21$, $P = 0.83$) as compared with untagged insects. Tags remained visible and allowed identification of individuals within a feeding, intermingling aggregation (maggot mass), especially when a high-contrast fluorescent colour was used. Tags were retained across life-stage changes and were easily found in dissected adults.

Résumé—Les implants visibles d'élastomère (VIE, Visible Implant Elastomer) ont servi avec succès à étiqueter individuellement une variété d'animaux marins et amphibies, des vers de terre et des scorpions. Des étiquettes visibles d'élastomère ont été injectées dans des larves de troisième stade de la mouche de la viande *Calliphora vicina* Robineau-Desvoidy (Diptera: Calliphoridae) pour évaluer leur compatibilité et leur rétention au cours des différents stades du cycle. L'injection dans la ligne dorsale médiane du 11^e segment (7^e segment abdominal) s'accompagne de taux de survie de 80% et les taux subséquents de développement ($z = 0,21$, $P = 0,83$) des insectes marqués ne diffèrent pas significativement de ceux des insectes nonmarqués. Les étiquettes demeurent visibles et permettent l'identification d'individus au sein d'une agrégation alimentaire (masse entremêlée d'asticots), particulièrement lorsqu'on utilise une couleur fluorescente à fort contraste. Les étiquettes sont retenues durant les changements du cycle biologique et se retrouvent facilement par dissection chez les adultes.

The ability to distinguish individual animals in a population has helped to progress many areas of zoology, and where natural appearance does not facilitate this, artificial marking or tagging can be a great help. Henderson (2003) defines “tagging” in relation to identifying animals individually, whereas “marking” is an umbrella term which also includes batch marking of cohorts.

A variety of marking methods has been used in population studies of insects (Southwood and Henderson 2000; Hagler and Jackson 2001), particularly to distinguish adult insects. Such techniques may involve the ingestion of batch-marking media as larvae, including protein markers (Hagler 1997), radio isotopes (Service 1993), and rubidium (Berry *et al.* 1972) among

others. Many of these have the advantage that the mark persists within the insect across moulting episodes. However, they are not designed to distinguish between individuals of a cohort and detection often requires specialist equipment and sophisticated techniques.

Tagging of insect larvae for its own sake, however, has received less attention and presents additional challenges. Many endopterygote larvae have a thin, translucent integument, which flexes continually and is shed repeatedly as the insect develops. While external marks can be applied to such larvae (*e.g.*, White and Singer 1987), or prolegs (where present) can be clipped (Weseloh 1985), these tags will not survive moulting.

Received 15 August 2012. Accepted 5 October 2012. First published online 27 March 2013.

C. Moffatt, School of Forensic and Investigative Sciences, University of Central Lancashire, Preston PR1 2HE, United Kingdom (e-mail: cmoffatt@uclan.ac.uk).
doi:10.4039/tce.2013.18


Currently, there appears to be no method of tagging insects that can survive moulting and allow discernment of individuals within a cohort. It was decided to attempt to address this using an injected visible implant in an endopterygote insect larva.

Visible Implant Elastomer (VIE), produced by Northwest Marine Technology (NMT) Inc. (Washington, United States of America), has been used for some time to tag marine organisms including fish (e.g., Frederick 1997), shrimp (e.g., Godin *et al.* 1996), lobster (e.g., Uglem *et al.* 1996), and crab (e.g., Davis *et al.* 2004). More recently, it has been used with squid (e.g., Replinger and Wood 2007), frogs (e.g., Nauwelaerts *et al.* 2000), salamanders (e.g., Kinkead *et al.* 2006), snakes (Hutchens *et al.* 2008), and turtles (Davy *et al.* 2010). It comprises a brightly coloured, medical-grade liquid polymer, and curing agent, which when mixed produce a harmless, gelatinous substance after around 24 hours (longer at lower temperatures). The elastomer can be injected hypodermically before curing, and remains visible through integument that is sufficiently translucent. A range of colours of VIE are available, some of which are fluorescent, facilitating observation in different light conditions with the use of an ultraviolet (black) lamp.

Butt and Lowe (2007) used VIE for marking earthworms and found that it had only a limited effect on survival, and tags were visible after many months. The method has been successfully extended to a group of terrestrial arthropods – scorpions (Chapin 2011). Based upon these successes, trials on an endopterygote insect were attempted to assess the potential use of the method with insects in general. The results of those trials are presented here.

While a “typical” endopterygote insect does not exist, blow flies (Diptera: Calliphoridae) were chosen based upon certain characteristics. Like all Cyclorrhapha, blow flies lack appendages and have three distinct and discrete larval developmental phases (instars) separated by moulting episodes (Ferrar 1987). Many calliphorid larvae feed in large intermingling aggregations – “maggot masses” – until reaching a point of satiation during the third instar phase, after which they move away from their food source (Erzinclioglu 1996), and undergo complete metamorphosis. Such an insect seemed to

Fig. 1. Rear of blow fly larva showing Visible Implant Elastomer injection locations. PVP, posterior ventral process; D11, dorsum of segment 11.




present a good test of tagging, while having the advantage of being relatively easy to culture. The tagging method specifically was required to: (i) allow a marked individual to be distinguishable in real time from otherwise ostensibly identical individuals in the maggot mass and (ii) allow the tag to be preserved across life stages, so having been retained as the insect moults.

Calliphora vicina Robineau-Desvoidy (a bluebottle fly), was used for all trials reported here, as it is common and relatively large (reaching a maximum larval size of about 18 mm (Greenberg and Kunich 2002)). Adult cultures were established from wild-caught individuals into which porcine liver was introduced as an oviposition medium. The liver was removed and kept at $\sim 22^{\circ}\text{C}$ at 55% relative humidity as the larvae hatched and began to develop.

After preliminary trials it was established that first and second instars were too small to inject successfully with VIE as the tip of the syringe needle was too large relative to the larvae. The gauge of the needle (a 29-gauge needle is supplied with the VIE) is limited by the viscosity of the elastomer. Young third instars (~ 8 mm long), however, showed more promise particularly at two injection sites (Fig. 1). The first, referred to here as the posterior ventral process (PVP), is the swollen area below the posterior spiracles on the terminal (12th) segment (abdominal segment 8). The second site was in the 11th segment, dorsally in the midline between two tissue masses (D11). These locations can only be approximate as it is difficult to maintain precise control over the tip of the syringe needle at a finer

Fig. 2. Posterior β density plots of the probability of survival for larvae tagged in one of two locations and their untagged cohabitants.



scale so more precise anatomical descriptions are superfluous.

A comparison of mortality of insects injected at the PVP and D11 sites was made using 120 feeding third instars. The larvae were randomly divided into six groups of 20. Ten of each group were injected with VIE at either PVP or D11, with each injection site represented by three of the six groups. Each group was then reared separately on 30 g of fresh porcine liver. The numbers that perished before the adult stage were significantly different for the two locations (generalised linear mixed-effects model: $z = 2.82$, $P = 0.005$); as Figure 2 shows, lower mortality resulted from injection into the D11 site. Tagging made no statistically significant difference to mortality ($z = 1.28$, $P = 0.20$), though Figure 2 suggests it may have had some effect (odds ratio = 1.7). The mortality of nontagged individuals appears to have been influenced by the mortality rate of the tagged individuals of the same cohort.


The tags were not readily visible through the cuticle of living adults, necessitating killing and dissection to establish if the tags were retained. Tags were visible in every case in the adult fly's abdomen (Fig. 3) in both male and female insects as a single, intact "blob" in almost all cases. However, in a few cases the tag had become a diffuse constellation of minute globules referred to as "scatter tags" by Butt *et al.* (2009). These were more difficult to see than an intact tag but were readily detectable under close scrutiny, again facilitated by using fluorescent tags and ultraviolet lighting.

Whether VIE injection affected rate of development was investigated using a further

Fig. 3. Green Visible Implant Elastomer tag clearly visible in a dissected adult blow fly.



Fig. 4. Posterior β density plot showing rate of development – measured by numbers reaching an arbitrary stage of pupation – was not affected by tagging.



60 larvae. The larvae were divided randomly into 12 groups of five, and larvae in six of the groups given an injection of VIE at D11. Each group was kept separately on a 30 g piece of fresh porcine liver and allowed to develop. Once a reasonable number of insects had reached an arbitrary pink-brown colour of puparium (mid-way through pupariation), this was used to simply classify insects into two developmental stages. Ignoring dead larvae (five tagged and one untagged; Fisher's exact test, $P = 0.19$) there was no difference in how these developmental groups were distributed in the tagged and untagged cohorts ($z = 0.21$, $P = 0.83$) (see Fig. 4).

Injecting the VIE was not straightforward. The small size of the larvae, and their constant

Fig. 5. Visible Implant Elastomer tagged larvae are discernible among untagged individuals. Here, three larvae have been tagged with fluorescent pink elastomer and photographed under an ultraviolet lamp.



movement made locating the syringe needle a challenge. Holding the larva between finger and thumb of one hand against a bench, with its posterior exposed was the most successful method. As the syringe needle pierces the integument, there is a release of fluid in every case. When first observed, this led to speculation that subsequent death was very likely, and it was at first surprising to discover this was not the case.

Individual tagged larvae were discernible within an untagged group, though tags are small, and it takes close inspection rather than a quick glance to pick them out (Fig. 5). However, use of a more contrasting and fluorescent coloured VIE (pink) under ultraviolet light made this somewhat easier.

Visible Implant Elastomer clearly has potential as a tag for use with insect larvae. Once injected under the integument it is visible and does not appear to affect subsequent development. It does affect mortality but not markedly, which is also what Butt and Lowe (2007) found when tagging earthworms, and like those authors it is believed that as the researcher becomes more experienced in the technique rates of mortality in the subjects would decrease. Thus, VIE injection into blow fly larvae satisfies at least partially, those requirements stipulated by Southwood and Henderson (2000). However, there is a size limitation for the subjects tagged by this method, with sizes of much less than about 8 mm likely to prove too difficult.

Acknowledgements

The author thanks Kevin Butt for introducing him to Visible Implant Elastomer and to Michael Wilson who collected some of the preliminary data.

References

- Berry, W.L., Stimman, M.W., and Wolf, W.W. 1972. Marking of native phytophagous insects with rubidium: a proposed technique. *Annals of the Entomological Society of America*, **65**: 236–238.
- Butt, K.R., Briones, M.J.I., and Lowe, C.N. 2009. Is tagging with visual implant elastomer a reliable technique for marking earthworms? *Pesquisa Agropecuária Brasileira*, **44**: 969–974.
- Butt, K.R. and Lowe, C.N. 2007. A viable technique for tagging earthworms using visible implant elastomer. *Applied Soil Ecology*, **35**: 454–457.
- Chapin, K.J. 2011. Suitability of a subcuticular permanent marking technique for scorpions. *Journal of Arachnology*, **39**: 194–196.
- Davis, J.L.D., Young-Williams, A.C., Hines, A.H., and Zmora, O. 2004. Comparing two types of internal tags in juvenile blue crabs. *Fisheries Research*, **67**: 265–274.
- Davy, C.M., Coombes, S.M., Whitear, A.K., and MacKenzie, A.S. 2010. Visible implant elastomer: a simple, non-harmful method for marking hatchling turtles. *Herpetological Review*, **41**: 442–445.
- Erzinclioglu, Z. 1996. *Blowflies naturalists handbooks number 23*. Richmond Publishing Co. Ltd, Slough, United Kingdom.
- Ferrar, P. 1987. A guide to the breeding habits and immature stages of Diptera Cyclorrhapha. *Entomograph 8*. E.J. Brill, Leiden, The Netherlands.
- Frederick, J.L. 1997. Evaluation of fluorescent elastomer injection as a method for marking small fish. *Bulletin of Marine Science*, **61**: 399–408.
- Godin, D.M., Carr, W.H., Hagino, G., Segura, F., Sweeney, J.N., and Blankenship, L. 1996. Evaluation of a fluorescent elastomer internal tag in juvenile and adult shrimp *Penaeus vannamei*. *Aquaculture*, **139**: 243–248.
- Greenberg, B. and Kunich, J.C. 2002. *Entomology and the law: flies as forensic indicators*. Cambridge University Press, Cambridge, United Kingdom.
- Hagler, J.R. 1997. Field retention of a novel mark-release-recapture method. *Environmental Entomology*, **26**: 1079–1086.
- Hagler, J.R. and Jackson, C.G. 2001. Methods for marking insects: current techniques and future prospects. *Annual Review of Entomology*, **46**: 511–543.
- Henderson, P.A. 2003. *Practical methods in ecology*. Blackwell Science, Oxford, United Kingdom.
- Hutchens, S.J., Deperno, C.S., Matthews, C.E., Pollock, K.H., and Woodward, D.K. 2008. Visible implant fluorescent elastomer: a reliable marking alternative for snakes. *Herpetological Review*, **39**: 301–303.

- Kinkead, K.E., Lanham, J.D., and Montanucci, M.R. 2006. Comparison of anesthesia and marking techniques on stress and behavioral responses in two *Desmognathus* salamanders. *Journal of Herpetology*, **40**: 323–328.
- Nauwelaerts, S., Coeck, J., and Aerts, P. 2000. Visible implant elastomers as a method for marking adult anurans. *Herpetological Review*, **31**: 154–155.
- Replinger, S.E. and Wood, J.B. 2007. A preliminary investigation of the use of subcutaneous tagging in Caribbean reef squid *Sepioteuthis sepioidea* (Cephalopoda: Loliginidae). *Fisheries Research*, **84**: 308–313.
- Service, M.W. 1993. *Mosquito ecology: field sampling methods*. Elsevier, New York, United States of America.
- Southwood, T.R.E. and Henderson, P.A. 2000. *Ecological methods*, 3rd edition. Blackwell, Oxford, United Kingdom.
- Uglen, I., Noess, H., Farestveit, E., and Jorstad, K.E. 1996. Tagging of juvenile lobsters (*Homarus gammarus* (L)) with visible implant elastomer tags. *Aquacultural Engineering*, **15**: 499–501.
- Weseloh, R.M. 1985. Dispersal, survival, and population abundance of gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae), larvae determined by releases and mark-recapture studies. *Annals of the Entomological Society of America*, **78**: 728–735.
- White, R.R. and Singer, M.C. 1987. Marking technique for larvae. *Pan-Pacific Entomologist*, **633**: 341–345.